ONELAX- bisacodyl suppository AKRON PHARMA INC

OneLAX
BISACODYL 10 mg Suppositories
Fast Acting Stimulant Laxative

Active Ingredients (per suppository)

Bisacodyl USP, 10 mg

Purpose

Laxative

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Uses

- for relief of occasional constipation
- this product usually produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only.

Do not use laxative products

- when abdominal pain, nausea or vomiting are present
- for a period longer than one week

Ask a doctor before use if you have

noticed a sudden change in bowel habits that persist over a period of 2 weeks

When using this product

it may cause abdominal discomfort, faintness, rectal burning and mild cramps

Stop use and ask a doctor if

rectal bleeding occurs, or you fail to have a bowel movement after using a laxative. This may indicate a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Directions

- adults and children 12 years of age and older
- detach one suppository from the strip
- remove from wrapper before inserting into the rectum
- the rectal suppository dose is one suppository per day or as directed by a doctor
- children under 12 years of age consult a doctor

Other information

store below 30° C (86° F)

Inactive ingredient

hydrogenated vegetable oil

Manufactured for:

Akron Pharma Inc., Fairfield, NJ-07004 www.akronpharma.com

INACTIVE INGREDIENT hydrogenated vegetable oil

Other information a store at temperatures below 30°C (86°F)

Children under 12 years and over 1 suppository once daily

The control of the suppository from the strip

at least of one suppository from the strip

at least of one suppository from the wings

at confinite tearing downward to almost the full length of the suppository (see illustration at the right)

at confinite tearing downward to almost the bull length of the suppository (see illustration at the right)

at confinite the confinite that suppository, with pointed end first, high into the rectum so it will not slip out

are retain it for 15 to 20 minutes

at confinite the confinite of the suppository seems soft, place in refrigerator for a short time before use

and the confinite that it is suppository to seems soft, place in refrigerator for a short time before use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away It pregnant or breast-feeding, ask a health professional before use.

■ you need to use a raxative tor more than one week

Stop use and ask a doctor if myou have rectal bleeding would be alial to have a bowel movement after using a laxative. These may be signs of a serious condition.

Myen using this product you may experience rectal burning mild cramps a shoominal discomfort raintness Ask a doctor before use if you have a sudden change in bowel habits that lasts over a period of 2 weeks.

Do not use when abdominal pain, nausea, or vomiting are present.

Warnings For rectal use only

Drug Facts (continued)

Bisacodyl 10mg Suppositories USGS Trelieves occasional constituation Trips product generally produces a bowel movement in 15 minutes to 1 hour

Stimulant laxative Purpose

Active ingredient (in each suppository)

Drug Facts

OneLAXTM

NDC 71399-8460-2

Bisacodyl 10mg **Suppositories**

Fast Acting Stimulant Laxative

TAMPER EVIDENT: SUPPOSITORIES ARE WRAPPED IN TAMPER EVIDENT SEALED WRAPPERS. DO NOT USE IF WRAPPER IS TORN OR OPEN WHEN PURCHASED.

12 **SUPPOSITORIES**



Akr**å**n Pharma

Manufactured for: Akron Pharma Inc., Fairfield, NJ-07004 www.akronpharma.com
Questions or comments? toll-free (877)225-6999

This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, owner of the registered trademark Dulcolax





Fast Acting Stimulant Laxative

ONELAX

bisacodyl suppository

HUMAN OTC DRUG	Item Code (Source)	NDC:71399-8460
RECTAL		

Active Ingredient/Active Moiety Ingredient Name BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y) BISACODYL BISACODYL BISACODYL 10 mg

Inactive Ingredients			
Ingredient Name	Strength		
FAT. HARD (UNII: 8334LX7S21)			

Product Characteristics			
Color	white (WHITE TO OFF WHITE)	Score	
Shape	OVAL (TORPEDO SHAPE)	Size	
Flavor		Imprint Code	
Contains			

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:71399- 8460-2	12 in 1 BOX	09/13/2024			
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product				
2	NDC:71399- 8460-5	50 in 1 BOX	09/13/2024			
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product				

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	09/13/2024	

Labeler - AKRON PHARMA INC (067878881)

Revised: 9/2024 AKRON PHARMA INC