

CLEMASTINE FUMARATE- clemastine fumarate tablet
Northwind Pharmaceuticals, LLC

Clemastine Fumarate

Description

Clemastine belongs to the benzhydryl ether group of antihistaminic compounds. The chemical name is (+)-(2R)-2-[2-[[[(R)-p-Chloro- α -methyl- α -phenylbenzyl]-oxy]ethyl]-1-methylpyrrolidine fumarate (1:1). Each tablet for oral administration contains 1.34 or 2.68 mg of clemastine fumarate.

Inactive Ingredients: Colloidal Silicon Dioxide, Corn Starch, Lactose, Povidone, Pregelatinized Starch and Stearic Acid.

Clinical Pharmacology

Clemastine fumarate is an antihistamine with anticholinergic (drying) and sedative side effects. Antihistamines appear to compete with histamine for cell receptor sites on effector cells. The inherently long duration of antihistaminic effects of clemastine fumarate has been demonstrated in wheal and flare studies. In normal human subjects who received histamine injections over a 24-hour period, the antihistaminic activity of clemastine reached a peak at 5-7 hours, persisted for 10-12 hours and, in some cases, for as long as 24 hours. Pharmacokinetic studies in man utilizing 3H and 14C labeled compound demonstrates that: clemastine is rapidly and nearly completely absorbed from the gastrointestinal tract, peak plasma concentrations are attained in 2-4 hours, and urinary excretion is the major mode of elimination.

Indications and Usage

Clemastine Fumarate Tablets 1.34 mg are indicated for the relief of symptoms associated with allergic rhinitis such as sneezing, rhinorrhea, pruritus, and lacrimation.

Clemastine Fumarate Tablets 2.68 mg are indicated for the relief of symptoms associated with allergic rhinitis such as sneezing, rhinorrhea, pruritus, and lacrimation. Clemastine Fumarate Tablets 2.68 mg are also indicated for the relief of mild, uncomplicated allergic skin manifestations of urticaria and angioedema.

It should be noted that clemastine fumarate is indicated for the dermatologic indications at the 2.68 mg dosage level only.

Contraindications

Usage in Nursing Mothers

Because of the higher risk of antihistamines for infants generally and for newborns and prematures in particular, antihistamine therapy is contraindicated in nursing mothers.

Usage in Lower Respiratory Disease

Antihistamines should not be used to treat lower respiratory tract symptoms including asthma.

Antihistamines are also contraindicated in the following conditions:

Hypersensitivity to clemastine fumarate or other antihistamines of similar chemical structure.

Monoamine oxidase inhibitor therapy (see Drug Interactions section).

Warnings

Antihistamines should be used with considerable caution in patients with: narrow angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction, symptomatic prostatic hypertrophy, and bladder neck obstruction.

Usage in Children

Safety and efficacy of clemastine fumarate have not been established in children under the age of 12 years.

Usage in Pregnancy

Experience with this drug in pregnant women is inadequate to determine whether there exists a potential for harm to the developing fetus.

Usage with CNS Depressants

Clemastine has additive effects with alcohol and other CNS depressants (hypnotics, sedatives, tranquilizers, etc.).

Usage in Activities Requiring Mental Alertness

Patients should be warned about engaging in activities requiring mental alertness such as driving a car or operating appliances, machinery, etc.

Usage in the Elderly (approximately 60 years or older)

Antihistamines are more likely to cause dizziness, sedation, and hypotension in elderly patients.

Precautions

Clemastine fumarate should be used with caution in patients with: history of bronchial asthma, increased intraocular pressure, hyperthyroidism, cardiovascular disease, and hypertension.

Drug Interactions

MAO inhibitors prolong and intensify the anticholinergic (drying) effects of antihistamines.

Adverse Reactions

Transient drowsiness, the most common adverse reaction associated with clemastine fumarate, occurs relatively frequently and may require discontinuation of therapy in some instances.

Antihistaminic Compounds

It should be noted that the following reactions have occurred with one or more antihistamines and, therefore, should be kept in mind when prescribing drugs belonging to this class, including clemastine. The most frequent adverse reactions are italicized.

General: Urticaria, drug rash, anaphylactic shock, photosensitivity, excessive perspiration, chills, dryness of the mouth, nose, and throat.

Cardiovascular System: Hypotension, headache, palpitations, tachycardia, extrasystoles.

Hematologic System: Hemolytic anemia, thrombocytopenia, agranulocytosis.

Nervous System: Sedation, sleepiness, dizziness, disturbed coordination, fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, paresthesias, blurred vision, diplopia, vertigo, tinnitus, acute labyrinthitis, hysteria, neuritis, convulsions.

GI System: Epigastric distress, anorexia, nausea, vomiting, diarrhea, constipation.

GU System: Urinary frequency, difficult urination, urinary retention, early menses.

Respiratory System: Thickening of bronchial secretions, tightness of chest and wheezing, nasal stuffiness.

Overdosage

Antihistamine overdosage reactions may vary from central nervous system depression to stimulation. Stimulation is particularly likely in children. Atropine-like signs and symptoms: dry mouth; fixed, dilated pupils; flushing; and gastrointestinal symptoms may also occur.

If vomiting has not occurred spontaneously the conscious patient should be induced to vomit. This is best done by having him drink a glass of water or milk after which he should be made to gag. Precautions against aspiration must be taken, especially in infants and children.

If vomiting is unsuccessful gastric lavage is indicated within 3 hours after ingestion and even later if large amounts of milk or cream were given beforehand. Isotonic and 1/2 isotonic saline is the lavage solution of choice.

Saline cathartics, such as milk of magnesia, by osmosis draw water into the bowel and therefore, are valuable for their action in rapid dilution of bowel content.

Stimulants should not be used.

Vasopressors may be used to treat hypotension.

Dosage and Administration

DOSAGE SHOULD BE INDIVIDUALIZED ACCORDING TO THE NEEDS AND RESPONSE OF THE PATIENT.

Clemastine Fumarate Tablets 1.34 mg

The recommended starting dose is one tablet twice daily. Dosage may be increased as required, but not to exceed six tablets daily.

Clemastine Fumarate Tablets 2.68 mg

The maximum recommended dosage is one tablet three times daily. Many patients respond favorably to a single dose which may be repeated as required, but not to exceed three tablets daily.

How supplied

Clemastine Fumarate Tablets 2.68 mg are white, round, scored tablets debossed with "93" and "308". Tablets are packaged in bottles of 100.

Store at controlled room temperature, between 20o and 25oC (68o and 77oF) (see USP).

Dispense in a tight, light-resistant container as defined in the USP/NF, with a child-resistant closure.

Label

NDC: 51655-425-04

MFG: 0093-0308-01

Clemastine Fumarate 2.68 mg

16 tablets

Rx Only

Lot#:

Exp. Date:

Each tablet contains Clemastine Fumarate, USP....2.68mg

Dosage: See package insert

Store between 68-77 degrees F.

Protect from moisture

Store in a tight, light-resistant container. Keep out of the reach of children.

Mfg. by: Teva Pharmaceuticals USA, Sellersburg, PA 18960 Lot#

Repackaged by Northwind Pharmaceuticals, Indianapolis, IN 46256



CLEMASTINE FUMARATE

clemastine fumarate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:51655-425(NDC:0093-0308)
Route of Administration	Oral		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLEMASTINE FUMARATE (UNII: 19259EGQ3D) (CLEMASTINE - UNII:95QN29S1ID)	CLEMASTINE FUMARATE	2.68 mg

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	93;308
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:51655-425-04	4 in 1 BOTTLE, DISPENSING		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA073283	04/28/2014		

Labeler - Northwind Pharmaceuticals, LLC (036986393)

Registrant - Northwind Pharmaceuticals, LLC (036986393)

Establishment

Name	Address	ID/FEI	Business Operations
Northwind Pharmaceuticals, LLC		036986393	repack(51655-425)

Revised: 7/2014

Northwind Pharmaceuticals, LLC