

CHILDRENS PAIN AND FEVER- acetaminophen suspension
Meijer Distribution Inc

Meijer Distribution, Inc. Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

72-95	11 years	15 mL
-------	----------	-------

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Principal Display Panel

meijer®

CHILDREN'S | age 2-11 years

pain & fever

acetaminophen 160 mg

per 5 mL | pain reliever

fever reducer

COMPARE TO CHILDREN'S TYLENOL® ACTIVE INGREDIENT

FAST, EFFECTIVE RELIEF OF CHILDREN'S PAIN & FEVER

GRAPE FLAVOR



Oral Suspension

Aspirin Free | Ibuprofen Free

Paraben Free | Gluten Free

Alcohol Free

4 FL OZ (118 mL)



meijer
 CHILDREN'S | age 2-11 years
pain & fever
 acetaminophen 160 mg per 5 mL
 pain reliever | fever reducer
FAST, EFFECTIVE RELIEF OF CHILDREN'S PAIN & FEVER
 GRAPE FLAVOR

Drug Facts

Active ingredient (in each 5 mL) *Purpose*
 Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses temporarily:
 ■ reduces fever
 ■ relieves minor aches and pains due to:
 ■ the common cold ■ flu ■ headache
 ■ sore throat ■ toothache

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 5 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed (see overdose warning)
 ■ shake well before using
 ■ mL = milliliter
 ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 ■ remove the child protective cap and squeeze your child's dose into the dosing cup
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information
 ■ each 5 mL contains: sodium 3 mg
 ■ store at 20-25°C (68-77°F)
 ■ do not use if printed neckband is broken or missing

Inactive ingredients
 anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-800-719-9260

**This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Tylenol®.

OUR QUALITY GUARANTEE
 The Meijer Family
 www.meijer.com/satisfaction

DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
 www.meijer.com

meijer NDC 41250-631-26
 CHILDREN'S | age 2-11 years
pain & fever
 acetaminophen 160 mg per 5 mL | pain reliever fever reducer
COMPARE TO CHILDREN'S TYLENOL® ACTIVE INGREDIENT**
FAST, EFFECTIVE RELIEF OF CHILDREN'S PAIN & FEVER
 GRAPE FLAVOR

Oral Suspension
 Aspirin Free | Ibuprofen Free
 Paraben Free | Gluten Free
 Alcohol Free

 
 4 FL OZ (118 mL)



Discard Seal, Empty & Replace Cap


 PAPER BOX PLASTIC BOTTLE PLASTIC CUP


how2recycle.info

DIRECTIONS:
STEP 1. Shake well before using

STEP 2. Remove child protective cap

STEP 3. Squeeze your child's dose into the dosing cup



IMPORTANT:
 Please see dosing chart or ask a doctor for your child's correct dose.



PID 461446

 0 41250 67394 1

13026 6E C15



CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-631
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	PURPLE (viscous)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-631-26	1 in 1 CARTON	05/05/1998	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-631-34	2 in 1 CARTON	05/05/1998	07/01/2025
2		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-631-00	1 in 1 CARTON	06/30/2025	
3		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M013	05/05/1998	

Labeler - Meijer Distribution Inc (006959555)

Revised: 7/2025

Meijer Distribution Inc