

MOTION SICKNESS RELIEF- dimenhydrinate tablet
WinCo Foods, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Winco 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to 12 years	1/2 to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to 6 years	1/2 tablet every 6-8 hours; do not exceed 1 1/2 tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions?

Call **1-800-824-1706** Monday - Friday 9am - 4pm MST

Principal Display Panel

WinCo™
FOODS

*Compare to the active ingredient in DRAMAMINE® original formula

Motion Sickness Relief

DIMENHYDRINATE 50 mg / ANTIEMETIC

12
Tablets

DISTRIBUTED BY:
WINCO FOODS, LLC, BOISE, ID 83704
50844 REV0518A19802 WF1810

*This product is not manufactured or distributed by
Medtech Products Inc., distributor of DRAMAINE® Original Formula.

The Supermarket low Price Leader®
100% SATISFACTION GUARANTEE
Comments & Questions?
800-824-1706 www.wincofoods.com

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING





B22239



Motion Sickness Relief

DIMENHYDRINATE 50 mg./ANTIEMETIC

12
Tablets

*Compare to the active ingredient in DRAMAMINE® Original Formula



100% SATISFACTION GUARANTEE

The Supermarket Low Price Leader®

DISTRIBUTED BY:
WINCO FOODS, LLC, BOISE, ID 83704
MEDTECH PRODUCTS INC., distributor of
DRAMAMINE® Original Formula.

8-2150-198-02
REV0518A19802

Comments & Questions?
800-824-1706 • www.wincofoods.com

No Print/No Varnish Area
Lot & Exp. date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient (in each tablet) Antiemetic
Dimenhydrinate 50 mg.

Uses for prevention and treatment of these symptoms associated with motion sickness:
nausea ■ vomiting ■ dizziness

Warnings
Do not use for children under 2 years of age unless directed by a doctor.
Ask a doctor before use if you have
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
■ difficulty in urination due to enlargement of the prostate gland
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
When using this product
■ marked drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity
adults and children 12 years and over
1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years
½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years
½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Drug Facts (continued)

Inactive ingredients
calcium phosphate diphosphate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Other information
■ each tablet contains: calcium 35 mg
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture
■ see end flap for expiration date and lot number

Questions? Call 1-800-824-1706 Monday - Friday
9am - 4pm MST

50844 REV0518A19802 WF1810
0 70552 23016 3

MOTION SICKNESS RELIEF

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-118
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-118-12	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part336	12/01/1992	

Labeler - WinCo Foods, LLC (056098817)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(67091-118)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867894	MANUFACTURE(67091-118)
-------------------------	--	-----------	------------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(67091-118)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(67091-118)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(67091-118)

Revised: 1/2020

WinCo Foods, LLC