MOTION SICKNESS- dimenhydrinate tablet Strategic Sourcing Services LLC

Foster and Thrive 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

 to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and	1 to 2 tablets every 4- 6 hours; do not
children	exceed 8 tablets in 24 hours, or as directed
-	by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- each tablet contains: calcium 35 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

Principal Display Panel

NDC 70677-1087-1

COMPARE TO DRAMAMINE®
ORIGINAL FORMULA ACTIVE INGREDIENT*

Foster & Thrive™

Motion Sickness

Dimenhydrinate 50 mg ANTIEMETIC

Prevents:

• Nausea • Vomiting • Dizziness for Children & Adults

ACTUAL SIZE **12 TABLETS**

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula. 50844 REV0518A19802 Rev. 10/23

Distributed by: McKesson Corp., via Strategic Sourcing Services LLC., Memphis, TN 38141 Money Back Guarantee www.fosterandthrive.com

© 2023 McKesson Corporation

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Foster & Thrive™ 🚄

Motion Sickness

Dimenhydrinate 50 mg ANTIEMETIC

Motion Sickness

NDC 70677-1087-1

Foster & Thrive™

COMPARE TO DRAMAMINE® ORIGINAL FORMULA ACTIVE INGREDIENT*

Motion Sickness

Dimenhydrinate 50 mg ANTIEMETIC

Prevents:

· Nausea · Vomiting · Dizziness for Children & Adults

ACTUAL SIZE

12 TABLETS

B-1243-198-02-R REV0518A19802

www.fosterandthrive.com **Молеу Васк Guarantee** Services LLC, Memphis, TN 38141 Distributed by: McKesson Corp., via Strategic Sourcing

50844 REV0518A19802 Rev. 10/23 Dramamine® Original Formula. Mediech Products Inc., owner of the registered trademark , I pis biognet is not manufactured or distributed by

T23 mq00:7 of ms00:9 ysbirit of ysbnoM Unestions or comments? Cal 833-358-6431

microcrystalline cellulose, silicon dioxide, stearic acid apeasic calcinm phosphate dihydrate, magnesium stearate Inactive ingredients croscamellose sodium,

see end flap for expiration date and lot number ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture

OPENED OR BLISTER IS TORN OR BROKEN ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS ■ each tablet contains: calcium 35 mg Other information

e years	% tablet every 6-8 hours; do not exceed 1% tablets in 24 hours, or as directed by a doctor
12 years	% to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
	or as directed by a doctor

12 years and over do not exceed 8 tablets in 24 hours, adults and children | 1 to 2 tablets every 4-6 hours; taken one-nail to one nour before starting activity

medical help or contact a Poison Control Center right Keep out of reach of children. In case of overdose, get

If pregnant or breast-feeding, ask a health professional

- machinery avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating
- QLOW SINESS alcohol, sedatives, and tranquitzers may increase
 - marked drowsiness may occur

When using this product sedatives or tranquilizers.

Ask a doctor or pharmacist before use if you are taking

 difficulty in unnation due to enlargement of the prostate pronchitis #giaucoma

a breathing problem such as emphysema or chronic Ask a doctor before use if you have

by a doctor.

Do not use for children under 2 years of age unless directed sвишем

s sauj zzip 🔳 Gugimov 🔳 🛚 easneu 🗖 922001916Q MILLI WOLION SICKNESS: USES for prevention and treatment of these symptoms



no print . no varnish area

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS PENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Foster and thrive 44-198

MOTION SICKNESS

dimenhydrinate tablet

D			
Prod	шст	Inform	ation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:70677-1087

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)

DIMENHYDRINATE 50 mg

Inactive Ingredients

mactive mgreatenes			
Ingredient Name	Strength		
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)			
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

Product Characteristics			
Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:70677- 1087-1	2 in 1 CARTON	05/04/2023		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information					
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
M009	05/04/2023				
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date			

Labeler - Strategic Sourcing Services LLC (116956644)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(70677-1087)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(70677-1087) , pack(70677-1087)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(70677-1087)

Revised: 4/2025 Strategic Sourcing Services LLC