

**AFTASEPTIC INSTANT PAIN RELIEF- salicylic acid 1% liquid**  
**Germa Products, LLC**

-----  
**Aftaseptic Analgesic Liquid**

**Active Ingredient**

Salicylic Acid 1% w/v

**Purpose**

Analgesic

**Uses**

For the temporary relief of pain associated with canker sores.

for the temporary relief of pain associated with canker sores.

**Warnings**

Alcohol warning: if you consume 3 or more alcoholic drinks everyday, consult with a specialist whether you should use Salicylic Acid or other pain reliever, fever reducers n Salicylic Acid may cause stomach bleeding.

Allergy alert: may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock.

**Warnings**

Flammable n Keep away from fire or flame. Do not use for teething or in children under 2 years of age

When using this product n avoid contact with eyes. Do not use longer than 1 week unless directed by a doctor. If pregnant or nursing contact a Doctor before using. Stop use and ask a doctor if condition persists or worsen

keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Other Information**

Store between 15° to 30°C (59° to 86°F). Cap bottle tightly. Do not accept if safety seal is broken or missing. Cap bottle tightly. Keep this box for future doubts.

**Directions**


Use enclosed applicator brush

Children 14 years of age to adult. Apply 2 brushstrokes 2 times a day to affected area of the mouth, gums or mucous membranes (throat). Do not rinse mouth after use. Children under 14 years of age consult a physician.

### Inactive Ingredients

Ethyl Alcohol, propylene glycol, purified water, rhubarb.

Aftaseptic



**GM**  
**GERMA**<sup>®</sup>  
NDC 59561-102-05

**AFTASEPTIC**<sup>®</sup>

**ANALGESIC LIQUID**

**INSTANT PAIN RELIEF**

Net Wt.: 0.5 FL. OZ. (15mL)

*Alivia Aftas y  
Ulceritas Bucuales*  
*Alleviates Canker Sores  
& Buccal Blisters*

Keep the bottle tightly closed

Distributed by:  
**Germa Products, LLC**  
13121 SW 122 Avenue, Miami, FL 33186  
Questions or Comments:  
1-305-278-0033, M-F, 9-5 Eastern  
[www.germa.net](http://www.germa.net)

Referred to box for "Drug Facts"  
or go to [www.germa.net](http://www.germa.net)



**GM**  
**GERMA**<sup>®</sup>  
NDC 73635-7102-5

**AFTASEPTIC**<sup>®</sup>

**ANALGESIC LIQUID**

**INSTANT PAIN RELIEF**  
*Alivio Instantaneo*

*Alivia las Aftas y Ulceritas Bucuales*



**GM**  
**GERMA**<sup>®</sup>  
*Alivia Aftas  
Dolorosas y  
Ulceritas Bucuales*  
**LIQUIDO ANALGESICO**  
Net Wt.: 0.5 FL. OZ. (14.8mL.)

**AFTASEPTIC**<sup>®</sup>



**GM**  
**GERMA**<sup>®</sup>  
*Alleviates  
Canker Sores &  
Buccal Blisters*  
**ANALGESIC LIQUID**  
Net Wt.: 0.5 FL. OZ. (14.8mL.)

**AFTASEPTIC**<sup>®</sup>

ANALGESIC LIQUID

**AFTASEPTIC**<sup>®</sup>

**GM**

**GERMA**<sup>®</sup>

**GM**

**GERMA**<sup>®</sup>

**AFTASEPTIC**<sup>®</sup>

ANALGESIC LIQUID



Distributed by: **Germa Products, LLC**  
13121 SW 122 Ave., Miami, Florida 33186  
Questions or Comments: 1-305-256-1464, M-F, 9-5 Eastern  
[www.germa.net](http://www.germa.net)



PLEASE RECYCLE



rev. 12/20

**Información Médica**  
**Propósito**  
 Ingredientes activos  
 Ácido Salicílico (USP) 1% w/v ..... Analgésico

**Usos** ■ para el alivio temporal del dolor y tratamiento de las aftas bucales (úlceras).

**Advertencia de alcohol** ■ si consume 3 o más bebidas alcohólicas por día, consulte con un especialista si debe usar ácido salicílico u otro analgésico, reductores de fiebre ■ El ácido salicílico puede causar sangramiento estomacal.

**Alerta de alergia** ■ puede causar una reacción alérgica grave que puede incluir: urticaria, hinchazón facial, asma (sibilancias), shock.

**Inflamable** ■ manténgase alejado del fuego

**No usar** ■ para la dentición ■ en niños menores de 2 años.

**Cuando use este producto** ■ evite el contacto con los ojos. ■ no use más de 1 semana a menos que lo indique un médico. ■ si esta embarazada o lactando consulte con su médico antes de usar este producto.

**Mantenga fuera del alcance de los niños** ■ en caso de ingestión accidental busque asistencia médica o llame al Centro de Tratamientos por Envenamamiento inmediatamente.

**Instrucciones** ■ use nuestro pincel aplicador para niños mayores de 14 años hasta adultos ■ usar 2 pinceladas 2 veces al día, sobre área afectada de la boca ■ no enjuagar la boca después de aplicar ■ menores de 14 años de edad consulte a un médico.

**Otras informaciones** ■ almacénelo entre 15° a 30°C (59° to 86°F). ■ no acepte si el sello de seguridad está roto o falta ■ cerrar bien después de cada uso. ■ Mantenga la caja para futura información. ■ **Ingredientes Inactivos** ■ Alcohol Etilico ■ propileno glicol ■ agua purificada ■ rubarbato

**Drug Facts**  
**Active ingredient**  
 Salicylic Acid (USP) 1% w/v ..... Analgesic

**Purpose**  
 ■ for the temporary relief of pain associated with canker sores.

**Alcohol warning** ■ if you consume 3 or more alcoholic drinks everyday, consult with a specialist whether you should use Salicylic Acid or other pain reliever, fever reducers ■ Salicylic Acid may cause stomach bleeding.

**Allergy alert** ■ may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock.

**Flammable** ■ Keep away from fire or flame.

**Do not use** ■ for teething ■ in children under 2 years of age

**When using this product** ■ avoid contact with eyes. ■ do not use longer than 1 week unless directed by a doctor. ■ if pregnant or nursing contact a doctor before using. ■ stop use and ask a doctor if condition persists or worsens

**Keep out of reach of children** ■ in case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ use enclosed applicator brush for children 14 years of age to adult ■ apply 2 brushstrokes 2 times a day to affected area of the mouth, gums or mucous membranes (throat) ■ do not rinse mouth after use ■ children under 14 years of age consult a physician.

**Other information** ■ store between 15° to 30°C (59° to 86°F). do not accept if safety seal is broken or missing ■ cap bottle tightly. ■ keep this box for future doubts. ■ **Inactive ingredients** ■ Ethyl Alcohol ■ propylene glycol ■ purified water ■ rubarbato.

Mantenga la botella bien cerrada

Keep the bottle tightly closed

# AFTASEPTIC INSTANT PAIN RELIEF

salicylic acid 1% liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73635-7102
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	10 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
ALCOHOL (UNII: 3K9958V90M)	
RHUBARB (UNII: G280W4MW6E)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73635-7102-5	1 in 1 BOX	03/29/2019	
1		14.8 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	03/29/2019	

**Labeler** - Germa Products, LLC (116626935)

Revised: 10/2023

Germa Products, LLC