

**MUSTELA MINERAL SUNSCREEN BROAD SPECTRUM SPF 50- zinc oxide,  
titanium dioxide spray**  
**Expanscience Laboratories d/b/a Mustela**

-----  
**MUSTELA MINERAL SUNSCREEN SPRAY BROAD SPECTRUM SPF 50**

**Drug Facts**

**Active ingredients**

Titanium Dioxide 4.3%

Zinc Oxide 14.7%

**Purpose**

Sunscreen

**Uses**

- Helps prevent sunburn
- If used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings**

**For external use only**

Avoid spraying in eyes. Contents under pressure. Do not puncture or incinerate. Do not store at temperature above 120F.

**Do not use**

on damaged or broken skin

**When using this product**

keep out of eyes. Rinse with water to remove

**Stop use and ask a doctor**

if rash occurs

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center immediately.

**Directions**

- Shake well before use
- Hold container 4 to 6 inches from the skin to apply
- Apply liberally 15 minutes before sun exposure
- Do not spray directly into face
- Do not spray in windy condition
- Use in well-ventilated area
- Reapply: after 80 minutes of swimming or sweating
- immediately after towel drying
- At least every 2 hours.
- **Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other protection measures including:
  - Limit time in the sun, especially from 10 a.m - 2p.m
  - Wear long-sleeved shirt, pants, hat, and sunglasses
  - Children under 6 months: ask a doctor.

### **Other information**

- Protect this product from excessive heat and direct sun.
- May stain some fabrics.

### **Inactive ingredients**

Water, Helianthus Annuus (Sunflower) Seed Oil, Olea Europaea (Olive) Fruit Oil, Butyloctyl Salicylate, Cocos Nucifera (Coconut) Oil, Capryloyl Glycerin/Sebacic Acid Copolymer, Heptyl Undecylenate, Methyl Dihydroabietate, Cetearyl Alcohol, Propanediol, Sodium Stearoyl Glutamate, Bentonite, Tocopherol, Caprylyl Glycol, Coco-Glucoside, Microcrystalline Cellulose, Bisabolol\*, Alumina, Polyhydroxystearic Acid, Citric Acid, Sodium Gluconate, Caprylhydroxamic Acid, Glycerin, Silica, Cellulose Gum, Aloe Barbadensis Leaf Juice\*, Calendula Officinalis Flower Extract\*, Chamomilla Recutita (Matricaria) Flower Extract\*, Spiraea Ulmaria Flower Extract\*, Nasturtium Officinale Flower/Leaf Extract\*, \*Certified Organic Ingredients

### **Questions or comments?**

Call toll free 1-800-422-2987  
Monday-Friday 9:00am-5:00pm EST

### **Company Information**

MADE IN THE USA from globally sourced materials for  
Expanscience Laboratories Inc.,  
Dbas Mustela, 60 E. 56th St. 6th Floor, New York, New York 10022  
[www.MustelaUSA.com](http://www.MustelaUSA.com)

### **Product Packaging**

## Drug Facts

**Active ingredients**  
Titanium Dioxide 4.3%.....Sunscreen  
Zinc Oxide 14.7%.....Sunscreen

**Purpose**  
Sunscreen

**Uses**  
■ Helps prevent sunburn. ■ If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings**  
For external use only  
Avoid spraying in eyes. Contents under pressure. Do not puncture or incinerate. Do not store at temperature above 120°F.  
Do not use on damaged or broken skin.  
When using this product keep out of eyes. Rinse with water to remove  
Stop use and ask a doctor if rash occurs  
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.

**Directions**  
■ Shake well before use ■ Hold container 4 to 6 inches from the skin to apply ■ Apply liberally 15 minutes before sun exposure ■ Do not spray directly into face ■ Do not spray in windy condition ■ Use in well-ventilated area ■ Reapply: after 80 minutes of swimming or sweating ■ Immediately after towel drying ■ At least every 2 hours  
**Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other protection measures including: ■ Limit time in the sun, especially from 10 a.m. - 2 p.m. ■ Wear long-sleeve shirts, pants, hats, and sunglasses ■ Children under 6 months: ask a doctor

**Other information**  
■ Protect this product from excessive heat and direct sun.  
■ May stain some fabrics.

**Inactive ingredients**  
Water, Helianthus Annuus (Sunflower) Seed Oil, Olea Europaea (Olive) Fruit Oil, Butyloctyl Salicylate, Cocos Nucifera (Coconut) Oil, Capryloyl Glycerin/Sebacic Acid Copolymer, Heptyl Undecylenate, Methyl Dihydroabietate, Cetearyl Alcohol, Propanediol, Sodium Stearoyl Glutamate, Bentonite, Tocopherol, Caprylyl Glycol, Coco-Glucoside, Microcrystalline Cellulose, Bisabolol\*, Alumina, Polyhydroxystearic Acid, Citric Acid, Sodium Gluconate, Caprylhydroxamic Acid, Glycerin, Silica, Cellulose Gum, Aloe Barbadensis Leaf Juice\*, Calendula Officinalis Flower Extract\*, Chamomilla Recutita (Matricaria) Flower Extract\*, Spiraea Ulmaria Flower Extract\*, Nasturtium Officinale Flower/Leaf Extract\*

**Questions or comments?**  
Call toll free 1-800-422-2987  
Monday-Friday 9:00 a.m. - 5:00 p.m. EST

\*Certified Organic Ingredients



child - family

**SPF 50**

**MINERAL SUNSCREEN SPRAY**

**BODY**

Broad spectrum SPF 50

Water resistant (80 minutes)

Ultra-sheer

Safe for sensitive skin





Net wt 170 g / 6.0 oz

MADE IN THE USA from globally sourced materials for Expanscience Laboratories Inc., Dba Mustela, 60 E. 56th St. 6th Floor, New York, New York 10022  
[www.MustelaUSA.com](http://www.MustelaUSA.com)

For sale only through select retailers. Best used before end of: see bottom



## MUSTELA MINERAL SUNSCREEN BROAD SPECTRUM SPF 50

zinc oxide, titanium dioxide spray

### Product Information

|                                |                |                           |                |
|--------------------------------|----------------|---------------------------|----------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:64768-2927 |
| <b>Route of Administration</b> | TOPICAL        |                           |                |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength      |
|---|-------------------|---------------|
| <b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)             | ZINC OXIDE        | 147 mg in 1 g |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) | TITANIUM DIOXIDE  | 43 mg in 1 g  |

| Inactive Ingredients  |                  |   |                      |                    |
|---|------------------|---|----------------------|--------------------|
| Ingredient Name   |                  |   |                      | Strength           |
| HEPTYL UNDECYLENATE (UNII: W77QUB6GXO)                                    |                  |   |                      |                    |
| METHYL DIHYDROABIETATE (UNII: 7666FJ0J9F)                                 |                  |   |                      |                    |
| CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)                                    |                  |   |                      |                    |
| BENTONITE (UNII: A3N5ZCN45C)  |                  |   |                      |                    |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U)  |                  |   |                      |                    |
| COCO GLUCOSIDE (UNII: ICS790225B)   |                  |   |                      |                    |
| ALUMINUM OXIDE (UNII: LMI26O6933)   |                  |   |                      |                    |
| SODIUM GLUCONATE (UNII: R6Q3791S76)                                       |                  |   |                      |                    |
| CAPRYLHYDROXAMIC ACID (UNII: UPY805K99W)                                  |                  |   |                      |                    |
| GLYCERIN (UNII: PDC6A3C0OX)   |                  |   |                      |                    |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)                                |                  |   |                      |                    |
| CHAMOMILE (UNII: FGL3685T2X)  |                  |   |                      |                    |
| POLYHYDROXYSTEARIC ACID (2300 MW) (UNII: YXH47AOU0F)                      |                  |   |                      |                    |
| LEVOMENOL (UNII: 24WE03BX2T)  |                  |   |                      |                    |
| NASTURTIUM OFFICINALE FLOWERING TOP (UNII: W1N2U8I64G)                    |                  |   |                      |                    |
| SUNFLOWER OIL (UNII: 3W1JG795YI)  |                  |   |                      |                    |
| FILIPENDULA ULMARIA FLOWER (UNII: 06L18L32G6)                             |                  |   |                      |                    |
| CALENDULA OFFICINALIS FLOWER (UNII: P0M7O4Y7YD)                           |                  |   |                      |                    |
| CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)        |                  |   |                      |                    |
| TOCOPHEROL (UNII: R0ZB2556P8)   |                  |   |                      |                    |
| WATER (UNII: 059QF0KO0R)  |                  |   |                      |                    |
| OLIVE OIL (UNII: 6UYK2W1W1E)  |                  |   |                      |                    |
| COCONUT OIL (UNII: Q9L0O73W7L)  |                  |   |                      |                    |
| CAPRYLOYL GLYCERIN/SEBACIC ACID COPOLYMER (2000 MPA.S) (UNII: N7YC58165T) |                  |   |                      |                    |
| BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)                                  |                  |   |                      |                    |
| ALOE VERA LEAF (UNII: ZY81Z83H0X)   |                  |   |                      |                    |
| PROPANEDIOL (UNII: 5965N8W85T)  |                  |   |                      |                    |
| SODIUM STEAROYL GLUTAMATE (UNII: 65A9F4P024)                              |                  |   |                      |                    |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)                             |                  |   |                      |                    |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)  |                  |   |                      |                    |
|   |                  |   |                      |                    |
| Packaging   |                  |   |                      |                    |
| #   | Item Code        | Package Description   | Marketing Start Date | Marketing End Date |
| 1   | NDC:64768-2927-1 | 170 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product | 05/01/2021           |                    |
|   |                  |   |                      |                    |
| Marketing Information   |                  |   |                      |                    |
| Marketing Category  |                  | Application Number or Monograph Citation                    | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug  |                  | M020  | 05/01/2021           |                    |

**Labeler** - Expanscience Laboratories d/b/a Mustela (181191057)

