

MULTI-SYMPTOM COLD- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Valu Merchandisers Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Best Choice 44-473C

Active ingredients (in each caplet)

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - nasal congestion
 - headache
 - cough
 - sinus congestion and pressure
 - sore throat
 - sneezing and runny nose
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or

followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- liver disease
- heart disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- diabetes
- a breathing problem such as emphysema or chronic bronchitis
- high blood pressure
- glaucoma
- thyroid disease

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole - do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

Best Choice®

NIGHTTIME

Multi-Symptom Cold

ACETAMINOPHEN - Pain Reliever/Fever Reducer

Chlorpheniramine maleate - Antihistamine

Dextromethorphan HBr - Cough Suppressant

Phenylephrine HCl - Nasal Decongestant

Relieves Fever/Headache,
Sore Throat, Runny Nose,
Coughing, Nasal Congestion

Pseudoephedrine Free

24 Cool Caplets

with Cool Blast Flavor

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

50844 REV0718E47308

**PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106**

Best Choice®

100% Guaranteed

www.bestchoicebrand.com

Best Choice®

NIGHTTIME Multi-Symptom Cold

Actual Size

ACETAMINOPHEN - Pain Reliever/Fever Reducer
Chlorpheniramine maleate - Antihistamine
Dextromethorphan HBr - Cough Suppressant
Phenylephrine HCl - Nasal Decongestant

Relieves Fever/Headache,
Sore Throat, Runny Nose,
Coughing, Nasal Congestion
Pseudoephedrine Free

24 Cool Caplets
with Cool Blast Flavor

Drug Facts

Active ingredients

(in each caplet)
Acetaminophen 325 mg... Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg... Antihistamine
Dextromethorphan HBr 10 mg... Cough suppressant
Phenylephrine HCl 5 mg... Nasal decongestant

Purpose

COMPLETE PRODUCT INFORMATION
KEEP OUTER PACKAGE FOR

Drug Facts (continued)

Uses ■ temporarily relieves these common cold and flu symptoms: ■ minor aches and pains

■ nasal congestion ■ headache ■ cough ■ sinus congestion and pressure ■ sore throat ■ sneezing and runny nose ■ helps clear nasal passages

Drug Facts (continued)

Questions or comments? 1-800-426-9391

No print/No varnish
Lot & Exp date

PROUDLY DISTRIBUTED BY:
VALU MERCHANDISERS, CO.
5000 KANSAS AVE
KANSAS CITY, KS 66106

Best Choice.
100% Guaranteed
www.bestchoicebrand.com

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING

50844 REV0718E47308

Drug Facts (continued)

Other information ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR
BLISTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients ■ corn starch, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

Do not use

■ if you have ever had an allergic reaction to this product or any of its ingredients
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if you have
■ cough that occurs with too much phlegm (mucus) ■ difficulty in urination due to enlargement of the prostate gland ■ liver disease ■ heart disease ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ diabetes

Directions

■ do not take more than directed
■ adults and children 12 years and over
■ take 2 caplets every 4 hours
■ swallow whole - do not crush, chew, or dissolve
■ do not take more than 10 caplets in 24 hours
■ children under 12 years: ask a doctor

more than 7 days ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

B-2115-4730-08CMR
REV0718E47308

BEST CHOICE





Best Choice 44-473C

MULTI-SYMPTOM COLD

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-473
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

STEARIC ACID (UNII: 4ELV7Z65AP)

SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)

Product Characteristics

Color	BLUE	Score	no score
Shape	OVAL	Size	17mm
Flavor	MINT (Cool Blast)	Imprint Code	44;473
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-473-08	2 in 1 CARTON	07/21/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	07/21/2005	

Labeler - Valu Merchandisers Company (868703513)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(63941-473)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(63941-473)

Revised: 10/2019

Valu Merchandisers Company