

**QUITA CALLOS CORN AND CALLUS REMOVER- salicylic acid 17% liquid**  
**Germa Products, LLC**

-----  
**Quita Callos Analgesic Liquid**

**Active Ingredient**

Salicylic Acid 17%

**Uses**

for the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance of the surface.

for the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance of the surface.

**Purpose**

Wart Remover Liquid

**Warnings**

For external use only. Flammable: Keep away from fire or flame.

Do not use: in or near eyes or mucous membranes. On irritated skin. If prone to allergic reaction to any ingredient in this product. On moles, birthmarks, warts with hair growing from them, genital warts, or warts on the face or mucous membranes. On any area that is infected or reddened. If you are a diabetic, or have poor blood circulation n on large area of the body, on wounds or damaged skin.

When using this product: If product gets in eyes, flush with water for 15 minutes. Do not inhale vapors. Cap bottle tightly and store at room temperature away from heat. If discomfort persists, see your doctor.

Stop use and ask a doctor if; Irritation occurs n condition worsens or does not improve or discomfort persists. If pregnant or breast-feeding. Ask a health professional before use. Keep out of reach of children. In case of accidental ingestion, get medical help or contact a Poison Control Center immediately.

**Warnings**

- Keep out of reach of children. In case of accidental ingestion, get medical help or contact a Poison Control Center immediately.

**Other Information**

Store between 20° to 25°C (68° to 77°F). Cap bottle tightly. Use product in well ventilated area. Avoid inhaling vapors. Retain packaging for full labeling.

## Directions

- Wash affected area and dry thoroughly.
- Apply one drop at a time with applicator to sufficiently cover each wart.
- Let dry.
- Repeat this procedure once or twice daily as needed for up to 2 weeks.

## Inactive Ingredients

Flexible Collodion (Alcohol, Camphor, Mineral Oil), and Isopropyl Alcohol.

INDICATIONS: For the removal of corns and calluses. Relieves pain by removing corns and calluses.

DIRECTIONS: Wash the affected area and dry thoroughly. Brush on corn or callus only, not on healthy skin. Let dry. Repeat this procedure once or twice daily as needed for up to 14 days (until corn/ callus is removed). You may also soak corn/callus in warm water for 5 minutes to assist in removal.

*Keep the bottle tightly closed*

**GERMA<sup>®</sup>**  
**QuitaCallos<sup>™</sup>**  
Removedor de Callos  
Corn & Callus Remover

**Salicylic Acid 17%**

Net Wt.:  
0.5 FL. OZ. (14.8mL.)

WARNINGS: For external use only. Do not use this product on irritated skin, or in any area that is infected or reddened, if you are a diabetic, or if you have poor blood circulation. If discomfort persists, see your doctor or a podiatrist. **Flammable.** Keep away from fire or flame. Cap bottle tightly and store at room temperature away from heat. If product gets into the eye, flush with water for 15 minutes. Avoid inhaling vapors. Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a poison control center immediately.

ACTIVE INGREDIENT: Salicylic Acid 17% (w/w).  
INACTIVE INGREDIENTS: Flexible Collodion, Isopropyl Alcohol.

Distributed by:  
**Germa Products, LLC**  
13121 SW 122 Avenue, Miami, Florida 33186  
Questions or Comments: 1-305-278-0030, M-F 9-5 Eastern  
[www.germa.net](http://www.germa.net)

GM NDC 59561-712-05

rev. 12/20

**GM** NDC 59561-712-05

**GERMA<sup>®</sup>**  
**QuitaCallos<sup>™</sup>**  
Removedor de Callos  
Corn & Callus Remover

**Salicylic Acid 17%**

**GM** **GERMA<sup>®</sup>**  
**QuitaCallos<sup>™</sup>**  
Removedor de Callos  
Corn & Callus Remover

**Salicylic Acid 17%**

Net Wt.:  
0.5 FL. OZ. (14.8mL.)

**GM** **GERMA<sup>®</sup>**  
**QuitaCallos<sup>™</sup>**  
Removedor de Callos  
Corn & Callus Remover

**Salicylic Acid 17%**

Net Wt.:  
0.5 FL. OZ. (14.8mL.)

**GM** **GERMA<sup>®</sup>**  
**QuitaCallos<sup>™</sup>**  
Removedor de Callos  
Corn & Callus Remover



Distributed by: **Germa Products, LLC**  
 13121 SW 122 Avenue, Miami, Florida 33186  
 Questions or Comments: 1-305-278-0033, M-F, 9-5 Eastern  
**www.germa.net**



rev. 12/20

**Drug Facts**

**Active ingredient**  
 Salicylic acid 17% w/w..... Corn & Callus Removal

**Purpose**

**Uses**

- for the removal of corns and calluses.
- relieves pain by removing corns and calluses.

**Warnings**

- For external use only. ■ Flammable: Keep away from fire or flame.
- Do not use
- if you are a diabetic. ■ if you have poor blood circulation.
- on irritated skin or any area that is irritated or reddened.

**When using this product**

- if product gets in eyes, flush with water for 15 minutes ■ do not inhale vapors. ■ cap bottle tightly and store at room temperature away from heat.
- if discomfort persists, see your doctor or podiatrist.

**Stop use and ask a doctor if discomfort persists.**

**Keep out of reach of children.** In case of accidental ingestion, get medical help or contact a Poison Control Center immediately.

**Directions**

- wash the affected area and dry thoroughly.
- apply one drop at a time with applicator to sufficiently cover each corn/callus. ■ let dry. ■ repeat this procedure once or twice daily as needed for up to 14 days (until corn/callus is removed). ■ may soak corn/callus in warm water for 5 minutes to assist in removal.

**Other information**

- store between 20° to 25°C (68° to 77°F). ■ cap bottle tightly. ■ use product in well-ventilated area. ■ avoid inhaling vapors. retain packaging for full labeling.
- protective pads may be used to cushion corn/callus during treatment. ■ Additional cushions may be purchased separately.

**Inactive ingredients** ■ flexible collodion ■ isopropyl alcohol



**Keep the bottle tightly closed**

## Quita Callos

### QUITA CALLOS CORN AND CALLUS REMOVER

salicylic acid 17% liquid

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:73635-5811
<b>Route of Administration</b>	TOPICAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	170 mg in 1 mL

#### Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	

<b>CAMPHOR (SYNTHETIC)</b> (UNII: 5TJD82A1ET)	
<b>CASTOR OIL</b> (UNII: D5340Y2I9G)	

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73635-5811-5	1 in 1 BOX	03/29/2019	
1		14.8 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M030	03/29/2019	

**Labeler** - Germa Products, LLC (116626935)

Revised: 10/2025

Germa Products, LLC