

FENTANYL CITRATE- fentanyl citrate injection, solution

Hospira, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use FENTANYL CITRATE INJECTION safely and effectively. See full prescribing information for FENTANYL CITRATE INJECTION.

FENTANYL CITRATE Injection, for Intravenous or Intramuscular use, CII

Initial U.S. Approval: 1968

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; CYTOCHROME P450 3A4 INTERACTION; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

See full prescribing information for complete boxed warning.

- Fentanyl Citrate Injection exposes users to risks of addiction, abuse, and misuse, which can lead to overdose and death. Assess patient's risk before prescribing and monitor regularly for these behaviors and conditions. (5.1)
- Serious, life-threatening, or fatal respiratory depression may occur. Monitor closely, especially upon initiation or following a dose increase. (5.2)
- Concomitant use with CYP3A4 inhibitors (or discontinuation of CYP3A4 inducers) can result in a fatal overdose of fentanyl. (5.3, 7, 12.3).
- Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing for use in patients for whom alternative treatment options are inadequate; limit dosages and durations to the minimum required; and follow patients for signs and symptoms of respiratory depression and sedation. (5.4, 7)

RECENT MAJOR CHANGES

Warnings and Precautions (5.2)

10/2019

INDICATIONS AND USAGE

Fentanyl Citrate Injection is an opioid agonist indicated for: (1)

- analgesic action of short duration during the anesthetic periods, premedication, induction and maintenance, and in the immediate postoperative period (recovery room) as the need arises.
- use as an opioid analgesic supplement in general or regional anesthesia.
- administration with a neuroleptic as an anesthetic premedication, for the induction of anesthesia and as an adjunct in the maintenance of general and regional anesthesia.
- use as an anesthetic agent with oxygen in selected high risk patients, such as those undergoing open heart surgery or certain complicated neurological or orthopedic procedures.

DOSAGE AND ADMINISTRATION

- Fentanyl Citrate Injection should be administered only by persons specifically trained in the use of intravenous anesthetics and management of the respiratory effects of potent opioids.
- Ensure that an opioid antagonist, resuscitative and intubation equipment, and oxygen are readily available. (2.1)
- Individualize dosing based on the factors such as age, body weight, physical status, underlying pathological condition, use of other drugs, type of anesthesia to be used, and the surgical procedure involved. (2.1)
- Initiate treatment in adults with 50 to 100 mcg (0.05 to 0.1 mg) (1 to 2 mL). (2.2)
- Initiate treatment in children 2 to 12 years of age with a reduced dose as low as 2 to 3 mcg/kg. (2.2)

DOSAGE FORMS AND STRENGTHS

- Injection: 50 mcg (0.05 mg)/mL in single-dose Flitop vial. (3)
- Injection: 50 mcg (0.05 mg)/mL in single-dose ampule. (3)

CONTRAINDICATIONS

- Hypersensitivity to fentanyl. (4)

WARNINGS AND PRECAUTIONS

- Risks of Skeletal Muscle Rigidity and Skeletal Muscle Movement: Manage with neuromuscular blocking agent. See full prescribing information for more detail on managing these risks. (5.5)
- Severe Cardiovascular Depression: Monitor during dosage initiation and titration. (5.6)
- Serotonin Syndrome: Potentially life-threatening condition could result from concomitant serotonergic drug administration. Discontinue Fentanyl Citrate Injection if serotonin syndrome is suspected. (5.7)
- Adrenal Insufficiency: If diagnosed, treat with physiologic replacement of corticosteroids, and wean patient off of the opioid. (5.8)
- Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness: Monitor for sedation and respiratory depression. (5.9)

ADVERSE REACTIONS

Most common serious adverse reactions were respiratory depression, apnea, rigidity, and bradycardia. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Hospira, Inc. at 1-800-441-4100, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Concomitant Use of CNS Depressants: May decrease pulmonary arterial pressure and may cause hypotension. See FPI for management instructions. For post-operative pain, start with the lowest effective dosage and monitor for potentiation of CNS depressant effects. (5.4, 7)
- Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics: Avoid use with Fentanyl Citrate Injection because they may reduce analgesic effect of Fentanyl Citrate Injection or precipitate withdrawal symptoms. (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: May cause fetal harm. (8.1)
- Lactation: Infants exposed to Fentanyl Citrate Injection through breast milk should be monitored for excess sedation and respiratory depression. (8.2)
- Geriatric Patients: Titrate slowly and monitor for CNS and respiratory depression. (8.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 10/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; CYTOCHROME P450 3A4 INTERACTION; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosage and Administration Instructions
- 2.2 Dosage

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Addiction, Abuse, and Misuse
- 5.2 Life-Threatening Respiratory Depression
- 5.3 Risks of Concomitant Use or Discontinuation of Cytochrome P450 3A4 Inhibitors and Inducers
- 5.4 Risks from Concomitant Use with Benzodiazepines or Other Central Nervous System Depressants
- 5.5 Risks of Muscle Rigidity and Skeletal Muscle Movement
- 5.6 Severe Cardiovascular Depression
- 5.7 Serotonin Syndrome with Concomitant Use of Serotonergic Drugs
- 5.8 Adrenal Insufficiency
- 5.9 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, or Head Injury
- 5.10 Risks of Use in Patients with Gastrointestinal Conditions

5.11 Increased Risk of Seizures in Patients with Seizure Disorders

5.12 Risks of Driving and Operating Machinery

5.13 Risks due to Interaction with Neuroleptic Agents

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

9.2 Abuse

9.3 Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; CYTOCHROME P450 3A4 INTERACTION; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

Fentanyl Citrate Injection exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing Fentanyl Citrate Injection, and monitor all patients regularly for the development of these behaviors and conditions [see *Warnings and Precautions (5.1)*].

Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of Fentanyl Citrate Injection. Monitor for respiratory depression, especially during initiation of Fentanyl Citrate Injection or following a dose increase [see *Warnings and Precautions (5.2)*].

Cytochrome P450 3A4 Interaction

The concomitant use of Fentanyl Citrate Injection with all cytochrome P450 3A4 inhibitors may result in an increase in fentanyl plasma concentrations, which could increase or prolong adverse reactions and may cause potentially fatal respiratory depression. In addition, discontinuation of a concomitantly used cytochrome P450 3A4 inducer may result in an increase in fentanyl plasma concentration. Monitor patients receiving Fentanyl Citrate Injection and any CYP3A4 inhibitor or inducer [see *Warnings and Precautions (5.3)*, *Drug Interactions (7)*, *Clinical Pharmacology (12.3)*].

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death [see *Warnings and Precautions (5.4)*, *Drug Interactions (7)*].

- Reserve concomitant prescribing of Fentanyl Citrate Injection and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

1 INDICATIONS AND USAGE

Fentanyl Citrate Injection is indicated for:

- analgesic action of short duration during the anesthetic periods, premedication, induction and maintenance, and in the immediate postoperative period (recovery room) as the need arises.
- use as an opioid analgesic supplement in general or regional anesthesia.
- administration with a neuroleptic as an anesthetic premedication, for the induction of anesthesia and as an adjunct in the maintenance of general and regional anesthesia.
- use as an anesthetic agent with oxygen in selected high risk patients, such as those undergoing open heart surgery or certain complicated neurological or orthopedic procedures.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Instructions

Fentanyl Citrate Injection should be administered only by persons specifically trained in the use of intravenous anesthetics and management of the respiratory effects of potent opioids.

- Ensure that an opioid antagonist, resuscitative and intubation equipment, and oxygen are readily available.
- Individualize dosage based on factors such as age, body weight, physical status, underlying pathological condition, use of other drugs, type of anesthesia to be used, and the surgical procedure involved.
- Monitor vital signs routinely.

As with other potent opioids, the respiratory depressant effect of fentanyl may persist longer than the measured analgesic effect. The total dose of all opioid agonists administered should be considered by the practitioner before ordering opioid analgesics during recovery from anesthesia.

If Fentanyl Citrate Injection is administered with a CNS depressant, become familiar with the properties of each drug, particularly each product's duration of action. In addition, when such a combination is used, fluids and other countermeasures to manage hypotension should be available *[see Warnings and Precautions (5.4)]*.

Inspect parenteral drug products visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

2.2 Dosage

50 mcg = 0.05 mg = 1 mL

Premedication in Adults

50 to 100 mcg (0.05 to 0.1 mg) (1 to 2 mL) may be administered intramuscularly 30 to 60 minutes prior to surgery.

Adjunct to General Anesthesia

See Dosage Range Charts below.

Table 1: Dosage Range Chart

TOTAL DOSAGE (expressed as fentanyl base)		
Low Dose – 2 mcg/kg (0.002 mg/kg) (0.04 mL/kg). For use in minor, but painful, surgical procedures. May also provide some pain relief in the immediate postoperative period.	Moderate Dose – 2–20 mcg/kg (0.002–0.02 mg/kg) (0.04–0.4 mL/kg). For use in major surgical procedures, in addition to adequate analgesia, may abolish some of the stress response. Expect respiratory depression requiring artificial ventilation during anesthesia and careful observation of ventilation postoperatively is essential.	High Dose – 20–50 mcg/kg (0.02–0.05 mg/kg) (0.4–1 mL/kg). For open heart surgery and certain more complicated neurosurgical and orthopedic procedures where surgery is more prolonged, and the stress response to surgery would be detrimental to the wellbeing of the patient. In conjunction with nitrous oxide/oxygen has been shown to attenuate the stress response as defined by increased levels of circulating growth hormone, catecholamine, ADH and prolactin. Expect the need for postoperative ventilation and observation are essential due to extended postoperative respiratory depression.

MAINTENANCE DOSE (expressed as fentanyl base)		
Low Dose – 2 mcg/kg (0.002 mg/kg) (0.04 mL/kg). Additional dosages are infrequently needed in these minor procedures.	Moderate Dose – 2–20 mcg/kg (0.002–0.02 mg/kg) (0.04–0.4 mL/kg). 25 to 100 mcg (0.025 to 0.1 mg) (0.5 to 2 mL) Administer intravenously or intramuscularly as needed when movement and/or changes in vital signs indicate surgical stress or lightening of analgesia.	High Dose – 20–50 mcg/kg (0.02–0.05 mg/kg) (0.4–1 mL/kg). Maintenance dosage [ranging from 25 mcg (0.025 mg) (0.5 mL) to one half the initial loading dose] as needed based on vital signs changes indicative of stress and lightening of analgesia. Individualize dosage especially if the anticipated remaining operative time is short.

Adjunct to Regional Anesthesia

50 to 100 mcg (0.05 to 0.1 mg) (1 to 2 mL) may be administered intramuscularly or slowly intravenously, over one to two minutes, when additional analgesia is required.

Postoperatively (recovery room)

50 to 100 mcg (0.05 to 0.1 mg) (1 to 2 mL) may be administered intramuscularly for the control of pain, tachypnea and emergence delirium. The dose may be repeated in one to two hours as needed.

For Induction and Maintenance in Children 2 to 12 Years of Age

A reduced dose as low as 2 to 3 mcg/kg is recommended.

As a General Anesthetic

As a technique to attenuate the responses to surgical stress without the use of additional anesthetic agents, doses of 50 to 100 mcg/kg (0.05 to 0.1 mg/kg) (1 to 2 mL/kg) may be administered with oxygen and a muscle relaxant. In certain cases, doses up to 150 mcg/kg (0.15 mg/kg) (3 mL/kg) may be necessary to produce this anesthetic effect. It has been used for open heart surgery and certain other major surgical procedures in patients for whom protection of the myocardium from excess oxygen demand is particularly indicated, and for certain complicated neurological and orthopedic procedures.

3 DOSAGE FORMS AND STRENGTHS

- Injection: 50 mcg (0.05 mg)/mL in single-dose Flitop vial.
- Injection: 50 mcg (0.05 mg)/mL in single-dose ampule.

4 CONTRAINDICATIONS

Fentanyl Citrate Injection is contraindicated in patients with:

- Hypersensitivity to fentanyl (e.g., anaphylaxis) [see *Adverse Reactions* (6)].

5 WARNINGS AND PRECAUTIONS

5.1 Addiction, Abuse, and Misuse

Fentanyl Citrate Injection contains fentanyl, a Schedule CII controlled substance. As an opioid, Fentanyl Citrate Injection exposes users to the risks of addiction, abuse, and misuse [see *Drug Abuse and Dependence* (9)].

Opioids are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Consider these risks when handling Fentanyl Citrate Injection. Strategies to reduce these risks include proper product storage and control practices for a C-II drug. Contact local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

5.2 Life-Threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Adequate facilities should be available for postoperative monitoring and ventilation of patients administered anesthetic doses of Fentanyl Citrate Injection. It is essential that these facilities be fully equipped to handle all degrees of respiratory depression. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status [see *Overdosage (10)*]. Carbon dioxide (CO₂) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

To reduce the risk of respiratory depression, proper dosing and titration of Fentanyl Citrate Injection are essential [see *Dosage and Administration (2)*]. As with other potent opioids, the respiratory depressant effect of fentanyl may persist longer than the measured analgesic effect. The total dose of all opioid agonists administered should be considered by the practitioner before ordering opioid analgesics during recovery from anesthesia.

Certain forms of conduction anesthesia, such as spinal anesthesia and some peridural anesthetics, can alter respiration by blocking intercostal nerves. Through other mechanisms [see *Clinical Pharmacology (12.2)*] Fentanyl Citrate Injection can also alter respiration. Therefore, when Fentanyl Citrate Injection is used to supplement these forms of anesthesia, the anesthetist should be familiar with the physiological alterations involved, and be prepared to manage them in the patients selected for these forms of anesthesia.

Patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased respiratory drive including apnea, even at recommended dosages of Fentanyl Citrate Injection. Elderly, cachectic, or debilitated patients may have altered pharmacokinetics or altered clearance compared to younger, healthier patients resulting in greater risk for respiratory depression.

Monitor such patients closely including vital signs, particularly when initiating and titrating Fentanyl Citrate Injection and when Fentanyl Citrate Injection is given concomitantly with other drugs that depress respiration. To reduce the risk of respiratory depression, proper dosing and titration of Fentanyl Citrate Injection are essential [see *Dosage and Administration (2)*].

Opioids can cause sleep-related breathing disorders including central sleep apnea (CSA) and sleep-related hypoxemia. Opioid use increases the risk of CSA in a dose-dependent fashion. In patients who present with CSA, consider decreasing the opioid dosage using best practices for opioid taper [see *Dosage and Administration (2.1)*].

5.3 Risks of Concomitant Use or Discontinuation of Cytochrome P450 3A4 Inhibitors and Inducers

Concomitant use of Fentanyl Citrate Injection with a CYP3A4 inhibitor, such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g., ketoconazole), and protease inhibitors (e.g., ritonavir), may increase plasma concentrations of fentanyl and prolong opioid adverse reactions, which may exacerbate respiratory depression [see *Warnings and Precautions (5.2)*], particularly when an inhibitor is added after a stable dose of Fentanyl Citrate Injection is achieved. Similarly, discontinuation of a CYP3A4 inducer, such as rifampin, carbamazepine, and phenytoin, in Fentanyl Citrate Injection-

treated patients may increase fentanyl plasma concentrations and prolong opioid adverse reactions. When using Fentanyl Citrate Injection with CYP3A4 inhibitors or discontinuing CYP3A4 inducers in Fentanyl Citrate Injection-treated patients, monitor patients closely at frequent intervals and consider dosage reduction of Fentanyl Citrate Injection [see *Dosage and Administration (2.1)*, *Drug Interactions (7)*].

Concomitant use of Fentanyl Citrate Injection with CYP3A4 inducers, or discontinuation of a CYP3A4 inhibitor, could result in lower than expected fentanyl plasma concentrations and decrease efficacy. When using Fentanyl Citrate Injection with CYP3A4 inducers or discontinuation of a CYP3A4 inhibitor, monitor patients closely at frequent intervals and consider increasing the fentanyl dosage. [see *Dosage and Administration (2.1)*, *Drug Interactions (7)*].

5.4 Risks from Concomitant Use with Benzodiazepines or Other Central Nervous System Depressants

When benzodiazepines or other CNS depressants are used with Fentanyl Citrate Injection, pulmonary arterial pressure may be decreased. This fact should be considered by those who conduct diagnostic and surgical procedures where interpretation of pulmonary arterial pressure measurements might determine final management of the patient. When high dose or anesthetic dosages of Fentanyl Citrate Injection are employed, even relatively small dosages of diazepam may cause cardiovascular depression.

When Fentanyl Citrate Injection is used with CNS depressants, hypotension can occur. If it occurs, consider the possibility of hypovolemia and manage with appropriate parenteral fluid therapy. When operative conditions permit, consider repositioning the patient to improve venous return to the heart. Exercise care in moving and repositioning of patients because of the possibility of orthostatic hypotension. If volume expansion with fluids plus other countermeasures do not correct hypotension, consider administration of pressor agents other than epinephrine. Epinephrine may paradoxically decrease blood pressure in patients treated with a neuroleptic that blocks alpha adrenergic activity.

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of Fentanyl Citrate Injection with benzodiazepines or other CNS depressants (e.g., non-benzodiazepine sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol).

If the decision is made to manage postoperative pain with Fentanyl Citrate Injection concomitantly with a benzodiazepine or other CNS depressant, start dosing with the lowest effective dosage and titrate based on clinical response. Follow patients closely for signs and symptoms of respiratory depression, sedation, and hypotension. Fluids or other measures to counter hypotension should be available. [see *Drug Interactions (7)*].

5.5 Risks of Muscle Rigidity and Skeletal Muscle Movement

Fentanyl Citrate Injection may cause muscle rigidity, particularly involving the muscles of respiration. The incidence and severity of muscle rigidity is dose related. These effects are related to the dose and speed of injection. Skeletal muscle rigidity also has been reported to occur or recur infrequently in the extended postoperative period usually following high dose administration. In addition, skeletal muscle movements of various groups in the extremities, neck, and external eye have been reported during induction of anesthesia with Fentanyl Citrate Injection; these reported movements have, on rare occasions, been strong enough to pose patient management problems.

These effects are related to the dose and speed of injection and its incidence can be reduced by: 1) administration of up to 1/4 of the full paralyzing dose of a non-depolarizing neuromuscular blocking agent just prior to administration of Fentanyl Citrate Injection; 2) administration of a full paralyzing dose of a neuromuscular blocking agent following loss of eyelash reflex when Fentanyl Citrate Injection is used in anesthetic doses titrated by slow intravenous infusion; or, 3) simultaneous administration of Fentanyl Citrate Injection and a full paralyzing dose of a neuromuscular blocking agent when Fentanyl

Citrate Injection is used in rapidly administered anesthetic dosages. The neuromuscular blocking agent used should be compatible with the patient's cardiovascular status.

5.6 Severe Cardiovascular Depression

Fentanyl Citrate Injection may cause severe bradycardia, severe hypotension including orthostatic hypotension, and syncope. There is increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (e.g., phenothiazines or general anesthetics) [see *Drug Interactions (7)*]. In patients with circulatory shock, Fentanyl Citrate Injection may cause vasodilation that can further reduce cardiac output and blood pressure. Monitor these patients for signs of hypotension after initiating or titrating the dosage of Fentanyl Citrate Injection.

5.7 Serotonin Syndrome with Concomitant Use of Serotonergic Drugs

Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of fentanyl with serotonergic drugs. Serotonergic drugs include selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT₃ receptor antagonists, drugs that affect the serotonergic neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), certain muscle relaxants (i.e., cyclobenzaprine, metaxalone), and drugs that impair metabolism of serotonin (including MAO inhibitors, both those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue) [see *Drug Interactions (7)*]. This may occur within the recommended dosage range.

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination, rigidity), and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). The onset of symptoms generally occurs within several hours to a few days of concomitant use, but may occur later than that. Discontinue Fentanyl Citrate Injection if serotonin syndrome is suspected.

5.8 Adrenal Insufficiency

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

5.9 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, or Head Injury

In patients who may be susceptible to the intracranial effects of CO₂ retention (e.g., those with evidence of increased intracranial pressure or brain tumors), Fentanyl Citrate Injection may reduce respiratory drive, and the resultant CO₂ retention can further increase intracranial pressure. Monitor such patients for signs of increasing intracranial pressure.

5.10 Risks of Use in Patients with Gastrointestinal Conditions

Fentanyl may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Monitor patients with biliary tract disease, including acute pancreatitis for worsening symptoms.

5.11 Increased Risk of Seizures in Patients with Seizure Disorders

Fentanyl may increase the frequency of seizures in patients with seizure disorders, and may increase the

risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during Fentanyl Citrate Injection therapy.

5.12 Risks of Driving and Operating Machinery

Fentanyl may impair the mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery after Fentanyl Citrate Injection administration.

5.13 Risks due to Interaction with Neuroleptic Agents

Many neuroleptic agents have been associated with QT prolongation, torsades de pointes, and cardiac arrest. Administer neuroleptic agents with extreme caution in the presence of risk factors for development of prolonged QT syndrome and torsades de pointes, such as: 1) clinically significant bradycardia (less than 50 bpm), 2) any clinically significant cardiac disease, including baseline prolonged QT interval, 3) treatment with Class 1 and Class III antiarrhythmics, 4) treatment with monoamine oxidase inhibitors (MAOI's), 5) concomitant treatment with other drug products known to prolong the QT interval and 6) electrolyte imbalance, in particular hypokalemia and hypomagnesemia, or concomitant treatment with drugs (e.g. diuretics) that may cause electrolyte imbalance.

Elevated blood pressure, with and without pre-existing hypertension, has been reported following administration of Fentanyl Citrate Injection combined with a neuroleptic. This might be due to unexplained alterations in sympathetic activity following large doses; however, it is also frequently attributed to anesthetic and surgical stimulation during light anesthesia.

ECG monitoring is indicated when a neuroleptic agent is used in conjunction with Fentanyl Citrate Injection as an anesthetic premedication, for the induction of anesthesia, or as an adjunct in the maintenance of general or regional anesthesia.

When Fentanyl Citrate Injection is used with a neuroleptic and an EEG is used for postoperative monitoring, the EEG pattern may return to normal slowly.

6 ADVERSE REACTIONS

The following serious adverse reactions are described, or described in greater detail, in other sections:

- Addiction, Abuse, and Misuse [*see Warnings and Precautions (5.1)*]
- Life-Threatening Respiratory Depression [*see Warnings and Precautions (5.2)*]
- Interactions with Benzodiazepines or Other CNS Depressants [*see Warnings and Precautions (5.4)*]
- Serotonin Syndrome [*see Warnings and Precautions (5.7)*]
- Severe Cardiovascular Depression [*see Warnings and Precautions (5.6)*]
- Gastrointestinal Adverse Reactions [*see Warnings and Precautions (5.10)*]
- Seizures [*see Warnings and Precautions (5.11)*]

The following adverse reactions associated with the use of fentanyl were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

As with other opioid agonists, the most common serious adverse reactions reported to occur with fentanyl are respiratory depression, apnea, rigidity, and bradycardia; if these remain untreated, respiratory arrest, circulatory depression or cardiac arrest could occur. Other adverse reactions that have been reported are hypertension, hypotension, dizziness, blurred vision, nausea, emesis, diaphoresis, pruritus, urticarial, laryngospasm, and anaphylaxis.

It has been reported that secondary rebound respiratory depression may occasionally occur postoperatively.

When a tranquilizer is used with fentanyl, the following adverse reactions can occur: chills and/or shivering, restlessness, and postoperative hallucinatory episodes (sometimes associated with transient periods of mental depression); extrapyramidal symptoms (dystonia, akathisia, and oculogyric crisis) have been observed up to 24 hours postoperatively. When they occur, extrapyramidal symptoms can usually be controlled with anti-parkinson agents. Postoperative drowsiness is also frequently reported following the use of neuroleptics with Fentanyl Citrate Injection.

Cases of cardiac dysrhythmias, cardiac arrest, and death have been reported following the use of Fentanyl Citrate Injection with a neuroleptic agent.

Serotonin syndrome: Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs.

Adrenal insufficiency: Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use.

Anaphylaxis: Anaphylaxis has been reported with ingredients contained in Fentanyl Citrate Injection.

Androgen deficiency: Cases of androgen deficiency have occurred with chronic use of opioids [see *Clinical Pharmacology* (12.2)].

7 DRUG INTERACTIONS

Table 2 includes clinically significant drug interactions with Fentanyl Citrate Injection.

Table 2: Clinically Significant Drug Interactions with Fentanyl Citrate Injection

Inhibitors of CYP3A4	
<i>Clinical Impact:</i>	The concomitant use of Fentanyl Citrate Injection and CYP3A4 inhibitors can increase the plasma concentration of fentanyl, resulting in increased or prolonged opioid effects, particularly when an inhibitor is added after a stable dose of Fentanyl Citrate Injection is achieved [see <i>Warnings and Precautions</i> (5.3)]. After stopping a CYP3A4 inhibitor, as the effects of the inhibitor decline, the fentanyl plasma concentration will decrease [see <i>Clinical Pharmacology</i> (12.3)], resulting in decreased opioid efficacy or a withdrawal syndrome in patients who had developed physical dependence to fentanyl.
<i>Intervention:</i>	If concomitant use is necessary, consider dosage reduction of Fentanyl Citrate Injection until stable drug effects are achieved [see <i>Dosage and Administration</i> (2)]. Monitor patients for respiratory depression and sedation at frequent intervals. If a CYP3A4 inhibitor is discontinued, consider increasing the Fentanyl Citrate Injection dosage until stable drug effects are achieved. Monitor for signs of opioid withdrawal.
<i>Examples:</i>	Macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g. ketoconazole), protease inhibitors (e.g., ritonavir), grapefruit juice.
CYP3A4 Inducers	
<i>Clinical Impact:</i>	The concomitant use of Fentanyl Citrate Injection and CYP3A4 inducers can decrease the plasma concentration of fentanyl [see <i>Clinical Pharmacology</i> (12.3)], resulting in decreased efficacy or onset of a withdrawal syndrome in patients who have developed physical dependence to fentanyl [see <i>Warnings and Precautions</i> (5.3)]. After stopping a CYP3A4 inducer, as the effects of the inducer decline, the fentanyl plasma concentration will increase [see <i>Clinical Pharmacology</i> (12.3)],

	which could increase or prolong both the therapeutic effects and adverse reactions, and may cause serious respiratory depression.
<i>Intervention:</i>	If concomitant use is necessary, consider increasing the Fentanyl Citrate Injection dosage until stable drug effects are achieved. Monitor for signs of opioid withdrawal. If a CYP3A4 inducer is discontinued, consider Fentanyl Citrate Injection dosage reduction and monitor for signs of respiratory depression.
<i>Examples:</i>	Rifampin , carbamazepine, phenytoin
Benzodiazepines and Other Central Nervous System (CNS) Depressants	
<i>Clinical Impact:</i>	The concomitant use of Fentanyl Citrate Injection with CNS depressants may result in decreased pulmonary artery pressure and may cause hypotension. Even small dosages of diazepam may cause cardiovascular depression when added to high dose or anesthetic dosages of Fentanyl Citrate Injection. As postoperative analgesia, concomitant use of Fentanyl Citrate Injection can increase the risk of hypotension, respiratory depression, profound sedation, coma, and death.
<i>Intervention:</i>	As postoperative analgesia, start with a lower dose of Fentanyl Citrate Injection and monitor patients for signs of respiratory depression, sedation, and hypotension. Fluids or other measures to counter hypotension should be available. [see <i>Warnings and Precautions (5.4)</i>].
<i>Examples:</i>	Benzodiazepines and other sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol.
Serotonergic Drugs	
<i>Clinical Impact:</i>	The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system has resulted in serotonin syndrome [see <i>Warnings and Precautions (5.7)</i>].
<i>Intervention:</i>	If concomitant use is warranted, carefully observe the patient, particularly during treatment initiation and dose adjustment. Discontinue Fentanyl Citrate Injection if serotonin syndrome is suspected.
<i>Examples:</i>	Selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT ₃ receptor antagonists, drugs that effect the serotonin neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), certain muscle relaxants (i.e., cyclobenzaprine, metaxalone), monoamine oxidase (MAO) inhibitors (those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue).
Monoamine Oxidase Inhibitors	
<i>Clinical Impact:</i>	MAOI interactions with opioids may manifest as serotonin syndrome [see <i>Warnings and Precautions (5.7)</i>] or opioid toxicity (e.g., respiratory depression, coma) [see <i>Warnings and Precautions (5.2)</i>]
<i>Intervention:</i>	The use of Fentanyl Citrate Injection is not recommended for patients taking MAOIs or within 14 days of stopping such treatment.
<i>Examples:</i>	Phenelzine, tranylcypromine, linezolid
Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics	
<i>Clinical Impact:</i>	May reduce the analgesic effect of Fentanyl Citrate Injection and/or precipitate withdrawal symptoms.
<i>Intervention:</i>	Avoid concomitant use.
<i>Examples:</i>	Butorphanol, nalbuphine, pentazocine, buprenorphine.
Muscle Relaxants	
<i>Clinical Impact:</i>	Fentanyl may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression.
	Monitor patients for signs of respiratory depression that may be greater than

<i>Intervention:</i>	otherwise expected and decrease the dosage of Fentanyl Citrate Injection and/or the muscle relaxant as necessary.
Diuretics	
<i>Clinical Impact:</i>	Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone.
<i>Intervention:</i>	Monitor patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage of the diuretic as needed.
Anticholinergic Drugs	
<i>Clinical Impact:</i>	The concomitant use of anticholinergic drugs may increase risk of urinary retention and/or severe constipation, which may lead to paralytic ileus.
<i>Intervention:</i>	Monitor patients for signs of urinary retention or reduced gastric motility when Fentanyl Citrate Injection is used concomitantly with anticholinergic drugs.
Neuroleptics	
<i>Clinical Impact:</i>	Elevated blood pressure, with and without pre-existing hypertension, has been reported following administration of Fentanyl Citrate Injection combined with a neuroleptic [see <i>Warnings and Precautions (5.13)</i>].
<i>Intervention:</i>	ECG monitoring is indicated when a neuroleptic agent is used in conjunction with Fentanyl Citrate Injection as an anesthetic premedication, for the induction of anesthesia, or as an adjunct in the maintenance of general or regional anesthesia.
Nitrous oxide	
<i>Clinical Impact:</i>	Nitrous oxide has been reported to produce cardiovascular depression when given with higher doses of Fentanyl Citrate Injection.
<i>Intervention:</i>	Monitor patients for signs of cardiovascular depression that may be greater than otherwise expected.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Prolonged use of opioid analgesics during pregnancy may cause neonatal opioid withdrawal syndrome. Available data with Fentanyl Citrate Injection in pregnant women are insufficient to inform a drug-associated risk for major birth defects and miscarriage.

In animal reproduction studies, fentanyl administration to pregnant rats during organogenesis was embryocidal at doses within the range of the human recommended dosing. No evidence of malformations was noted in animal studies completed to date [see Data]. The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

Clinical Considerations

Fetal/Neonatal Adverse Reactions

Prolonged use of opioid analgesics during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth.

Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns

for symptoms of neonatal opioid withdrawal syndrome and manage accordingly.

Labor or Delivery

Opioids cross the placenta and may produce respiratory depression and psycho-physiologic effects in neonates. An opioid antagonist, such as naloxone, must be available for reversal of opioid-induced respiratory depression in the neonate. Fentanyl Citrate Injection is not recommended for use in pregnant women during or immediately prior to labor, when other analgesic techniques are more appropriate. Opioid analgesics, including Fentanyl Citrate Injection, can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression.

Data

Animal Data

Fentanyl has been shown to be embryocidal in pregnant rats at doses of 30 mcg/kg intravenously (0.05 times the human dose of 100 mcg/kg on a mg/m² basis) and 160 mcg/kg subcutaneously (0.26 times the human dose of 100 mcg/kg on a mg/m² basis). There was no evidence of teratogenicity reported.

No evidence of malformations or adverse effects on the fetus was reported in a published study in which pregnant rats were administered fentanyl continuously via subcutaneously implanted osmotic minipumps at doses of 10, 100, or 500 mcg/kg/day starting 2-weeks prior to breeding and throughout pregnancy. The high dose was approximately 0.81 times the human dose of 100 mcg/kg on a mg/m² basis.

8.2 Lactation

Risk Summary

Fentanyl is present in breast milk. However, there is insufficient information to determine the effects of fentanyl on the breastfed infant and the effects of fentanyl on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Fentanyl Citrate Injection and any potential adverse effects on the breastfed infant from Fentanyl Citrate Injection or from the underlying maternal condition.

Clinical Considerations

Monitor infants exposed to Fentanyl Citrate Injection through breast milk for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

8.3 Females and Males of Reproductive Potential

Infertility

Chronic use of opioids may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [see *Adverse Reactions* (6), *Clinical Pharmacology* (12.2), *Nonclinical Toxicology* (13.1)].

8.4 Pediatric Use

The safety and efficacy of Fentanyl Citrate Injection in children under two years of age has not been established.

Rare cases of unexplained clinically significant methemoglobinemia have been reported in premature neonates undergoing emergency anesthesia and surgery which included the combined use of fentanyl, pancuronium and atropine. A direct cause and effect relationship between the combined use of these drugs and the reported cases of methemoglobinemia has not been established.

8.5 Geriatric Use

Elderly patients (aged 65 years or older) may have increased sensitivity to fentanyl. In general, use caution when selecting a dosage for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

Respiratory depression is the chief risk for elderly patients treated with opioids, and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration. Titrate the dosage of Fentanyl Citrate Injection slowly in geriatric patients and monitor closely for signs of central nervous system and respiratory depression [*see Warnings and Precautions (5.2)*].

Fentanyl is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

8.6 Hepatic Impairment

Fentanyl Citrate Injection should be administered with caution to patients with liver dysfunction because of the extensive hepatic metabolism. Reduce the dosage as needed and monitor closely for signs of respiratory depression, sedation, and hypotension.

8.7 Renal Impairment

Fentanyl Citrate Injection should be administered with caution to patients with kidney dysfunction because of the renal excretion of fentanyl and its metabolites. Reduce the dosage as needed and monitor for signs of respiratory depression, sedation, and hypotension.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Fentanyl Citrate Injection contains fentanyl, a Schedule II controlled substance.

9.2 Abuse

Fentanyl Citrate Injection contains fentanyl, a substance with a high potential for abuse similar to other opioids including hydrocodone, hydromorphone, methadone, morphine, oxycodone, oxymorphone, and tapentadol. Fentanyl Citrate Injection can be abused and is subject to misuse, addiction, and criminal diversion [*see Warnings and Precautions (5.1)*].

All patients treated with opioids require careful monitoring for signs of abuse and addiction, since use of opioid analgesic products carries the risk of addiction even under appropriate medical use.

Prescription drug abuse is the intentional non-therapeutic use of a prescription drug, even once, for its rewarding psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that develop after repeated substance use and includes: a strong desire to take the drug, difficulties in controlling its use, persisting in its use despite harmful consequences, a higher priority given to drug use than to other activities and obligations, increased tolerance, and sometimes a physical withdrawal.

Fentanyl Citrate Injection, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Risks Specific to Abuse of Fentanyl Citrate Injection

Abuse of Fentanyl Citrate Injection poses a risk of overdose and death. The risk is increased with concurrent use of Fentanyl Citrate Injection with alcohol and other central nervous system depressants.

Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

9.3 Dependence

Both tolerance and physical dependence can develop during chronic opioid therapy. Tolerance is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). Tolerance may occur to both the desired and undesired effects of drugs, and may develop at different rates for different effects.

Physical dependence results in withdrawal symptoms after abrupt discontinuation or a significant dosage reduction of a drug. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene), mixed agonist/antagonist analgesics (e.g., pentazocine, butorphanol, nalbuphine), or partial agonists (e.g., buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

10 OVERDOSAGE

Clinical Presentation

Acute overdose with Fentanyl Citrate Injection can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and in some cases, pulmonary edema, bradycardia, hypotension, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations [see *Clinical Pharmacology* (12.2)].

Treatment of Overdose

In case of overdose, priorities are the reestablishment of a patent and protected airway and institution of assisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life-support techniques.

The opioid antagonists, naloxone or nalmefene, are specific antidotes to respiratory depression resulting from opioid overdose. For clinically significant respiratory or circulatory depression secondary to fentanyl overdose, administer an opioid antagonist. Opioid antagonists should not be administered in the absence of clinically significant respiratory or circulatory depression secondary to fentanyl overdose.


Because the duration of opioid reversal is expected to be less than the duration of action of fentanyl in Fentanyl Citrate Injection, carefully monitor the patient until spontaneous respiration is reliably re-established. If the response to an opioid antagonist is suboptimal or only brief in nature, administer additional antagonist as directed by the product's prescribing information.

In an individual physically dependent on opioids, administration of the recommended usual dosage of the antagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms experienced will depend on the degree of physical dependence and the dose of the antagonist administered. If a decision is made to treat serious respiratory depression in the physically dependent patient, administration of the antagonist should be initiated with care and by titration with smaller than usual doses of the antagonist.

11 DESCRIPTION

Fentanyl Citrate Injection is an opioid agonist. Fentanyl Citrate Injection, is a sterile, nonpyrogenic

solution of fentanyl citrate in water for injection, available as 50 mcg (0.05 mg) per mL which is administered only by the intravenous or intramuscular routes of injection. The chemical name is N-(1-phenethyl-4-piperidyl) propionanilide citrate (1:1). The molecular weight is 528.60. Its molecular formula is $C_{22}H_{28}N_2O \cdot C_6H_8O_7$, and it has the following chemical structure.



Fentanyl citrate, a white powder which is sparingly soluble in water. Each milliliter contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5).

The solution contains no bacteriostat, antimicrobial agent or added buffer and is intended only for use as a single-dose injection. When smaller doses are required, the unused portion should be discarded in an appropriate manner.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Fentanyl is an opioid agonist whose principal actions of therapeutic value are analgesic and sedation.

12.2 Pharmacodynamics

Effects on the Central Nervous System

Fentanyl produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation.

Fentanyl causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

Effects on the Gastrointestinal Tract and Other Smooth Muscle

Fentanyl causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone may be increased to the point of spasm resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, and transient elevations in serum amylase.

Effects on the Cardiovascular System

Fentanyl produces peripheral vasodilation which may result in orthostatic hypotension or syncope. Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes, sweating, and/or orthostatic hypotension.

Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotrophic hormone (ACTH), cortisol, and luteinizing

hormone (LH) in humans [see *Adverse Reactions* (6)]. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see *Adverse Reactions* (6)].

Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system in *in vitro* and animal models. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

Concentration–Efficacy Relationships

A dose of 100 mcg (0.1 mg) (2.0 mL) of Fentanyl Citrate Injection is approximately equivalent in analgesic activity to 10 mg of morphine or 75 mg of meperidine.

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with potent agonist opioids. The minimum effective analgesic concentration of fentanyl for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome and/or the development of analgesic tolerance [see *Dosage and Administration* (2.1, 2.2)].

The onset of action of fentanyl is almost immediate when the drug is given intravenously; however, the maximal analgesic effect may not be noted for several minutes. The usual duration of action of the analgesic effect is 30 to 60 minutes after a single intravenous dose of up to 100 mcg (0.1 mg) (2 mL). Following intramuscular administration, the onset of action is from seven to eight minutes, and the duration of action is one to two hours.

Concentration–Adverse Reaction Relationships

There is a relationship between increasing fentanyl plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions [see *Dosage and Administration* (2.1, 2.2)].

The onset of action of fentanyl is almost immediate when the drug is given intravenously; however, the maximal respiratory depressant effect may not be noted for several minutes. As with longer acting opioid analgesics, the duration of the respiratory depressant effect of fentanyl may be longer than the analgesic effect. The following observations have been reported concerning altered respiratory response to CO₂ stimulation following administration of fentanyl citrate:

- Diminished sensitivity to CO₂ stimulation may persist longer than depression of respiratory rate. (Altered sensitivity to CO₂ stimulation has been demonstrated for up to four hours following a single dose of 600 mcg (0.6 mg) (12 mL) fentanyl citrate to healthy volunteers). Fentanyl frequently slows the respiratory rate, duration and degree of respiratory depression being dose-related.
- The peak respiratory depressant effect of a single intravenous dose of Fentanyl Citrate Injection is noted 5 to 15 minutes following injection [see *Warnings and Precautions* (5.2)].

12.3 Pharmacokinetics

Fentanyl Citrate Injection is administered by the intravenous or intramuscular route. The pharmacokinetics of fentanyl can be described as a three-compartment model.

Distribution

Fentanyl plasma protein binding decreases with increasing ionization of the drug. Alterations in pH may affect its distribution between plasma and the central nervous system. It accumulates in skeletal muscle

and fat, and is released slowly into the blood. The volume of distribution for fentanyl is 4 L/kg. It has a distribution time of 1.7 minutes and redistribution time of 13 minutes.

Elimination

The terminal elimination half-life is 219 minutes.

Fentanyl, which is primarily transformed in the liver, demonstrates a high first-pass clearance and releases approximately 75% of an intravenous dose in urine, mostly as metabolites with less than 10% representing the unchanged drug. Approximately 9% of the dose is recovered in the feces, primarily as metabolites.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies in animals to evaluate the carcinogenic potential of Fentanyl Citrate Injection have not been conducted.

Mutagenesis

Studies in animals to evaluate the mutagenic potential of fentanyl citrate injection have not been conducted.

Impairment of Fertility

Decreased pregnancy rates occurred in a multigenerational study in which pregnant rats were treated subcutaneously during the first 21 days of pregnancy with 160 mcg/kg to 1250 mcg/kg fentanyl (0.26 times to 2.0 times a human dose of 100 mcg/kg based on body surface area).

Studies in animals to characterize the effect of fentanyl on male fertility have not been conducted.

16 HOW SUPPLIED/STORAGE AND HANDLING

Fentanyl Citrate Injection, USP equivalent to 50 mcg (0.05 mg) fentanyl per mL, is supplied in single-dose glass containers as follows:

Unit of Sale	Concentration
NDC 0409-9093-32 Clamcell of 10, 2 mL Single-dose Ampules	100 mcg Fentanyl/2 mL (50 mcg/mL)
NDC 0409-9094-22 Tray containing 25, 2 mL Single-dose Flip-top Vials	100 mcg Fentanyl/2 mL (50 mcg/mL)
NDC 0409-9093-35 Clamcell of 10, 5 mL Single-dose Ampules	250 mcg Fentanyl/5 mL (50 mcg/mL)
NDC 0409-9094-25 Tray containing 25, 5 mL Single-dose Flip-top Vials	250 mcg Fentanyl/5 mL (50 mcg/mL)
NDC 0409-9093-36 Carton containing 5, 10 mL Single-dose Ampules	500 mcg Fentanyl/10 mL (50 mcg/mL)
NDC 0409-9094-28 Tray containing 25, 10 mL Single-dose Flip-top Vials	500 mcg Fentanyl/10 mL (50 mcg/mL)
NDC 0409-9093-38 Carton containing 5, 20 mL Single-dose Ampules	1000 mcg Fentanyl/20 mL (50 mcg/mL)
NDC 0409-9094-31 Tray containing 25, 20 mL Single-dose Flip-top Vials	1000 mcg Fentanyl/20 mL (50 mcg/mL)
NDC 0409-9094-61	2500 mcg Fentanyl/50 mL

Tray containing 25, 50 mL Single-dose Flip-top Vials

(50 mcg/mL)

Protect from light. Retain in carton until time of use.

Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

17 PATIENT COUNSELING INFORMATION

Serotonin Syndrome

Inform patients that opioids could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. Instruct patients to inform their healthcare provider if they are taking, or plan to take serotonergic medications [see *Warnings and Precautions (5.7)*, *Drug Interactions (7)*].

Constipation

Advise patients of the potential for severe constipation [see *Clinical Pharmacology (12.2)*].

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

LAB-0843-5.0

PRINCIPAL DISPLAY PANEL - 2 mL Ampule Label

2 mL

Rx only

NDC 0409-9093-37

Fentanyl Citrate

Injection, USP

100 mcg

Fentanyl/2 mL

(50 mcg/mL)

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA

RL-7711

CII


For Intravenous or

Intramuscular use.

Protect from light.

Retain in carton

until time of use.



PRINCIPAL DISPLAY PANEL - 2 mL Ampule Cello Pack Label

2 mL Single-dose 10 Ampuls

Rx only

NDC 0409-9093-32

Contains 10 of NDC 0409-9093-37

Fentanyl Citrate Injection, USP

100 mcg Fentanyl/2 mL

(50 mcg/mL)

CII

Protect from light. Keep ampuls in tray until time of use.

For Intravenous or Intramuscular use.

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). Usual dosage: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

Distributed by

Hospira, Inc., Lake Forest, IL 60045 USA

Hospira

RL-7313

LOT ##-###-AA

EXP DMMYYYYY

Tamper Evident Tray

Directions for ampul verification:

1. Verify shrink-wrap is intact and all ampuls present.

2. Do not unwrap tray prior to dispensing.

For your convenience in recording narcotic use

INITIAL/DATE

- 1.
- 2.
- 3.
- 4.
- 5.

INITIAL/DATE

- 6.
- 7.
- 8.
- 9.
- 10.

2 mL Single-dose 10 Ampuls		Rx only NDC 0409-9093-32 Contains 10 of NDC 0409-9093-37	Tamper Evident Tray												
Fentanyl Citrate Injection, USP 100 mcg Fentanyl/2 mL (50 mcg/mL)			Directions for ampul verification: 1. Verify shrink-wrap is intact and all ampuls present. 2. Do not unwrap tray prior to dispensing. For your convenience in recording narcotic use <table style="width: 100%;"><thead><tr><th style="width: 50%;">INITIAL/DATE</th><th style="width: 50%;">INITIAL/DATE</th></tr></thead><tbody><tr><td>1. _____</td><td>6. _____</td></tr><tr><td>2. _____</td><td>7. _____</td></tr><tr><td>3. _____</td><td>8. _____</td></tr><tr><td>4. _____</td><td>9. _____</td></tr><tr><td>5. _____</td><td>10. _____</td></tr></tbody></table>	INITIAL/DATE	INITIAL/DATE	1. _____	6. _____	2. _____	7. _____	3. _____	8. _____	4. _____	9. _____	5. _____	10. _____
INITIAL/DATE	INITIAL/DATE														
1. _____	6. _____														
2. _____	7. _____														
3. _____	8. _____														
4. _____	9. _____														
5. _____	10. _____														
Protect from light. Keep ampuls in tray until time of use. For Intravenous or Intramuscular use. Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). Usual dosage: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]		 (01)20304099093320													
Distributed by Hospira, Inc., Lake Forest, IL 60045 USA															
<div style="display: flex; justify-content: space-between;"><div>RL-7313</div><div>LOT ##-##-AA EXP DMMYYYY</div></div>															

PRINCIPAL DISPLAY PANEL - 5 mL Ampule Label

5 mL

NDC 0409-9093-45

Rx only

Fentanyl Citrate

Injection, USP

250 mcg Fentanyl/5 mL

CII

(50 mcg/mL)

For Intravenous or Intramuscular use. Protect from light.

Retain in carton until time of use.

Dist. by Hospira, Inc., Lake Forest, IL 60045 USA

RL-7698

LOT ##-##-AA

EXP DMMYYYY



PRINCIPAL DISPLAY PANEL - 5 mL Ampule Cello Pack Label

5 mL Single-dose 10 Ampuls

NDC 0409-9093-35

Contains 10 of NDC 0409-9093-45

Fentanyl Citrate Injection, USP

Rx only

250 mcg Fentanyl/5 mL

CII

(50 mcg/mL)

Protect from light. Keep ampuls in tray until time of use.

For Intravenous or Intramuscular use.

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). Usual dosage: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

Hospira

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

RL-7315

LOT ##-###-AA

EXP DMMYYYYY

Tamper Evident Tray

Directions for ampul verification:

1. Verify shrink-wrap is intact and all ampuls present.
2. Do not unwrap tray prior to dispensing.



For your convenience in recording narcotic use


INITIAL/DATE

- 1.
- 2.
- 3.
- 4.
- 5.

INITIAL/DATE

- 6.
- 7.
- 8.
- 9.
- 10.

5 mL Single-dose 10 Ampuls		NDC 0409-9093-35													
Fentanyl Citrate		Contains 10 of NDC 0409-9093-45													
250 mcg Fentanyl/5 mL		Injection, USP Rx only													
(50 mcg/mL)															
Protect from light. Keep ampuls in tray until time of use.		<div style="border: 1px solid black; padding: 2px; text-align: center; margin-bottom: 5px;">Tamper Evident Tray</div> <div style="font-size: 0.8em;">Directions for ampul verification: 1. Verify shrink-wrap is intact and all ampuls present. 2. Do not unwrap tray prior to dispensing.</div> <div style="margin-top: 10px;">For your convenience in recording narcotic use</div> <table style="width: 100%; border-collapse: collapse;"><thead><tr><th style="width: 50%; text-align: center;">INITIAL/DATE</th><th style="width: 50%; text-align: center;">INITIAL/DATE</th></tr></thead><tbody><tr><td>1. _____</td><td>6. _____</td></tr><tr><td>2. _____</td><td>7. _____</td></tr><tr><td>3. _____</td><td>8. _____</td></tr><tr><td>4. _____</td><td>9. _____</td></tr><tr><td>5. _____</td><td>10. _____</td></tr></tbody></table>		INITIAL/DATE	INITIAL/DATE	1. _____	6. _____	2. _____	7. _____	3. _____	8. _____	4. _____	9. _____	5. _____	10. _____
INITIAL/DATE	INITIAL/DATE														
1. _____	6. _____														
2. _____	7. _____														
3. _____	8. _____														
4. _____	9. _____														
5. _____	10. _____														
For Intravenous or Intramuscular use.															
Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). Usual dosage: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]															
Distributed by Hospira, Inc., Lake Forest, IL 60045 USA															
RL-7315		LOT 88-888-AA EXP 08/2011													


FP0/7.5 mL
(01)20304099093351

PRINCIPAL DISPLAY PANEL - 10 mL Ampule Label

10 mL Single-dose

Rx only

Discard unused portion

Fentanyl Citrate

Injection, USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)

CII

FOR INTRAVENOUS USE BY HOSPITAL
PERSONNEL SPECIFICALLY TRAINED IN
THE USE OF NARCOTIC
ANALGESICS.

10 mL Single-dose

Discard unused portion

Rx only

NDC 0409-9093-41

Fentanyl Citrate

Injection, USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)

FOR INTRAVENOUS USE BY HOSPITAL
PERSONNEL SPECIFICALLY TRAINED IN
THE USE OF NARCOTIC
ANALGESICS.



Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain NaOH and/or HCl for pH adjustment. pH 4.7 (4.0 to 7.5). Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

Protect from light. Retain in carton until time of use.

Hospira, Inc.

Lake Forest, IL 60045 USA



RL-5195

PRINCIPAL DISPLAY PANEL - 10 mL Ampule Carton

10 mL

Single-dose

5 Ampuls

Discard unused portion

NDC 0409-9093-36

Fentanyl Citrate Injection, USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)

CII

Hospira

Protect from light. Retain in carton until time of use.

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid

for pH adjustment. pH 4.7 (4.0 to 7.5). Sterile, nonpyrogenic.

Usual dose: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

10 mL Single-dose
5 Ampuls

Discard unused portion

NDC 0409-9093-36

Fentanyl Citrate Injection, USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)



Hospira

Protect from light. Retain in carton until time of use.

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). Sterile, nonpyrogenic.

Usual dose: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

CA-4227



(01) 20304099093368

For your convenience in recording narcotic use
INITIAL/DATE

1	
2	
3	
4	
5	

NDC 0409-9093-36

10 mL Single-dose
5 Ampuls

Fentanyl Citrate Injection, USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)

Hospira



(50 mcg/mL)

500 mcg Fentanyl/10 mL

Fentanyl Citrate Injection, USP

10 mL Single-dose
5 Ampuls

NDC 0409-9093-36



Hospira

Hospira

Protect from light. Retain in carton until time of use.

500 mcg Fentanyl/10 mL

(50 mcg/mL)

Fentanyl Citrate Injection, USP

10 mL Single-dose
5 Ampuls

Discard unused portion

NDC 0409-9093-36



INSPECT™ Tamper Evident Carton

Directions for ampul verification:

1. Upon receipt, inspect carton. Verify tamper evident tape is not broken. Do not break tape prior to dispensing.
2. Lift front flap. Verify carton contains 5 ampuls. Reclose the flap.

10 mL Single-dose
5 Ampuls

NDC 0409-9093-36

Discard unused portion

Rx only

Fentanyl Citrate
Injection, USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)



Protect from light.

Retain in carton until time of use.

**FOR INTRAVENOUS ADMINISTRATION BY
HOSPITAL PERSONNEL SPECIFICALLY TRAINED
IN THE USE OF NARCOTIC ANALGESICS.**

Hospira, Inc., Lake Forest, IL 60045 USA

Hospira

20 mL Single-dose

Discard unused portion

Rx only

Fentanyl

Citrate

Injection, USP

1000 mcg Fentanyl/20 mL

(50 mcg/mL)

FOR INTRAVENOUS USE BY HOSPITAL
PERSONNEL SPECIFICALLY TRAINED IN
THE USE OF NARCOTIC ANALGESICS.

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

NDC 0409-9093-31

Each mL contains fentanyl (as
the citrate) 50 mcg (0.05 mg).
May contain NaOH and/or HCl
for pH adjustment. pH 4.7 (4.0
to 7.5). Protect from light.

Retain in carton until time of
use. Store at 20 to 25°C (68 to
77°F). [See USP Controlled
Room Temperature.]

CII

Hospira

RL-7316

LOT ##-###-AA

EXP DMMMYYYY

20 mL Single-dose

Discard unused portion

Rx only

NDC 0409-9093-31

**Fentanyl
Citrate
Injection, USP**

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain NaOH and/or HCl for pH adjustment. pH 4.7 (4.0 to 7.5). **Protect from light.** Retain in carton until time of use. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

FPO / 7.5 Mil

1000 mcg Fentanyl/20 mL
(50 mcg/mL)

FOR INTRAVENOUS USE BY HOSPITAL
PERSONNEL SPECIFICALLY TRAINED IN
THE USE OF NARCOTIC ANALGESICS.

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA



Hospira

RL-7316

LOT ##-##-AA
EXP DMMYYY

PRINCIPAL DISPLAY PANEL - 20 mL Ampule Carton

20 mL

Single-dose

5 Ampuls

NDC 0409-9093-38

Contains 5 of NDC 0409-9093-31

Discard unused portion

Rx only

Fentanyl Citrate

Injection, USP

1000 mcg Fentanyl/20 mL

(50 mcg/mL)

CII

Protect from light.

Retain in carton until time of use.

FOR INTRAVENOUS ADMINISTRATION BY
HOSPITAL PERSONNEL SPECIFICALLY TRAINED
IN THE USE OF NARCOTIC ANALGESICS.

Distributed by

Hospira, Inc., Lake Forest, IL 60045 USA

Hospira

20 mL Single-dose
5 Ampuls

Discard unused portion



NDC 0409-9093-38



Fentanyl Citrate Injection, USP

1000 mcg Fentanyl/20 mL
(50 mcg/mL)

Protect from light. Retain in carton until time of use.

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). Sterile, nonpyrogenic.
Usual dose: See insert. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]

20 mL Single-dose
5 Ampuls
Fentanyl Citrate Injection, USP
1000 mcg Fentanyl/20 mL
(50 mcg/mL)

NDC 0409-9093-38



CA-7144

GTIN 20304099093382



LOT #####A
SN #####
EXP DMMYYYY



FPO/10 mL

(01)20304099093382

For your convenience in recording narcotic use
INITIAL/DATE

1	_____
2	_____
3	_____
4	_____
5	_____



Protect from light. Retain in carton until time of use.

1000 mcg Fentanyl/20 mL
(50 mcg/mL)

Fentanyl Citrate Injection, USP

20 mL Single-dose
5 Ampuls

Discard unused portion



NDC 0409-9093-38

INSPECT™ Tamper Evident Carton

Directions for ampul verification:

1. Upon receipt, inspect carton. Verify tamper evident tape is not broken. Do not break tape prior to dispensing.
2. Lift front flap. Verify carton contains 5 ampuls. Reclose the flap.

20 mL Single-dose
5 Ampuls

NDC 0409-9093-38

Contains 5 of NDC 0409-9093-31

Discard unused portion

Rx only

Fentanyl Citrate Injection, USP

1000 mcg Fentanyl/20 mL
(50 mcg/mL)

Protect from light.

Retain in carton until time of use.

FOR INTRAVENOUS ADMINISTRATION BY
HOSPITAL PERSONNEL SPECIFICALLY TRAINED
IN THE USE OF NARCOTIC ANALGESICS.



NDC 0409-9093-38

Fentanyl Citrate Injection, USP

1000 mcg Fentanyl/20 mL
(50 mcg/mL)

20 mL Single-dose
5 Ampuls



Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA



PRINCIPAL DISPLAY PANEL - 2 mL Vial Label

2 mL Single-dose
NDC 0409-9094-12
Rx only

Intravenous or
Intramuscular Use

Fentanyl Citrate
Inj., USP
CII
100 mcg Fentanyl/2 mL
(50 mcg/mL)

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

Hospira
RL-7310

2 mL Single-dose
Intravenous or
Intramuscular Use

NDC 0409-9094-12
Rx only

Fentanyl Citrate
Inj., USP
100 mcg Fentanyl/2 mL
(50 mcg/mL)

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA



RL-7310

LOT ###-##-AA

EXP DMMYYY

PRINCIPAL DISPLAY PANEL - 2 mL Vial Tray Top


2 mL
Single-dose
25 Fliptop Vials
Discard unused portion.
NDC 0409-9094-22
Contains 25 of NDC 0409-9094-12
Fentanyl Citrate

Injection, USP
100 mcg Fentanyl/2 mL
(50 mcg/mL)
CII

Protect from light.
Retain in tray until time of use.
For Intravenous or Intramuscular use.
Rx only

CA-7136

Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA
Hospira




PRINCIPAL DISPLAY PANEL - 2 mL Vial Tray

2 mL
Single-dose 25 Fliptop Vials
NDC 0409-9094-22
Discard unused portion.

Fentanyl Citrate Inj., USP
100 mcg Fentanyl/2 mL
(50 mcg/mL)
Rx only
CII

Retain in tray until time of use.
Protect from light.
For Intravenous or Intramuscular use.

Hospira



PRINCIPAL DISPLAY PANEL - 5 mL Vial Label

5 mL

Single-dose Vial

NDC 0409-9094-18

Rx only

Fentanyl Citrate

Injection, USP

CII

250 mcg Fentanyl/5 mL

(50 mcg/mL)

Distributed by Hospira, Inc.

Lake Forest, IL 60045 USA

Hospira

5 mL NDC 0409-9094-18
Single-dose Vial Rx only

Fentanyl Citrate
Injection, USP



250 mcg Fentanyl/5 mL

(50 mcg/mL)

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA



**For Intravenous
or Intramuscular
Use**

**Protect from
light.**

**Retain in tray
until time
of use.**



RL-7308

LOT
##-##-AA
EXP
DDMMYY

PRINCIPAL DISPLAY PANEL - 5 mL Vial Tray Top

5 mL Single-dose

25 Fliptop Vials

Rx only

NDC 0409-9094-25

Contains 25 of NDC 0409-9094-18

Fentanyl Citrate

Injection, USP

CII

250 mcg Fentanyl/5 mL

(50 mcg/mL)

For Intravenous or Intramuscular use.

Protect from light.

Retain in tray until time of use.

Discard unused portion.

Distributed by

Hospira, Inc., Lake Forest, IL 60045 USA

Hospira

CA-7140

5 mL Single-dose
25 Flip-top Vials

Rx only NDC 0409-9094-25
Contains 25 of NDC 0409-9094-18

Fentanyl Citrate

Injection, USP



250 mcg Fentanyl/5 mL

(50 mcg/mL)

**For Intravenous or Intramuscular use.
Protect from light.
Retain in tray until time of use.
Discard unused portion.**

Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA




CA-7140

PRINCIPAL DISPLAY PANEL - 5 mL Vial Tray

5 mL Single-dose Vials
25 Units
NDC 0409-9094-25
Rx only

Fentanyl Citrate Injection, USP
CII
250 mcg Fentanyl/5 mL
(50 mcg/mL)

INTRAVENOUS or INTRAMUSCULAR USE. Protect from light.
Retain in tray until time of use. Discard unused portion.



PRINCIPAL DISPLAY PANEL - 10 mL Vial Label

10 mL Single-dose Vial NDC 0409-9094-17

Discard unused portion. Rx only

Fentanyl Citrate Inj., USP

500 mcg Fentanyl/10 mL

CII

(50 mcg/mL)

Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA

10 mL Single-dose Vial NDC 0409-9094-17

Discard unused portion. Rx only

Fentanyl Citrate Inj., USP

500 mcg Fentanyl/10 mL

(50 mcg/mL)

Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA



Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain NaOH and/or HCl for pH adjustment. pH 4.7 (4.0 to 7.5). **FOR I.V. USE BY HOSPITAL PERSONNEL SPECIFICALLY TRAINED IN THE USE OF NARCOTIC ANALGESICS.** Usual dosage: See insert. **Protect from light.** Retain in carton until time of use. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]



RL-7306

LOT ##-##-##-AA
EXP DMMYYY

PRINCIPAL DISPLAY PANEL - 10 mL Vial Tray

10 mL Single-dose 25 Fliptop Vials

Discard unused portion

NDC 0409-9094-28

Contains 25 of NDC 0409-9094-17

Fentanyl Citrate Injection, USP CII


500 mcg Fentanyl/10 mL

(50 mcg/mL)

Rx only

Protect from light. Retain in carton until time of use.

Hospira



PRINCIPAL DISPLAY PANEL - 20 mL Vial Label

20 mL Single-dose
Vial NDC 0409-9094-16

Discard unused portion.
Rx only

Fentanyl Citrate
Injection, USP
CII

1000 mcg Fentanyl/20 mL
(50 mcg/mL)

Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA
Hospira

20 mL Single-dose Vial NDC 0409-9094-16

Discard unused portion. Rx only

**Fentanyl Citrate
Injection, USP**

**1000 mcg Fentanyl/20 mL
(50 mcg/mL)**

Distributed by
Hospira, Inc., Lake Forest, IL 60045 USA



Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain NaOH and/or HCl for pH adjustment. pH 4.7 (4.0 to 7.5). **FOR INTRAVENOUS USE BY HOSPITAL PERSONNEL SPECIFICALLY TRAINED IN THE USE OF NARCOTIC ANALGESICS.** Usual dosage: See insert. **Protect from light.** Retain in carton until time of use. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]



RL-7307

LOT ##-##-AA
EXP DMMYYY

PRINCIPAL DISPLAY PANEL - 20 mL Vial Tray

20 mL Single-dose 25 Fliptop Vials

Discard unused portion.

Rx only

NDC 0409-9094-31

Contains 25 of NDC 0409-9094-16


Fentanyl Citrate Injection, USP

CII

1000 mcg Fentanyl/20 mL (50 mcg/mL)

Protect from light. Retain in carton until time of use.

Hospira



PRINCIPAL DISPLAY PANEL - 50 mL Vial Label

50 mL
 Single-dose Vial
 Discard unused portion.

Fentanyl Citrate
 Injection, USP
 2500 mcg Fentanyl/50 mL
 CII

(50 mcg/mL)
 Rx only

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA

Hospira

50 mL Single-dose Vial Discard unused portion. NDC 0409-9094-41

**Fentanyl Citrate
Injection, USP
2500 mcg Fentanyl/50 mL**

(50 mcg/mL)



Rx only

Each mL contains fentanyl (as the citrate) 50 mcg (0.05 mg). May contain sodium hydroxide and/or hydrochloric acid for pH adjustment. pH 4.7 (4.0 to 7.5). **FOR INTRAVENOUS USE BY HOSPITAL PERSONNEL SPECIFICALLY TRAINED IN THE USE OF NARCOTIC ANALGESICS.** Usual dosage: See insert. **Protect from light.** Retain in carton until time of use. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.]



RL-7311

LOT ##-##-AA
EXP DMMYY

Distributed by Hospira, Inc.
Lake Forest, IL 60045 USA



PRINCIPAL DISPLAY PANEL - 50 mL Vial Carton

One/NDC 0409-9094-41

Rx only

50 mL

Single-dose

Fliptop Vial

Discard unused portion.

Fentanyl

Citrate

Injection, USP


2500 mcg Fentanyl/50 mL

CII

(50 mcg/mL)

Protect from light.

Retain in carton
until time of use.




PRINCIPAL DISPLAY PANEL - 50 mL Vial Carton Tray

50 mL
 Single-dose
 25 Fliptop Vials
 Discard unused portion.
 NDC 0409-9094-61
 Contains 25 of NDC 0409-9094-41
 Rx only
 Fentanyl Citrate Injection, USP
 2500 mcg Fentanyl/50 mL
 (50 mcg/mL)
 CII

Protect from light. Retain in carton until time of use.

Hospira



FENTANYL CITRATE

fentanyl citrate injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-9093

Route of Administration		INTRAMUSCULAR, INTRAVENOUS		DEA Schedule		CII	
Active Ingredient/Active Moiety							
Ingredient Name					Basis of Strength		Strength
FENTANYL CITRATE (UNII: MUN5LYG46H) (FENTANYL - UNII:UF599785JZ)					FENTANYL		50 ug in 1 mL
Inactive Ingredients							
Ingredient Name						Strength	
SODIUM HYDRO XIDE (UNII: 55X04QC32I)							
HYDROCHLORIC ACID (UNII: QTT17582CB)							
WATER (UNII: 059QF0KO0R)							
Packaging							
#	Item Code	Package Description			Marketing Start Date		Marketing End Date
1	NDC:0409-9093-32	10 in 1 CELLO PACK			11/15/2005		
1	NDC:0409-9093-37	2 mL in 1 AMPULE; Type 0: Not a Combination Product					
2	NDC:0409-9093-35	10 in 1 CELLO PACK			12/14/2005		
2	NDC:0409-9093-45	5 mL in 1 AMPULE; Type 0: Not a Combination Product					
3	NDC:0409-9093-36	5 in 1 CARTON			07/20/2006		09/01/2011
3	NDC:0409-9093-41	10 mL in 1 AMPULE; Type 0: Not a Combination Product					
4	NDC:0409-9093-38	5 in 1 CARTON			03/01/2006		09/01/2018
4	NDC:0409-9093-31	20 mL in 1 AMPULE; Type 0: Not a Combination Product					
Marketing Information							
Marketing Category		Application Number or Monograph Citation			Marketing Start Date		Marketing End Date
NDA		NDA019115			11/15/2005		

FENTANYL CITRATE

fentanyl citrate injection, solution

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0409-9094
Route of Administration	INTRAMUSCULAR, INTRAVENOUS	DEA Schedule	CII
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
FENTANYL CITRATE (UNII: MUN5LYG46H) (FENTANYL - UNII:UF599785JZ)		FENTANYL	50 ug in 1 mL

Inactive Ingredients				
Ingredient Name				Strength
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
HYDROCHLORIC ACID (UNII: QTT17582CB)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0409-9094-22	25 in 1 TRAY	12/02/2010	
1	NDC:0409-9094-12	2 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:0409-9094-25	25 in 1 TRAY	11/09/2005	
2	NDC:0409-9094-18	5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
3	NDC:0409-9094-28	25 in 1 TRAY	02/17/2006	
3	NDC:0409-9094-17	10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
4	NDC:0409-9094-31	25 in 1 TRAY	09/27/2005	
4	NDC:0409-9094-16	20 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
5	NDC:0409-9094-61	25 in 1 TRAY	12/22/2005	
5		1 in 1 CARTON		
5	NDC:0409-9094-41	50 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA019115	09/27/2005	

Labeler - Hospira, Inc. (141588017)

Establishment			
Name	Address	ID/FEI	Business Operations
Hospira, Inc.		093132819	ANALYSIS(0409-9093, 0409-9094) , LABEL(0409-9093, 0409-9094) , MANUFACTURE(0409-9093, 0409-9094) , PACK(0409-9093, 0409-9094)

Establishment			
Name	Address	ID/FEI	Business Operations
Hospira, Inc.		827731089	ANALYSIS(0409-9093, 0409-9094)