

**MIGRAINE RELIEF- acetaminophen, aspirin, caffeine tablet
Family Dollar (FAMILY WELLNESS)**

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen. If this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraine
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing, bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warnings applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for

- diabetes,
- gout,
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stool
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask a doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- close cap tightly after use
- read all product information before using

Inactive ingredients

carnauba wax, hydroxypropyl cellulose, hypromellose, microcrystalline cellulose, polyethylene glycol, sodium stearyl fumarate, titanium dioxide

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENTS IN EXCEDRIN® MIGRAINE†

MIGRAINE RELIEF

Acetaminophen Aspirin and Caffeine Tablets, USP

250 mg, 250 mg and 65 mg

Pain reliever/Pain reliever aid

CAPLETS**

(**Capsule-shaped tablets)

†This product is not manufactured or distributed by GSK Consumer Healthcare Holdings (US) LLC, distributor of Excedrin® Migraine

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY

MIDWOOD BRANDS, LLC

500 VOLVO PKWY,

CHESAPEAKE, VA 23320

Package Label



Alcohol-free. By Migraine.™
Pain reliever. Pain reliever aid.
Ask a doctor before use if you have ever had:

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month. Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

alcoholic drinks every day while using this product.
■ Take more or for a longer time than directed
■ Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.
■ Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use ■ if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer. ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if ■ you have ever had migraines diagnosed by a health professional. ■ you have a headache that is different from your usual migraines. ■ you have the worst headache of your life. ■ you have fever and stiff neck. ■ you have headaches beginning after or caused by head injury, exertion, coughing or bending. ■ you experience your first headache after the age of 50. ■ you have daily headaches. ■ you have a migraine so severe as to require bed rest. ■ you have liver disease. ■ stomach bleeding warning applies to you. ■ you have a history of stomach problems, such as heartburn, or kidney disease. ■ you are taking a diuretic. ■ you have asthma. ■ you have problems or serious side effects from taking pain relievers or fever reducers. ■ you have vomiting with your migraine headache.

Ask a doctor or pharmacist before use if you are ■ taking a prescription drug for diabetes

■ do not use more than directed
■ adults: take 2 caplets with a glass of water
■ if symptoms persist or worsen, ask your doctor
■ do not take more than 2 caplets 24 hours, unless directed by a doctor. ■ under 18 years of age: ask a doctor notice any signs or symptoms.

Directions ■ store at controlled room temperature 20° to 25°C (68° to 77°F). ■ close cap tightly ►

Other information ■ store all product information before using after use. ■ read all product information before using

►

Drug Facts (continued)
Inactive ingredients: camphor, wax, hydroxypropyl cellulose, hydrophilic, microcrystalline cellulose, polyethylene glycol, sodium stearyl fumarate, titanium dioxide.

Questions or comments?
Call 1-877-753-3535 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by GSK Consumer Healthcare Products (US) LLC, distributor of Excedrin® Migraine.
TAMPER EVIDENCE: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUT OF CARTON FOR COMPLETE
WARNINGS AND PRODUCT INFORMATION.

Drug Facts

Active ingredients (in each caplet) Purposes

Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
Transdermal anti-inflammatory drug

Use ■ Treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 2 caplets in 24 hours, which is the maximum daily amount. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older. ■ have had stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug. ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others). ■ have 3 or more

COMPARE TO THE ACTIVE INGREDIENTS
IN EXCEDRIN® MIGRAINE!

MIGRAINE RELIEF

Acetaminophen, Aspirin and Caffeine Tablets, USP
250 mg, 250 mg and 65 mg

Pain Reliever/Pain Reliever Aid

24 CAPLETS**
(**Capsule-shaped tablets)

ACTUAL SIZE

NDC 55319-970-24

FAMILY WELLNESS Migraine Relief

MIGRAINE RELIEF

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-970(NDC:62207-680)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0E1Z)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Capsule shaped Tablet)	Size	17mm
Flavor		Imprint Code	I1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-970-24	1 in 1 BOX	07/01/2023	12/31/2026
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55319-970-10	1 in 1 BOX	07/01/2023	12/31/2026
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214039	07/01/2023	12/31/2026

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)