

IBUPROFEN- ibuprofen tablet, film coated
CHAIN DRUG MARKETING ASSOCIATION INC

Quality Choice 44-291

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - toothache
 - the common cold
 - backache
 - headache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - slurred speech
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

**QC[®]
Quality
Choice**

NDC 63868-773-50

†Compare to the
Active Ingredient in
Advil[®] Tablets

Ibuprofen

Ibuprofen Tablets USP, 200 mg

Pain Reliever,
Fever Reducer **(NSAID)**

50 Coated Tablets

actual size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by
PF Consumer Healthcare 1 LLC, owner of the
registered trademark Advil® Tablets.

50844 REV1221C29115

• *SATISFACTION*

100%

QC®

GUARANTEED•

Distributed by CDMA, Inc.
43157 W Nine Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

No Print / No Varnish
Lot no. & Exp. date



QUALITY CHOICE
SATISFACTION
GUARANTEED
Distributed by CDMA, Inc.
12517 W. New Mile
Nashville, TN 37217
www.qualitychoice.com
Questions: 800-935-2382

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active Ingredient
(in each brown tablet)
Ibuprofen USP
200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose
Ibuprofen USP
200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
• menstrual cramps
• toothache
• the common cold
• backache
• headache
• muscular aches
• minor pain of arthritis
• temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• rash
• skin redness
• asthma (wheezing)
• facial swelling
• shock
• blisters
• hives
• If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
• take more or for a longer time than directed
• take a blood thinning (anticoagulant) or steroid drug
• are age 60 or older
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have had stomach ulcers or bleeding problems
• have 3 or more alcoholic drinks every day while using this product

NDC 63868-773-50

Compare to the Active Ingredient in Advil® Tablets

Ibuprofen

Ibuprofen Tablets USP, 200 mg

Pain Reliever, Fever Reducer (NSAID)

50 Coated Tablets

actual size

Drug Facts (continued)
microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide
Questions or comments?
Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday
PF Consumer Healthcare 1 LLC, owner of the registered trademark Advil® Tablets.
50844 REV1221C29115

Drug Facts (continued)
• pain gets worse or lasts more than 10 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present in the painful area
• any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 36 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• do not take more than directed
• the smallest effective dose should be used
• adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
• if pain or fever does not respond to 1 tablet, 2 tablets may be used
• do not exceed 6 tablets in 24 hours, unless directed by a doctor
• children under 12 years: ask a doctor

Other information
• store between 20°-25°C (68°-77°F)
• avoid excessive heat 40°C (104°F)
• see end flap for expiration date and lot number

Inactive ingredients carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate,

Drug Facts (continued)
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use
• if you have ever had an allergic reaction to any other pain reliever/fever reducer
• right before or after heart surgery
Ask a doctor before use if
• you have a history of stomach problems, such as heartburn
• you are taking a diuretic
• you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
• stomach bleeding warning applies to you
• you have problems or serious side effects from taking pain relievers or fever reducers
Ask a doctor or pharmacist before use if you are
• under a doctor's care for any serious condition
• taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
• taking any other drug
When using this product
• take with food or milk if stomach upset occurs
Stop use and ask a doctor if
• you experience any of the following signs of stomach bleeding:
• feel faint
• vomit blood
• have bloody or black stools
• have stomach pain that does not get better
• you have symptoms of heart problems or stroke:
• chest pain
• trouble breathing
• leg swelling
• slurred speech
• weakness in one part or side of body

Quality Choice 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:63868-773

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CARNAUBA WAX (UNII: R12CBM0EIZ)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STARCH, CORN (UNII: O8232NY3SJ)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)				
MAGNESIUM STEARATE (UNII: 70097M6I3O)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	brown	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;291	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-773-10	1 in 1 CARTON	05/24/1988	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63868-773-05	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	
3	NDC:63868-773-24	1 in 1 CARTON	05/24/1988	06/01/2027
3		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:63868-773-50	1 in 1 CARTON	05/24/1988	06/01/2027
4		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA075010	05/24/1988	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(63868-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63868-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(63868-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(63868-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63868-773)

Revised: 10/2025

CHAIN DRUG MARKETING ASSOCIATION INC