

PAIN RELIEF- acetaminophen tablet, film coated, extended release
Meijer Distribution Inc

Meijer Distribution, Inc. Pain Relief Drug Facts

Active ingredient (in each caplet)

Acetaminophen 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- muscular aches
- backache
- minor pain of arthritis
- toothache
- premenstrual and menstrual cramps
- headache
- the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

- pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> • take 2 caplets every 8 hours with water • swallow whole; do not crush, chew, split or dissolve • do not take more than 6 caplets in 24 hours • do not use for more than 10 days unless directed by a doctor
children under 12 years	<ul style="list-style-type: none"> • do not use

Other information

- store at 20-25°C (68-77°F)
- **do not use if printed foil under cap is broken or missing**
- meets the requirements of USP Dissolution Test 2

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

meijer®

Compare to Tylenol® 8HR active ingredient

pain relief

Acetaminophen Extended-Release Tablets, 650 mg | Pain Reliever | Fever Reducer
8 HOUR Pain Relief

For up to 8 Hours Relief of Minor Muscle Aches & Pain

24 Caplets*

650 mg each

*Capsule-Shaped Tablets | actual size

meijer

NDC 41250-802-62

Compare to Tylenol® 8HR active ingredient**

pain relief


Acetaminophen Extended-Release Tablets,
650 mg | Pain Reliever | Fever Reducer

8 HOUR Pain Relief

24 Caplets*

650 mg each

*Capsule-Shaped Tablets



actual size

For up to 8 Hours
Relief of Minor
Muscle Aches & Pain

Contains No Aspirin
U.S. Patent 7,887,172

DIST. BY MEIJERDISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



WWW.MEIJER.COM/
SATISFACTION



**This product is not manufactured or distributed
by Johnson & Johnson, owner of the registered
trademark Tylenol®.

Drug Facts

Active ingredient **Purpose**
(in each caplet)

Acetaminophen 650 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
 - minor pain of arthritis
 - toothache
 - premenstrual and menstrual cramps
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor
children under 12 years	do not use

Other information store at 20-25°C (68-77°F)

- do not use if printed foil under cap is broken or missing
- meets the requirements of USP Dissolution Test 2

Inactive ingredients carnauba wax, colloidal silicon dioxide, croscarmellose sodium, hypromellose, magnesium stearate, methylcellulose, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, pregelatinized starch, stearic acid, titanium dioxide

Questions or comments? 1-800-719-9260

PID 4039720



7 13733 92100 8



PAIN RELIEF

acetaminophen tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-802
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L544
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-802-62	1 in 1 CARTON	03/03/2017	
1		24 in 1 BOTTLE; Type 0: Not a Combination		

1		Product		
2	NDC:41250-802-78	1 in 1 CARTON	03/03/2017	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075077	03/03/2017	

Labeler - Meijer Distribution Inc (006959555)

Revised: 10/2022

Meijer Distribution Inc