# TUSSIN CF- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid AptaPharma Inc.

-----

#### **Drug Facts**

# Active ingredients (in each 5 mL tsp)

Dextromethorphan HBr, USP 10 mg Guaifenesin, USP 100 mg Phenylephrine HCL, USP 5 mg

#### Purpose

Dextromethorphan HBr USP ......Cough Suppressant Guailenesin USP ......Expectorant Phenylephrine HCI, USP ......Nasal Decongestant

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation.

#### Warnings

Do not use if you are now taking aprescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychialric or emotional conditions or Parkinson's disease), or for 2 weeks aMer stopping the MAOI drug. If you do not know il your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- heart disease
  high blood pressure
- thyroid disease
  diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use il you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

#### Stop use and ask a doctor if

- you get nervous, diuy or sleepless
- symptoms do not get better within 7 days or are

accompanied by fever

cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of aserious condition.

#### If pregnant or breast feeding,

ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact aPoison Control Center right away.

#### **Directions**

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age

AGE	DOSE	
Adults & children 12 years & over	2 teaspoons every 4 hours	
Children under 12 years	2 teaspoons every 4 hours	

#### Other information

- **■** Keep carton for full Direction for use
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

#### Inactive ingredients

anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at **1-877·798-5944** 

**Product Label** 

AP *SAFE* NDC 76821-504-24

**COMPARE TO THE ACTIVE** 

**INGREDIENT IN ROBITUSSIN®** 

**PEAK COLD** 

**MULTI-SYMPTOM COLD\*** 

**Adult Tussin** 

#### **Multi-Symptom Cold**

Dextromethorphan HBr / Guaifensesin /

**Phenylephrine HCL** 

Cough Supressant / Expectorant /

**Nasal Decongestant** 

**Non-Drowsy** 

#### Relieves:

- Cough
- Mucus
- Nasal Congestion
- For Ages 12 and over

4 FL OZ (118 mL)

CF

# DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

5 1013430 3/20

\*This product is not manufactured or distributed by Plizer, owner of the registered trademark Robitussin<sup>®</sup>

Manufactured by: AptaPharma Inc., 1533 Union Ave, Pennsauken, NJ 08110

Ap·BX·05

Carton



#### **Bottle**

Nasal Congestion Cough · Mucus





For Ages 12 and ove

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

### Drug Facts

Active ingredients (in each 5 mL tsp) Purpose Phenylephrine HCI, USP 5 mg Nasal Decongestant

Uses ■ helps loosen phlegm (mucus) and bronchial secretions to drain bronchial tubes temporarily relieves these symptoms occurring with a cold: nasal congestion cough due to minor throat and bronchial irritation.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Stop use and ask a doctor if wou get nervous, dizzy or sleepless

- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children.

#### Directions - do not take more than 6 doses in any 24 hour period

AGE	DOSE
Adults & children 12 years & over	2 teaspoons every 4 hours
Children under 12 years	do not use

Other information ■ store at 20-25°C (68-77°F) ■ do not refrigerate ■ sodium 3 mg per teaspoonful ■ dosage cup provided ■ See carton for full labeling

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

#### Questions? 1-877-798-5944

Manufactured by: AptaPharma Inc.,

1533 Union Ave. Pennsauken, NJ 08110 AP-LR-05

EXP:

LOT:

# **TUSSIN CF**

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76281-504
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	<b>Basis of Strength</b>	Strength		
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL		
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL		

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6A3C0OX)			
MENTHOL (UNII: L7T10EIP3A)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SORBITOL SOLUTION (UNII: 8KW3E207O2)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				

Packaging				
# Item Code Package Description		Marketing Start Date	Marketing End Date	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	

N	<b>darketing</b> In	formation		
	Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M012	09/30/2020	

### Labeler - AptaPharma Inc. (790523323)

# **Registrant -** AptaPharma Inc. (790523323)

Establishment				
Name	Address	ID/FEI	Business Operations	
AptaPharma Inc.		790523323	manufacture(76281-504)	

Revised: 12/2023 AptaPharma Inc.