ANTI-DIARRHEAL- loperamide hcl tablet L.N.K. International, Inc.

Convenience Valet 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl. **Heart alert:** Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

• drink plenty of clear fluids to help prevent dehydration caused by diarrhea

• find right dose on chart. If possible, use weight to dose; otherwise, use age.

	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
5	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

1-844-428-2538

Principal Display Panel

24/7 life

BY 7-ELEVEN_{TM}

Anti-

Diarrheal

Looperamide HCl Tablets, 2 mg

Anti-Diarrheal

Controls the Symptoms

of Diarrhea

compare to

Imodium® A-D

active ingredient*

12

CAPLETS

Actual Size

QUALITY GUARANTEED

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D. 50844 REV0619A37502

DISTRIBUTED BY 7-ELEVEN, INC. IRVING, TX 75063 WWW.7-ELEVEN.COM

Satisfaction Guaranteed 1-800-255-0711

1. To open, tear at perforations and remove a section.

2. Use 1 to cut through plastic for caplet.

USB controls symptoms of diarrhea, including Travelers' Diarrhea Anti-diarrheal Loperamide HCI 2 mg Purpose Active ingredient (in each caplet) Drug Facts KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION



7-ELEVEN

Anti-Diarrheal

Loperamide HCl Tablets, 2 mg Anti-Diarrheal

Controls the Symptoms of Diarrhea

compare to **Imodium® A-D** active ingredient*





This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D. 50844 REV0619A37502

DISTRIBUTED BY 7-ELEVEN, INC. IRVING, TX 75063 WWW.7-ELEVEN.COM

Satisfaction Guaranteed 1-800-255-0711

2. Use % to cut through plastic for caplet.

B-1205-375-02 REV0619A37502

0

Questions or comments? 1-844-428-2538

FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide Inactive ingredients com starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate,

> ■ see end flap for expiration date and lot number ■ store between 20°-25° (68°-77°F)

> > **Drug Facts** (continued)

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN Other information

əsu fon ob	children under 2 years (up to 33 lbs)
азк а doctor	children 2-5 years (34-47 lbs)
1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours	children 6-8 years (48-59 lbs)
1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours	children 9-11 years (60-95 lbs)
2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours	adults and children 12 years and over


find right dose on chart. If possible, use weight to dose; otherwise, use age. ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

.1-800-222-1222) right away.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center If pregnant or breast-feeding, ask a health professional before use.

■ you get abdominal swelling or bulging. These may be signs of a serious condition. Stop use and ask a dodor if symptoms get worse diarrhea lasts for more than 2 days

When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating



7-11 44-375

ANTI-DIARRHEAL

loperamide hcl tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-259
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg		

Inactive Ingredients				
Ingredient Name	Strength			
STARCH, CORN (UNII: O8232NY3SJ)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
DIBASIC CALCIUM PHO SPHATE DIHYDRATE (UNII: O7TSZ97GEP)				
FD&C BLUE NO. 1ALUMINUM LAKE (UNII: J9EQA3S2JM)				
MAGNESIUM STEARATE (UNII: 70097M6130)				
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)				
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)				

Product Characteristics				
Color	GREEN (light)	Score	2 pieces	
Shape	OVAL	Size	10 mm	
Flavor		Imprint Code	44;375	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50844-259-02	2 in 1 CARTON	07/15/2019		
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA076497	07/15/2019		

Labeler - L.N.K. International, Inc. (038154464)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(50844-259)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(50844-259)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(50844-259)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(50844-259)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(50844-259)

Revised: 12/2019 L.N.K. International, Inc.