

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) LIGHT BEIGE-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) FAIR IVORY-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) LIGHT IVORY-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) LIGHT OCHRE-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) MEDIUM BEIGE-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) MEDIUM IVORY-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) MEDIUM OCHRE-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) DARK BEIGE-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) DARK IVORY-** octinoxate, octocrylene, and titanium dioxide powder

**SHISEIDO AMERICAS CORPORATION**

-----

## **SHISEIDO UV PROTECTIVE COMPACT (REFILL)**

### ***Drug Facts***

<b><i>ACTIVE INGREDIENTS:</i></b>	<b><i>Purpose</i></b>
OCTINOXATE 7.4%	Sunscreen
OCTOCRYLENE 5.0%	Sunscreen
TITANIUM DIOXIDE 11.8%	Sunscreen

### **Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

### **Warnings**

**For external use only**

**Do not use** on damaged or broken skin

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor** if rash occurs

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. – 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

### **Inactive Ingredients**

SYNTHETIC FLUORPHLOGOPITE, DIMETHICONE, SILICA, NYLON-12, VINYL DIMETHICONE/METHICONE SILSESQUIOXANE  
CROSSPOLYMER, PETROLATUM, METHICONE, TRIETHYLHEXANOIN, PEG/PPG-14/7 DIMETHYL ETHER, TOCOPHERYL ACETATE, SAXIFRAGA SARMENTOSA EXTRACT, SODIUM HYALURONATE, SOPHORA ANGUSTIFOLIA ROOT EXTRACT, SORBITAN SESQUIESTEARATE, HYDROGEN DIMETHICONE, ALUMINUM HYDROXIDE, WATER, POLYSILICONE-2, ALCOHOL, SODIUM MAGNESIUM SILICATE, HYDRATED SILICA, TRIETHOXYCAPRYLYLSILANE, ALUMINUM DISTEARATE, BUTYLENE GLYCOL, GLYCERIN, TOCOPHEROL, BHT, DISTEARYLDIMONIUM CHLORIDE, SYZYGIIUM JAMBOS LEAF EXTRACT, TIN OXIDE, CHLORPHENESIN, FRAGRANCE, MICA, TITANIUM DIOXIDE, IRON OXIDES, ZINC OXIDE,

### **Other information**

- protect this product in this container from excessive heat and direct sun.

### **Questions or comments?**

Call toll free 1-800-906-7503

### **PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Fair Ivory**

#### **SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Fair Ivory  
(SP90)

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes  
before sun exposure
- use a water resistant  
sunscreen if swimming or  
sweating
- reapply at least every 2 hours

■ **Sun Protection Measures.**

Spending time in the sun  
increases your risk of skin  
cancer and early skin aging.

To decrease this risk, regularly  
use a sunscreen with a broad  
spectrum SPF of 15 or higher  
and other sun protection  
measures including:

- limit time in the sun,  
especially from 10 a.m. – 2  
p.m.

- wear long-sleeve shirts,  
pants, hats, and sunglasses

- children under 6 months: Ask  
a doctor

SHISEIDO

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHLOGOPITE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHICONE/METHICONE SILSESQUOXANE COPOLYMER • PETROLATUM • METHICONE • TRIETHYLHEXANOLIN • PEG/PPG-14/7 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAXIFRAGA SAPMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBITAN SESQUISTEARATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIMETHOXYCAPRYLYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEAROYL DIMONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLOROPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs ▶

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11180  
www.shiseido.com



7 30852 11180 6

11180-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Light Beige**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Light Beige  
(SP20)

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36  
SUNSCREEN

12g NET WT. .42 OZ.

SHISEIDO

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- **Sun Protection Measures.**  
Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. – 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: Ask a doctor

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHOSPHATE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHICONE/METHICONE SILSESQUOXANE COPOLYMER • PETROLATUM • METHICONE • TRIMETHYLHEXANDIOL • PEG/PPG-14/7 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAXIFRAGA SAPRMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBETAN SESQUISTEARATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIETHOXYCAPRYLYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEAROYL DIMONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLORPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish.
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

**For external use only**

**Do not use** on damaged or broken skin

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor** if rash occurs ▶

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11181  
[www.shiseido.com](http://www.shiseido.com)



7 30852 11181 3

11181-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Light Ivory**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Light Ivory

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

SHISEIDO

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes  
before sun exposure
- use a water resistant  
sunscreen if swimming or  
sweating
- reapply at least every 2 hours

■ **Sun Protection Measures.**

Spending time in the sun  
increases your risk of skin  
cancer and early skin aging.  
To decrease this risk, regularly  
use a sunscreen with a broad  
spectrum SPF of 15 or higher  
and other sun protection  
measures including:

- limit time in the sun,  
especially from 10 a.m. – 2  
p.m.
- wear long-sleeve shirts,  
pants, hats, and sunglasses

- children under 6 months: Ask  
a doctor

12g NET WT. .42 OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHLOGOPITE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHACRYLATE • DIMETHACRYLATE • SILSESQUOXANE CROSSPOLYMER • PETROLATUM • METHICONE • TRIETHYLHEXANOLIN • PEG/PPG-14/7 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBETAN SESQUISOESTEATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIMETHOXYCAPRYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEAROYL MONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLORPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish.
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

Active ingredients	Purpose
OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11182  
www.shiseido.com

7 30852 11182 0

11182-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Light Ochre**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Light Ochre  
(SP30)

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours

■ **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. – 2 p.m.

- wear long-sleeve shirts, pants, hats, and sunglasses

- children under 6 months: Ask a doctor ▶

SHISEIDO

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHOSPHATE - DIMETHICONE - SILICA - NYLON-12 - VINYL DIMETHICONE/METHICONE SILSESQUOXANE COPOLYMER - PETROLATUM - METHICONE - TRIMETHYLHEXANDIN - PEG/PPG-14/7 DIMETHYL ETHER - TOCOPHERYL ACETATE - SAXIFRAGA SAPRMENTOSA EXTRACT - SODIUM HYALURONATE - SOPHORA ANGUSTIFOLIA ROOT EXTRACT - SORBETAN SESQUISTEARATE - HYDROGEN DIMETHICONE - ALUMINUM HYDROXIDE - WATER - POLYSILICONE-2 - ALCOHOL - SODIUM MAGNESIUM SILICATE - HYDRATED SILICA - TRIETHOXYCAPRYLYLSILANE - ALUMINUM DISTEARATE - BUTYLENE GLYCOL - GLYCERIN - TOCOPHEROL - BHT - DISTEARYL DIMONIUM CHLORIDE - SYZYGIUM JAMBOS LEAF EXTRACT - TIN OXIDE - CHLORPHENESIN - FRAGRANCE - MICA - TITANIUM DIOXIDE - IRON OXIDES - ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish.
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

**For external use only**

**Do not use on damaged or broken skin**

**When using this product keep out of eyes. Rinse with water to remove.**

**Stop use and ask a doctor if rash occurs** ▶

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11183  
www.shiseido.com



7 30852 11183 7

11183-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Medium Beige**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Medium Beige  
(SP60)

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

SHISEIDO

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours

■ **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. – 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

- children under 6 months: Ask a doctor

12g NET WT. .42 OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHLOGOPITE - DIMETHICONE - SILICA - NYLON-12 - VINYL DIMETHICONE/METHICONE SILSESQUOXANE CROSSPOLYMER - PETROLATUM - METHICONE - TRIETHYLHEXANOLIN - PEG/PPG-14/7 DIMETHYL ETHER - TOCOPHERYL ACETATE - SAXIFRAGA SARMENTOSA EXTRACT - SODIUM HYALURONATE - SOPHORA ANGUSTIFOLIA ROOT EXTRACT - SORBITAN SESQUISULFATE - HYDROGEN DIMETHICONE - ALUMINUM HYDROXIDE - WATER - POLYSILOXANE-2 - ALCOHOL - SODIUM MAGNESIUM SILICATE - HYDRATED SILICA - TRIMETHOXYCAPRYLYLSILANE - ALUMINUM DISTEARATE - BUTYLENE GLYCOL - GLYCERIN - TOCOPHEROL - BHT - DISTEARYL DIMONIUM CHLORIDE - SYZYGIUM JAMBOS LEAF EXTRACT - TIN OXIDE - CHLORPHENESIN - FRAGRANCE - MICA - TITANIUM DIOXIDE - IRON OXIDES - ZINC OXIDE -

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish.
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11184  
www.shiseido.com

7 30852 11184 4

11184-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Medium Ivory**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Medium Ivory  
(SP50)

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes  
before sun exposure
- use a water resistant  
sunscreen if swimming or  
sweating
- reapply at least every 2 hours

■ **Sun Protection Measures.**

Spending time in the sun  
increases your risk of skin  
cancer and early skin aging.

To decrease this risk, regularly  
use a sunscreen with a broad  
spectrum SPF of 15 or higher  
and other sun protection  
measures including:

- limit time in the sun,  
especially from 10 a.m. – 2  
p.m.

- wear long-sleeve shirts,  
pants, hats, and sunglasses

- children under 6 months: Ask  
a doctor

SHISEIDO

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHOSPHATE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHICONE/METHACRYLATE • SILSESQUOXANE CROSSPOLYMER • PETROLATUM • METHACRYLATE • TRIETHYLHEXANOLIN • PEG/PPG-14/7 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAXIFRAGA SAPMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBETAN SESQUISOESTEATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIETHOXYCAPRYLYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEARYLDIMONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLOROPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
- Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
- Provides a silky touch and a matte finish.
- NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs ▶

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11185  
www.shiseido.com



7 30852 11185 1

11185-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Medium Ochre**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation

(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Medium Ochre  
(SP40)

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes  
before sun exposure
- use a water resistant  
sunscreen if swimming or  
sweating
- reapply at least every 2 hours
- **Sun Protection Measures.**  
Spending time in the sun  
increases your risk of skin  
cancer and early skin aging.  
To decrease this risk, regularly  
use a sunscreen with a broad  
spectrum SPF of 15 or higher  
and other sun protection  
measures including:
  - limit time in the sun,  
especially from 10 a.m. – 2  
p.m.
  - wear long-sleeve shirts,  
pants, hats, and sunglasses
- children under 6 months: Ask  
a doctor

SHISEIDO

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36  
SUNSCREEN

12g NET WT. .42OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHOSPHATE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHICONE METHACRYLATE • SILSESQUOXANE CROSSPOLYMER • PETROLATUM • METHICONE • TRIMETHYLHEXANDIN • PEG/PPG-14/7 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAMIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBITAN SESQUISULFATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIETHOXYCAPRYLYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEAROYL DIMONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLORPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil UV 360™ technology.
  - Formulated with Profense CELL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish.
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
G.L.O. 11186  
www.shiseido.com

7 30852 11186 8

11186-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Dark Beige**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation

(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Dark Beige

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42OZ.

SHISEIDO

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes  
before sun exposure
- use a water resistant  
sunscreen if swimming or  
sweating
- reapply at least every 2 hours
- **Sun Protection Measures.**  
Spending time in the sun  
increases your risk of skin  
cancer and early skin aging.  
To decrease this risk, regularly  
use a sunscreen with a broad  
spectrum SPF of 15 or higher  
and other sun protection  
measures including:
  - limit time in the sun,  
especially from 10 a.m. – 2  
p.m.
  - wear long-sleeve shirts,  
pants, hats, and sunglasses
- children under 6 months: Ask  
a doctor ▶

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHLOGOPITE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHICONE METHACRYLATE • SILSESQUOXANE COPOLYMER • PETROLATUM • METHICONE • TRIETHYLHEXANON • PEG/PPG-147 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAXIFRAGA SARMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBETAN SESQUISULFATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIETHOXYCAPRYLYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEARYLDIMONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLOROPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**  
Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil-UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish.
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% .....	Sunscreen
OCTOCRYLENE 5.0% .....	Sunscreen
TITANIUM DIOXIDE 11.8% .....	Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11187  
www.shiseido.com

7 30852 11187 5

11187-41-5040

AAA

**PRINCIPAL DISPLAY PANEL - 12 g Tray Carton - Dark Ivory**

**SHI SEIDO**

Anti-Aging Foundation

**36**

UV Protective  
Compact Foundation

(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42 OZ.

UV Protective Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

Dark Ivory  
(SP70)

**Drug Facts**  
(continued)

**Warnings**

Keep out of reach of children.  
If product is swallowed, get  
medical help or contact a Poison  
Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes  
before sun exposure
- use a water resistant  
sunscreen if swimming or  
sweating
- reapply at least every 2 hours

■ **Sun Protection Measures.**

Spending time in the sun  
increases your risk of skin  
cancer and early skin aging.  
To decrease this risk, regularly  
use a sunscreen with a broad  
spectrum SPF of 15 or higher  
and other sun protection  
measures including:

- limit time in the sun,  
especially from 10 a.m. – 2  
p.m.
- wear long-sleeve shirts,  
pants, hats, and sunglasses

- children under 6 months: Ask  
a doctor

SHISEIDO

Anti-Aging Foundation



36

UV Protective  
Compact Foundation  
(Refill)

BROAD SPECTRUM  
SPF 36

SUNSCREEN

12g NET WT. .42OZ.

**Drug Facts**  
(continued)

**Inactive ingredients**

SYNTHETIC FLUOROPHOSPHATE • DIMETHICONE • SILICA • NYLON-12 • VINYL DIMETHICONE/METHICONE SILSESQUOXANE CROSSPOLYMER • PETROLATUM • METHICONE • TRIETHYLHEXANOLIN • PEG/PPG-14/7 DIMETHYL ETHER • TOCOPHERYL ACETATE • SAXIFRAGA SAPRIMENTOSA EXTRACT • SODIUM HYALURONATE • SOPHORA ANGUSTIFOLIA ROOT EXTRACT • SORBITAN SESQUISTEARATE • HYDROGEN DIMETHICONE • ALUMINUM HYDROXIDE • WATER • POLYSILOXANE-2 • ALCOHOL • SODIUM MAGNESIUM SILICATE • HYDRATED SILICA • TRIETHOXYCAPRYLYLSILANE • ALUMINUM DISTEARATE • BUTYLENE GLYCOL • GLYCERIN • TOCOPHEROL • BHT • DISTEAROYL DIMONIUM CHLORIDE • SYZYGIUM JAMBOS LEAF EXTRACT • TIN OXIDE • CHLORPHENESIN • FRAGRANCE • MICA • TITANIUM DIOXIDE • IRON OXIDES • ZINC OXIDE

**Other Information**

■ protect this product in this container from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-800-906-7503

SHISEIDO  
UV PROTECTIVE COMPACT FOUNDATION (REFILL)  
BROAD SPECTRUM  
SPF 36

A foundation that protects skin from damage caused by external aggressors such as UV rays and dryness. Resistant to water and sebum, it provides a long lasting, natural-looking finish, while safeguarding skin's beauty for the future.

- Protects skin from UVA/UVB rays and helps reduce the risk of visible photo-aging, with exclusive SuperVeil UV 360™ technology.
  - Formulated with Profense CEL™, a complex of ingredients that reduces enzyme activity,\* cell damage\* and DNA damage\* linked to the formation of wrinkles and dark spots.
  - Provides a silky touch and a matte finish
  - NON-COMEDOGENIC AND DERMATOLOGIST-TESTED
- \* in vitro test

**Drug Facts**

**Active ingredients**

**Purpose**

OCTINOXATE 7.4% ..... Sunscreen  
OCTOCRYLENE 5.0% ..... Sunscreen  
TITANIUM DIOXIDE 11.8% ..... Sunscreen

**Uses**

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

SHISEIDO AMERICAS CORPORATION DIST.  
NEW YORK, NY 10022  
MADE IN U.S.A.  
GLO. 11188  
www.shiseido.com

7 30852 11188 2

11188-41-5040

AAA

**SHISEIDO UV PROTECTIVE COMPACT (REFILL) LIGHT BEIGE**

octinoxate, octocrylene, and titanium dioxide powder

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58411-122
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>HYDRATED SILICA</b> (UNII: Y6O7T4G8P9)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)	
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:58411-122-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) FAIR IVORY

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-121
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

### Inactive Ingredients

Ingredient Name	Strength
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>HYDRATED SILICA</b> (UNII: Y6O7T4G8P9)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573Z X3X)	
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	

<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)
<b>MICA</b> (UNII: V8A1AW0880)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-121-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M020	02/01/2014		

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) LIGHT IVORY

octinoxate, octocrylene, and titanium dioxide powder

Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58411-123
<b>Route of Administration</b>	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g	
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g	

Inactive Ingredients		
Ingredient Name	Strength	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)		
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)		
<b>NYLON-12</b> (UNII: 446U8J075B)		
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)		
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)		
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)		
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)		
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: VR1WPI7EW8)		

<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>ALCOHOL</b> (UNII: 3K9958V90M)
<b>HYDRATED SILICA</b> (UNII: Y607T4G8P9)
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)
<b>MICA</b> (UNII: V8A1AW0880)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>FERROSO FERRIC OXIDE</b> (UNII: XM0M87F357)
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-123-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) LIGHT OCHRE

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58411-124
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

### Inactive Ingredients

Ingredient Name	Strength
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>HYDRATED SILICA</b> (UNII: Y607T4G8P9)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)	
<b>SYZYGIIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-124-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) MEDIUM BEIGE

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-125
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

### Inactive Ingredients

Ingredient Name	Strength
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>HYDRATED SILICA</b> (UNII: Y6O7T4G8P9)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)	
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)	

MICA (UNII: V8A1AW0880)

FERRIC OXIDE RED (UNII: 1K09F3G675)

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

FERROSFERRIC OXIDE (UNII: XM0M87F357)

ZINC OXIDE (UNII: SOI2LOH54Z)

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-125-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) MEDIUM IVORY

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-126
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
OCTOCRYLENE (UNII: 5A68WGF6VM) (OCTOCRYLENE - UNII:5A68WGF6VM)	OCTOCRYLENE	600 mg in 12 g
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

### Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
NYLON-12 (UNII: 446U8J075B)	
PETROLATUM (UNII: 4T6H12BN9U)	
METHICONE (20 CST) (UNII: 6777U11MKT)	
TRIETHYLHEXANOIN (UNII: 7K3W1BIU6K)	
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)	
.ALPHA.-TOCOPHEROL ACETATE, DL- (UNII: WR1WPI7EW8)	
SAXIFRAGA STOLONIFERA LEAF (UNII: O3TMV4903H)	

<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>ALCOHOL</b> (UNII: 3K9958V90M)
<b>HYDRATED SILICA</b> (UNII: Y607T4G8P9)
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)
<b>MICA</b> (UNII: V8A1AW0880)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-126-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) MEDIUM OCHRE

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58411-127
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

## Inactive Ingredients

Ingredient Name	Strength
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>HYDRATED SILICA</b> (UNII: Y607T4G8P9)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)	
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-127-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) DARK BEIGE

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-128
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

### Inactive Ingredients

Ingredient Name	Strength
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	
<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>HYDRATED SILICA</b> (UNII: Y6O7T4G8P9)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573Z X3X)	
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)	
<b>MICA</b> (UNII: V8A1AW0880)	

<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-128-30	1 in 1 CARTON	02/01/2014	05/01/2024
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	05/01/2024

## SHISEIDO UV PROTECTIVE COMPACT (REFILL) DARK IVORY

octinoxate, octocrylene, and titanium dioxide powder

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:58411-129
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	888 mg in 12 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	600 mg in 12 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1416 mg in 12 g

### Inactive Ingredients

Ingredient Name	Strength
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>NYLON-12</b> (UNII: 446U8J075B)	
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>TRIETHYLHEXANOIN</b> (UNII: 7K3W1BIU6K)	
<b>PEG/PPG-14/7 DIMETHYL ETHER</b> (UNII: 6DNW9T7YT2)	
<b>.ALPHA.-TOCOPHEROL ACETATE, DL-</b> (UNII: WR1WPI7EW8)	
<b>SAXIFRAGA STOLONIFERA LEAF</b> (UNII: O3TMV4903H)	
<b>HYALURONATE SODIUM</b> (UNII: YSE9PPT4TH)	

<b>SOPHORA FLAVESCENS ROOT</b> (UNII: IYR6K8KQ5K)
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>ALCOHOL</b> (UNII: 3K9958V90M)
<b>HYDRATED SILICA</b> (UNII: Y607T4G8P9)
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)
<b>ALUMINUM DISTEARATE</b> (UNII: 7P1HP1B9UI)
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)
<b>SYZYGIUM JAMBOS LEAF</b> (UNII: 407Z4W5LFF)
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)
<b>MICA</b> (UNII: V8A1AW0880)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-129-30	1 in 1 CARTON	02/01/2014	
1		12 g in 1 TRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	02/01/2014	

**Labeler** - SHISEIDO AMERICAS CORPORATION (193691821)

### Establishment

Name	Address	ID/FEI	Business Operations
Shiseido America Inc.		782677132	analysis(58411-121, 58411-122, 58411-123, 58411-124, 58411-125, 58411-126, 58411-127, 58411-128, 58411-129) , manufacture(58411-121, 58411-122, 58411-123, 58411-124, 58411-125, 58411-126, 58411-127, 58411-128, 58411-129)