

ACETAMINOPHEN- acetaminophen tablet
BluePoint Laboratories

Acetaminophen Extended-Release Tablets, USP 650mg

Drug Facts

Active Ingredient (in each extended release tablet)

Acetaminophen USP 650 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure

whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right

away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you

do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning).

Adults:

- take 2 tablets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 tablets in 24 hours
- do not use for more than 10 days unless directed by a doctor.

Under 18 years of age:

- ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F). Avoid excessive heat 40°C (104°F)
- **do not use if carton is opened or foil inner seal is broken**
- USP Dissolution test is pending

Inactive ingredients

colloidal silicon dioxide, hydroxyethyl cellulose, hypromellose, magnesium stearate,

microcrystalline

cellulose, povidone, pregelatinized starch (maize), sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call 1-855-274-4122

Manufactured by:
Aurobindo Pharma Limited
Unit-VII (SEZ), Mahabubnagar (Dt)-509302, India.


For BluePoint Laboratories

Rev: 03/2021

M.L. No.: 22/MN/AP/2009/F/R

Package Label - Principal Display Panel

Acetaminophen Extended-Release tablets USP 650mg (100 tablets Carton) NDC: 68001-495-00



Acetaminophen Extended-Release tablets USP 650mg (100 tablets Container Label)
 NDC: 68001-495-00

Unwinding Direction →

NDC 68001-495-00

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

8 HOUR ARTHRITIS PAIN
Acetaminophen
 Extended-Release Tablets USP

650 mg

Pain Reliever/Fever Reducer
 For the Temporary Relief of Minor Arthritis Pain

100 Extended-release tablets

Do not use if foil inner seal is broken.

Drug Facts

Active ingredient (in each extended-release tablet)
 Acetaminophen USP 650 mg

Purpose
 Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis ■ muscular aches ■ backache ■ premenstrual and menstrual cramps ■ the common cold ■ headache ■ toothache ■ temporarily reduces fever

Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 tablets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness/itching ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if: ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Dose adjustment may be needed. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning). Adults: ■ take 2 tablets every 8 hours with water ■ swallow whole, do not crush, chew, split or dissolve ■ do not take more than 6 tablets in 24 hours ■ do not use for more than 10 days unless directed by a doctor. Under 18 years of age: ■ ask a doctor

Other information ■ store at 20° to 25°C (68° to 77°F). Avoid excessive heat 40°C (104°F). ■ USP Dissolution test is pending

Inactive ingredients colloidal silicon dioxide, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, pregelatinized starch (maize), sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? call 1-855-274-4122

Manufactured by: Aurbindo Pharma Limited
 Unit VII (SEZ), Mahabubnagar (Dist-509302), India.
 For BluePoint Laboratories M.L. No.: 22MM/AP2009/FR
 Rev. 03/2021

P 142.7863 LM-4527

NWZ

* Lot: XXXXXXXXX
 EXP: MM/YYYY
 Prefix & Variables of L
 printed online during J

A/s: 120 x 45 mm

ACETAMINOPHEN			
acetaminophen tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68001-495
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg	
Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
TRIACETIN (UNII: XHX3C3X673)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			
HYDROXYETHYL CELLULOSE (140 MPA.S AT 5%) (UNII: 8136Y38GY5)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
SODIUM STARCH GLYCOLATE TYPE B POTATO (UNII: 27NA468985)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
MAGNESIUM STEARATE (UNII: 70097M6130)			

Product Characteristics

Color	white ((White to Off-White))	Score	no score
Shape	CAPSULE ((Caplet))	Size	19mm
Flavor		Imprint Code	l;06
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68001-495-00	1 in 1 CARTON	04/30/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207229	04/30/2021	

Labeler - BluePoint Laboratories (985523874)

Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Ltd		650381903	analysis(68001-495) , manufacture(68001-495)

Revised: 6/2024

BluePoint Laboratories