

STOMACH RELIEF- bismuth subsalicylate tablet, chewable
SDA Laboratories, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredient (in each tablet)

Bismuth subsalicylate 262mg

Purposes

Anti-diarrheal/ Upset stomach reliever

Uses

- controls diarrhea
- relieves upset stomach due to overindulgence in food or drink

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

- if you have bloody or black stool
- if you have an ulcer or bleeding problem

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse

- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- repeat dosage every 1/2 to 1 hour as needed
- do not take more than 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- **adults and children 12 years and over:** 2 tablets
- **children under 12 years:** ask a doctor

Other information

- **each tablet contains:** calcium 77 mg, salicylate 102 mg
- store at 15 to 30⁰°C (59 to 86⁰F)
- protect from moisture

Inactive ingredients

calcium carbonate, D&C red #27(Al-lake), dextrose, flavor(cherry), magnesium stearate, maltodextrin, silicon dioxide, sorbitol

Questions or comments?

Adverse drug event call 1-(714)-903-7952

Principal Display Panel

NDC 66424-045-30

Stomach Relief

Bismuth Subsalicylate 262 mg

Anti-Diarrheal/ Upset Stomach Reliever

Helps to Relieve:

- Upset Stomach
- Diarrhea
- Indigestion
- Heartburn
- Nausea

†Compare to the Active Ingredients in Pepto-Bismol

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by Proctor & Gamble, owner of the registered trademark Pepto-Bismol

NDC 66424-045-30

Stomach Relief

Bismuth Subsalicylate 262 mg

ANTI-DIARRHEAL / UPSET STOMACH RELIEVER

Helps to Relieve:
Upset Stomach • Diarrhea
Indigestion • Heartburn • Nausea

CMP 30 Chewable Tablets 

†Compare to the Active Ingredient in Pepto-Bismol

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Bismuth subsalicylate 262 mg.....Upset stomach reliever

Purposes
Anti-diarrheal/Upset stomach reliever

Uses
■ controls diarrhea
■ relieves upset stomach due to overindulgence in food or drink

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Do not take if you are
■ allergic to salicylates (including aspirin)
■ taking other salicylate products

Do not use
■ if you have bloody or black stool
■ if you have an ulcer or bleeding problem

Drug Facts continued under label

†This product is not manufactured or distributed by Proctor & Gamble, owner of the registered trademark Pepto-Bismol®.

 Manufactured in the U.S.A.
for CMP, Inc.,
Garden Grove, CA 92841
L506/112

Lot No.: 3
Exp. Date: 1

66424-04530



Drug Facts (continued)

Ask a doctor before use if you have
■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
■ anticoagulation (thinning the blood)
■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if
■ symptoms get worse
■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ repeat dosage every 1/2 to 1 hour as needed
■ do not take more than 8 doses in 24 hours
■ use until diarrhea stops but not more than 2 days
■ adults and children 12 years and over: 2 tablets
■ children under 12 years: ask a doctor

Other information
■ each tablet contains: calcium 77 mg, salicylate 102 mg
■ store at 15 to 30°C (59 to 86°F)
■ protect from moisture

Inactive ingredients calcium carbonate, D&C red #27 (Al-lake), dextrose, flavor (cherry), magnesium stearate, maltodextrin, silicon dioxide, sorbitol

Questions or comments? Adverse drug event call 1-(714)-903-7952

STOMACH RELIEF			
bismuth subsalicylate tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66424-045
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8 YE)	BISMUTH SUBSALICYLATE	262 mg	
Inactive Ingredients			
Ingredient Name	Strength		
CALCIUM CARBONATE (UNII: H0G9379FGK)			
D&C RED NO. 27 (UNII: 2LRS185U6K)			
DEXTROSE (UNII: IY9XDZ35W2)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			

MALTODEXTRIN (UNII: 7CVR7L4A2D)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
SORBITOL (UNII: 506T60A25R)			
Product Characteristics			
Color	pink (light)	Score	no score
Shape	ROUND	Size	16 mm
Flavor	CHERRY	Imprint Code	AP;045
Contains			
Packaging			
#	Item Code	Package Description	Marketing Start Date
1	NDC:66424-045-30	30 in 1 BOTTLE, PLASTIC	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	09/27/2012	

Labeler - SDA Laboratories, Inc. (948067889)

Revised: 8/2013

SDA Laboratories, Inc.