EQUATE BENZ-DEX COUGH AND THROAT LOZENGES- benzocaine lozenge Walmart

Benzocaine 7.5 mg Dextromethorphan Hydrobromide 5 mg

Active ingredients (in each lozenge)

Benzocaine 7.5 mg

Dextromethorphan hydrobromide 5 mg

Purposes

Oral Pain Reliever

Cough Suppressant

Uses temporarily relieves these symptoms:

- occasional minor mouth irritation, sore mouth and sore throat
- cough due to minor throat and bronchial irritation as may occur with the common cold

Warnings

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Sore throat warning: If sore threat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly.

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- for teething
- in children under 2 years of age

Ask a doctor before use if you have:

- persistent or chronic cough such as accurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling develops
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant of breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center right away.

Directions

adults and children 12 years and older - Allow 2 lozenges to dissolve slowly in mouth, one at a time; may be repeated every 4 hours, not to exceed 12 lozenges in 24 hours, or as directed by a doctor

children 6 to 12 years of age - take 1 lozenge and allow to dissolve slowly in mouth; may be repeated every 4 hours, not to exceed 6 lozenges in 24 hours, or as directed by a doctor

children under 6 years of age - do not use

Other information

- store at room temperature
- protect from moisture

Inactive ingredients: Acesulfame potassium, corn starch, corn syrup, FD&C Blue 1, FD&C Red 40, glycerin, medium chain triglycerides, natural and artificial flavors, propylene glycol, soybean oil, sucrose and water.

Ouestions? 1-888-287-1915



Benzocaine 7.5 mg Dextromethorphan HBr 5 mg





hotion guaranteed - Or well ce it or give you your money For guestions or comment



18 Lozenges

EXTRA STRENGTH

- · Relieves sore throat pain
- Quiets cough
- Maximum numbing[†]
- Take every 4 hours

Gluten-Free

WARNING: TAMPER-EVIDENT PACKAGING. DO NOT USE IF OUTER BOX OR BLISTER IS OPEN OR DAMAGED.

3161-785-888-1 SanoitzeuD

Inactive ingredients Acesuliame potsasium, com aterch, com syrup, FD&C Blue 1, FD&C Red 40, glycerin, medium chain trighycerides, natural and artificial flavors, propylene glycol, soybean oil, aucrose and water.

Other information a store at room temperature protect from moisture

Mixed Berry

†based on amount of benzocalne per adult dose

Drug Facts (continued)

No link or Coating

z eau for de	children under 6 years of age
take 1 lozenge and allow to dissolve slowly in the mouth; may be repeated every 4 hours, not to exceed 6 lozenges in 24 hours, or as directed by a doctor	children 6 to 12 years of age
Allow 2 lozenges to dissolve slowly in mouth, one at a time; may be repeated every 4 hours, not to exceed 12 lozenges in 24 hours, or as directed by a doctor	

if pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children, in case of overdose, get medical help or contact Poison Control Center right away.

These could be signs of a serious illness

Stop use and sak a doctor if some mouth symptoms do not improve in 7 days. Intitation, pain, or redness persists or worsens swelling develops a cough leats more than 7 days, comes back, or occurs with fever, rash, or persistent headsche.

condu rust occnis with too much bhiedm (muchs)

Ask a doctor before use if you have: persistent or chronic cough such as occurs with smoking, asthms, or emphysema

■ for teething ■ in children under 2 years of age

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional contitions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not forour prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

benzocelne, or other "calne" anesthetics.

Sore threat warming: If sore threat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nauses, or vomiting, consult a doctor promptly, intration, pain, or redness persist or worsens, see your dentist or doctor promptly.

Allergy slevt: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine,

Weamings

Methopolopinemia warning:

Methopolopinemia

Me

Orug Facts (continued)

No luk or Coading

EQUATE BENZ-DEX COUGH AND THROAT LOZENGES

benzocaine lozenge

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:79903-042

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength		
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	7.5 mg		
	DEXTROMETHORPHAN HYDROBROMIDE	5 mg		

Inactive Ingredients			
	Ingredient Name	Strength	
WATER (UNII: 059QF0KO0R)			

Product Characteristics			
Color	red (Dark Red)	Score	no score
Shape	ROUND	Size	17mm
Flavor	BERRY	Imprint Code	В
Contains			

ı	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l		IDC:79903- 142-18	18 in 1 BLISTER PACK; Type 0: Not a Combination Product	04/23/2021	



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/23/2021	

Labeler - Walmart (051957769)

Registrant - Bestco LLC (002149136)

Establishment			
Name	Address	ID/FEI	Business Operations
Bestco LLC		002149136	manufacture(79903-042)

Revised: 12/2024 Walmart