

MAXIMUM STRENGTH COLD AND FLU- acetaminophen,dextromethorphan hydrobromide,guaifenesin,phenylephrine hydrochloride capsule, liquid filled CVS Pharmacy, Inc

628T CVS 69842 789 ACETAMINOPHEN,DEXTROMETHORPHAN HYDROBROMIDE,GUAIFENESIN,PHENYLEPHRINE HYDROCHLORIDE 16CT

DRUG FACTS

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- Temporarily relieves these common cold and flu symptoms:

nasal congestion

headache

cough

minor aches and pains

sore throat

- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 softgels in 24 hrs, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product, do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 softgels in any 24-hour period
- adults and children 12 years of age and older: take 2 softgels every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

FD&C yellow no. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sorbitol sorbitan solution, titanium dioxide

Questions? call 1-877-290-4008

SAFETY SEALED: DO NOT USE IF THE INDIVIDUAL BLISTER UNIT IS OPEN OR TORN

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION



Maximum Strength[†]
COLD & FLU
Acetaminophen, 325 mg
Pain Reliever, Fever Reducer

Dextromethorphan HBr, 10 mg, Cough Suppressant
Guaifenesin, 200 mg, Expectorant
Phenylephrine HCl, 5 mg, Nasal Decongestant

Compare to Maximum Strength Mucinex[®] Fast-Max[®] Cold & Flu active ingredients*

Multi-Symptom[†]
Relieves: Sore throat, Sinus pressure, Sinus congestion, Headache, Fever, Cough, Nasal congestion, Body pain

16 SOFTGELS

Ages 12 & Over

†Under OTC monograph.

Directions or comments: Call 1-877-290-4008

Use only if you have a cold or flu. Do not use if you have a fever higher than 102°F (38.9°C) or if you have been told by a doctor not to take these medicines. Use only as directed. Total of 16 softgels in 24-hour period. Use only as directed.

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 12 softgels in 24 hours, which is the maximum daily amount; with other drugs containing acetaminophen; 3 or more alcoholic drinks daily while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, hives, rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI), certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: liver disease, heart disease, diabetes, high blood pressure, thyroid disease, trouble urinating due to an enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, cough that occurs with too much phlegm (mucus).

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed: Stop use and ask a doctor if: nervousness, dizziness, or sleepiness occur; pain, nasal congestion, or cough gets worse or lasts more than 7 days; persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema; fever gets worse or lasts more than 3 days; redness or swelling is present; new symptoms occur; cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

LOT: _____

EXP.: _____

Print/Varnish Omit Area

Varnish Omit Area

Varnish Omit Area

MAXIMUM STRENGTH COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-789
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
GELATIN (UNII: 2G86QN327L)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SHELLAC (UNII: 46N107B71O)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	26mm
Flavor		Imprint Code	78
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-789-16	1 in 1 CARTON	07/10/2021	
1		8 in 1 BLISTER PACK		
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/10/2021	

Labeler - CVS Pharmacy, Inc (062312574)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(69842-789)

Revised: 3/2026

CVS Pharmacy, Inc