

ASPIRIN- aspirin tablet, film coated
H E B

HEB 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal Display Panel

**Compare to Bayer® Genuine
Aspirin** active ingredient†

NDC 37808-157-12

H-E-B®

Aspirin
325 mg (**NSAID**)

Pain Reliever/Fever Reducer

Safe Pain Relief

- Headaches
- Aches & Pains

actual
size

100 COATED TABLETS

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

100% GUARANTEE
promise

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Genuine Aspirin.

**MADE WITH PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204**

50844 REV0122A15712

Compare to Bayer® Genuine Aspirin active ingredient†
NDC 37808-157-12

H-E-B

Aspirin

325 mg (NSAID)

Pain Reliever/Fever Reducer

Safe Pain Relief

- Headaches
- Aches & Pains



actual size

100 COATED TABLETS

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Genuine Aspirin.
MADE WITH PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204

Drug Facts

Active ingredient (in each tablet)
Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses Temporarily relieves: ■ minor pain of arthritis ■ headache ■ muscle pain ■ menstrual pain ■ toothache ■ pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product

Directions ■ do not take more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

Other information ■ use by expiration date on package ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391

50844 REV0122A15712
34910 -2207
100% GUARANTEE
0 41220 45703 8
PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

product ■ are age 60 or older

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you are taking a diuretic ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

Other information ■ use by expiration date on package ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391

STOP PEELING.

Adhesive Only

HEB 157

ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-157
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength

STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44;157
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-157-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
2	NDC:37808-157-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
3	NDC:37808-157-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/10/1996	

Labeler - H E B (007924756)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(37808-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(37808-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(37808-157)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(37808-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(37808-157)

Revised: 8/2025

H E B