

**TRAMADOL HYDROCHLORIDE EXTENDED-RELEASE- tramadol  
hydrochloride capsule  
Trigen Laboratories, LLC**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use Tramadol Hydrochloride Extended-Release Capsules safely and effectively. See full prescribing information for Tramadol Hydrochloride Extended-Release Capsules.

**Tramadol Hydrochloride Extended-Release Capsules for oral use, CIV**

**Initial U.S. Approval: 1995**

**WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF TRAMADOL  
HYDROCHLORIDE EXTENDED-RELEASE CAPSULES**

*See full prescribing information for complete boxed warning.*

- **Tramadol Hydrochloride Extended-Release Capsules exposes users to risks of addiction, abuse, and misuse, which can lead to overdose and death. Assess patient's risk before prescribing and reassess regularly for these behaviors and conditions. (5.1)**
- **Serious, life-threatening, or fatal respiratory depression may occur, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration are essential. Instruct patients to swallow Tramadol Hydrochloride Extended-Release Capsules intact, and not to split, chew, crush, or dissolve content of the capsules to avoid exposure to a potentially fatal dose of tramadol. (2.1, 5.2)**
- **Accidental ingestion of Tramadol Hydrochloride Extended-Release Capsules, especially by children, can result in a fatal overdose of tramadol. (5.2)**
- **Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing for use in patients for whom alternative treatment options are inadequate. (5.3, 7)**
- **Advise pregnant women using opioids for an extended period of time of the risk of Neonatal Opioid Withdrawal Syndrome, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery. (5.4)**
- **Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription. (5.5)**
- **Tramadol Hydrochloride Extended-Release Capsules are contraindicated in children younger than 12 years of age and in children younger than 18 years of age following tonsillectomy and/or adenoidectomy. (4) Avoid the use of Tramadol Hydrochloride Extended-Release Capsules in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of tramadol. (5.6)**
- **The effects of concomitant use or discontinuation of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with tramadol are complex. Use of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with Tramadol Hydrochloride Extended-Release Capsules requires careful consideration of the effects on the parent drug, tramadol, and the active metabolite, M1. (5.7, 7)**

----- **RECENT MAJOR CHANGES** -----

Boxed Warning	12/2025
Indications and Usage (1)	12/2025
Dosage and Administration (2.2, 2.3, 2.5)	12/2025

**INDICATIONS AND USAGE**

Tramadol Hydrochloride Extended-Release Capsules are an opioid agonist indicated for the management of severe and persistent pain that requires an opioid analgesic and that cannot be adequately treated with alternative options, including immediate-release opioids.

**Limitations of Use**

- Because of the risks of addiction, abuse, misuse, overdose, and death, which can occur at any dosage or duration and persist over the course of therapy, reserve opioid analgesics, including Tramadol Hydrochloride Extended-Release Capsules, for use in patients for whom alternative treatment options are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain. (1, 5.1)
- Tramadol Hydrochloride Extended-Release Capsules are not indicated as an as-needed (prn) analgesic. (1)

**DOSAGE AND ADMINISTRATION**

- Tramadol Hydrochloride Extended-Release Capsules should be prescribed only by healthcare professionals who are knowledgeable about the use of extended-release/long-acting opioids and how to mitigate the associated risks. (2.1)
- Use the lowest effective dosage for the shortest duration of time consistent with individual patient treatment goals. Reserve titration to higher doses of Tramadol Hydrochloride Extended-Release Capsules for patients in whom lower doses are insufficiently effective and in whom the expected benefits of using a higher dose opioid clearly outweigh the substantial risks. (2, 5)
- Initiate the dosing regimen for each patient individually, taking into account the patient's underlying cause and severity of pain, prior analgesic treatment and response, and risk factors for addiction, abuse, and misuse. (2.1, 5.1)
- Respiratory depression can occur at any time during opioid therapy, especially when initiating and following dosage increases with Tramadol Hydrochloride Extended-Release Capsules. Consider this risk when selecting an initial dose and when making dose adjustments. (2.1, 5.2)
- Tramadol Hydrochloride Extended-Release Capsules are administered orally once daily. (2.1)
- Discuss opioid overdose reversal agents and options for acquiring them with the patient and/or caregiver, both when initiating and renewing treatment with Tramadol Hydrochloride Extended-Release Capsules, especially if the patient has additional risk factors for overdose, or close contacts at risk for exposure and overdose. (2.2, 5.1, 5.2, 5.3)
- For patients currently on tramadol IR: Calculate total 24-hr IR dose, and initiate Tramadol Hydrochloride Extended-Release Capsules at a dose rounded down to next lower 100 mg increment; then adjust dose according to need and tolerance. See full prescribing information for instructions on conversion, titration, and maintenance of therapy. (2.3, 2.4)
- For patients converting from other opioid analgesics: Discontinue all other opioid analgesics other than as needed for breakthrough pain and initiate Tramadol Hydrochloride Extended-Release Capsules at a dose of 100 mg once daily, then titrate up by 100 mg increments every 5 days according to need and tolerance. (2.3, 2.4)
- Do not exceed a daily dose of 300 mg tramadol. Do not use with other tramadol products. (2.4)
- Periodically reassess patients receiving CONZIP to evaluate the continued need for opioid analgesics to maintain pain control, for the signs or symptoms of adverse reactions, and for the development of addiction, abuse, or misuse. (2.4)
- Do not rapidly reduce or abruptly discontinue Tramadol Hydrochloride Extended-Release Capsules in a physically-dependent patient because rapid reduction or abrupt discontinuation of opioid analgesics has resulted in serious withdrawal symptoms, uncontrolled pain, and suicide. (2.5, 5.18)

**DOSAGE FORMS AND STRENGTHS**

Extended-release capsules: 100 mg, 200 mg, and 300 mg (3)

**CONTRAINDICATIONS**

- Children younger than 12 years of age. (4)
- Postoperative management in children younger than 18 years of age following tonsillectomy and/or adenoidectomy. (4)
- Significant respiratory depression. (4)
- Acute or severe bronchial asthma in an unmonitored setting or in absence of resuscitative equipment. (4)
- Known or suspected gastrointestinal obstruction, including paralytic ileus. (4)
- Hypersensitivity to tramadol. (4)

- Concurrent use of monoamine oxidase inhibitors (MAOIs) or use within the last 14 days. (4)

### ----- **WARNINGS AND PRECAUTIONS** -----

- Opioid-Induced Hyperalgesia and Allodynia: Opioid-Induced Hyperalgesia (OIH) occurs when an opioid analgesic paradoxically causes an increase in pain, or an increase in sensitivity to pain. If OIH is suspected, carefully consider appropriately decreasing the dose of the current opioid analgesic or opioid rotation. (5.8)
- Serotonin Syndrome Risk: Potentially life-threatening condition could result from use of Tramadol Hydrochloride Extended-Release Capsules, particularly during concomitant use of serotonergic drugs. (5.9)
- Increased Risk of Seizures: Present within recommended dosage range. Risk is increased with higher than recommended doses and concomitant use of SSRIs, SNRIs, anorectics, tricyclic antidepressants and other tricyclic compounds, other opioids, MAOIs, neuroleptics, other drugs that reduce seizure threshold, in patients with epilepsy or at risk for seizures. (5.10, 7)
- Suicide Risk: Do not use Tramadol Hydrochloride Extended-Release Capsules in suicidal or addiction-prone patients. Use with caution in those taking tranquilizers, antidepressants or abusing alcohol. (5.11)
- Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients: Regularly evaluate, particularly during initiation and titration. (5.12)
- Adrenal Insufficiency: If diagnosed, treat with physiologic replacement of corticosteroids, and wean patient off of the opioid. (5.13)
- Severe Hypotension: Regularly evaluate during dosage initiation and titration. Avoid use of Tramadol Hydrochloride Extended-Release Capsules in patients with circulatory shock. (5.14)
- Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness: Monitor for sedation and respiratory depression. Avoid use of Tramadol Hydrochloride Extended-Release Capsules in patients with impaired consciousness or coma. (5.15)

### ----- **ADVERSE REACTIONS** -----

Most common adverse reactions (incidence  $\geq 10\%$  and twice placebo) are nausea, constipation, dry mouth, somnolence, dizziness, and vomiting. (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Trigen Laboratories, LLC at 1-800-541-4802 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### ----- **DRUG INTERACTIONS** -----

Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics: Avoid use with Tramadol Hydrochloride Extended-Release Capsules because they may reduce analgesic effect of Tramadol Hydrochloride Extended-Release Capsules or precipitate withdrawal symptoms. (5.18, 7)

### ----- **USE IN SPECIFIC POPULATIONS** -----

- Pregnancy: May cause fetal harm. (8.1)
- Lactation: Breastfeeding not recommended. (8.2)
- Severe Hepatic or Renal Impairment: Use not recommended. (8.6, 8.7)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 12/2025**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF TRAMADOL HYDROCHLORIDE EXTENDED-RELEASE CAPSULES**

#### **1 INDICATIONS AND USAGE**

#### **2 DOSAGE AND ADMINISTRATION**

2.1 Important Dosage and Administration Instructions

2.2 Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose

2.3 Initial Dosage

2.4 Titration and Maintenance Therapy

2.5 Safe Reduction or Discontinuation of Tramadol Hydrochloride Extended-Release Capsules

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Addiction, Abuse, and Misuse

5.2 Life-Threatening Respiratory Depression

5.3 Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants

5.4 Neonatal Opioid Withdrawal Syndrome

5.5 Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

5.6 Ultra-Rapid Metabolism of Tramadol and Other Risk Factors for Life-Threatening Respiratory Depression in Children

5.7 Risks of Interactions with Drugs Affecting Cytochrome P450 Isoenzymes

5.8 Opioid-Induced Hyperalgesia and Allodynia

5.9 Serotonin Syndrome Risk

5.10 Increased Risk of Seizures

5.11 Suicide Risk

5.12 Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients

5.13 Adrenal Insufficiency

5.14 Severe Hypotension

5.15 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness

5.16 Risks of Gastrointestinal Complications

5.17 Anaphylaxis and Other Hypersensitivity Reactions

5.18 Withdrawal

5.19 Risks of Driving and Operating Machinery

5.20 Hyponatremia

5.21 Hypoglycemia

### **6 ADVERSE REACTIONS**

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

### **7 DRUG INTERACTIONS**

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

### **9 DRUG ABUSE AND DEPENDENCE**

9.1 Controlled Substance

9.2 Abuse

9.3 Dependence

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

**FULL PRESCRIBING INFORMATION**

## **WARNING: SERIOUS AND LIFE-THREATENING RISKS FROM USE OF TRAMADOL HYDROCHLORIDE EXTENDED-RELEASE CAPSULES**

### **Addiction, Abuse, and Misuse**

Because the use of Tramadol Hydrochloride Extended-Release Capsules exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death, assess each patient's risk prior to prescribing and reassess all patients regularly for the development of these behaviors and conditions [*see Warnings and Precautions (5.1)*].

### **Life-Threatening Respiratory Depression**

Serious, life-threatening, or fatal respiratory depression may occur with use of Tramadol Hydrochloride Extended-Release Capsules, especially during initiation or following a dosage increase. To reduce the risk of respiratory depression, proper dosing and titration of Tramadol Hydrochloride Extended-Release Capsules are essential. Instruct patients to swallow Tramadol Hydrochloride Extended-Release Capsules intact, and not to split, break, chew, crush, or dissolve the contents of the capsules to avoid exposure to a potentially fatal dose of tramadol. [*see Dosage and Administration (2.1), Warnings and Precautions (5.2)*].

### **Accidental Ingestion**

Accidental ingestion of even one dose of Tramadol Hydrochloride Extended-Release Capsules, especially by children, can result in a fatal overdose of tramadol [*see Warnings and Precautions (5.2)*].

### **Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants**

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing of Tramadol Hydrochloride Extended-Release Capsules and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate [*see Warnings and Precautions (5.3), Drug Interactions (7)*].

### **Neonatal Opioid Withdrawal Syndrome (NOWS)**

Advise pregnant women using opioids for an extended period of time of the risk of Neonatal Opioid Withdrawal Syndrome, which may be life-threatening if not recognized and treated. Ensure that management by neonatology experts will be available at delivery [*see Warnings and Precautions (5.4)*].

### **Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)**

Healthcare providers are strongly encouraged to complete a REMS-compliant education program and to counsel patients and caregivers on serious risks, safe use, and the importance of reading the Medication Guide with each prescription [*see Warnings and Precautions (5.5)*].

## **Ultra-Rapid Metabolism Of Tramadol And Other Risk Factors for Life-Threatening Respiratory Depression in Children**

**Life-threatening respiratory depression and death have occurred in children who received tramadol. Some of the reported cases occurred following tonsillectomy and/or adenoidectomy, and in at least one case, the child had evidence of being an ultra-rapid metabolizer of tramadol due to a CYP2D6 polymorphism [see *Warnings and Precautions (5.6)*]. Tramadol Hydrochloride Extended-Release Capsules are contraindicated in children younger than 12 years of age and in children younger than 18 years of age following tonsillectomy and/or adenoidectomy [see *Contraindications (4)*]. Avoid the use of Tramadol Hydrochloride Extended-Release Capsules in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of tramadol [see *Warnings and Precautions (5.6)*].**

### **Interactions with Drugs Affecting Cytochrome P450 Isoenzymes**

**The effects of concomitant use or discontinuation of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with tramadol are complex. Use of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with Tramadol Hydrochloride Extended-Release Capsules requires careful consideration of the effects on the parent drug, tramadol, and the active metabolite, M1 [see *Warnings and Precautions (5.7)*, *Drug Interactions (7)*].**

## **1 INDICATIONS AND USAGE**

Tramadol Hydrochloride Extended-Release Capsules are indicated for the management of severe and persistent pain that requires an opioid analgesic and that cannot be adequately treated with alternative options, including immediate-release opioids.

### **Limitation of Use**

- Because of the risks of addiction, abuse, misuse, overdose, and death, which can occur at any dosage or duration and persist over the course of therapy [see *Warnings and Precautions (5.1)*], reserve opioid analgesics, including Tramadol Hydrochloride Extended-Release Capsules, for use in patients for whom alternative treatment options are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Tramadol Hydrochloride Extended-Release Capsules are not indicated as an as-needed (prn) analgesic.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Important Dosage and Administration Instructions**

- Tramadol Hydrochloride Extended-Release Capsules should be prescribed only by healthcare professionals who are knowledgeable about the use of extended-release/long-acting opioids and how to mitigate the associated risks.
- Do not use Tramadol Hydrochloride Extended-Release Capsules concomitantly with

other tramadol products [see *Warnings and Precautions (5.7), (5.15)*].

- Do not administer Tramadol Hydrochloride Extended-Release Capsules at a dose exceeding 300 mg per day.
- Use the lowest effective dosage for the shortest duration of time consistent with individual patient treatment goals [see *Warnings and Precautions (5)*]. Because the risk of overdose increases as opioid doses increase, reserve titration to higher doses of Tramadol Hydrochloride Extended-Release Capsules for patients in whom lower doses are insufficiently effective and in whom the expected benefits of using a higher dose opioid clearly outweigh the substantial risks.
- Initiate the dosing regimen for each patient individually, taking into account the patient's underlying cause and severity of pain, prior analgesic treatment and response, and risk factors for addiction, abuse, and misuse [see *Warnings and Precautions (5.1)*].
- Respiratory depression can occur at any time during opioid therapy, especially when initiating and following dosage increases with Tramadol Hydrochloride Extended-Release Capsules. Consider this risk when selecting an initial dose and when making dose adjustments [see *Warnings and Precautions (5.2)*].
- Tramadol Hydrochloride Extended-Release Capsules are administered orally once daily.
- Instruct patients to swallow Tramadol Hydrochloride Extended-Release Capsules capsules whole, and to take it with liquid. Breaking, chewing, splitting, or dissolving Tramadol Hydrochloride Extended-Release Capsules will result in uncontrolled delivery of tramadol and can lead to overdose or death [see *Warnings and Precautions (5.1)*].
- Tramadol Hydrochloride Extended-Release Capsules may be taken without regard to food. It is recommended that Tramadol Hydrochloride Extended-Release Capsules be taken in a consistent manner [see *Clinical Pharmacology (12.3)*].

## **2.2 Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose**

Inform patients and caregivers about opioid overdose reversal agents (e.g., naloxone, nalmefene). Discuss the importance of having access to an opioid overdose reversal agent, especially if the patient has risk factors for overdose (e.g., concomitant use of CNS depressants, a history of opioid use disorder, or prior opioid overdose) or if there are household members (including children) or other close contacts at risk for accidental ingestion or opioid overdose. The presence of risk factors for overdose should not prevent the management of pain in any patient [see *Warnings and Precautions (5.1, 5.2, 5.3)*].

Discuss the options for obtaining an opioid overdose reversal agent (e.g., prescription, over-the-counter, or as part of a community-based program) [see *Warnings and Precautions (5.2)*].

There are important differences among the opioid overdose reversal agents, such as route of administration, product strength, approved patient age range, and pharmacokinetics. Be familiar with these differences, as outlined in the approved labeling for those products, prior to recommending or prescribing such an agent.

## **2.3 Initial Dosage**

It is safer to underestimate a patient's 24-hour tramadol requirements and provide rescue medication (e.g., immediate-release opioid) than to overestimate the 24-hour

tramadol dosage and manage an adverse reaction due to an overdose. While useful tables of opioid equivalents are readily available, there is inter-patient variability in the potency of opioid drugs and opioid formulations. Frequently reevaluate patients for signs and symptoms of opioid withdrawal and for signs of oversedation/toxicity after converting patients to Tramadol Hydrochloride Extended-Release Capsules.

### Use of Tramadol Hydrochloride Extended-Release Capsules

The initial dose of Tramadol Hydrochloride Extended-Release Capsules is 100 mg once daily.

### Patients Currently on Tramadol Immediate-Release (IR) Products

Calculate the 24-hour tramadol IR dose and initiate a total daily dose of Tramadol Hydrochloride Extended-Release Capsules rounded down to the next lowest 100 mg increment. The dose should be taken once daily. The dose may subsequently be individualized according to patient need.

Due to limitations in flexibility of dose selection with Tramadol Hydrochloride Extended-Release Capsules, some patients maintained on tramadol IR products may not be able to convert to Tramadol Hydrochloride Extended-Release Capsules.

### Conversion from Other Opioid Analgesics to Tramadol Hydrochloride Extended-Release Capsules

When Tramadol Hydrochloride Extended-Release Capsules therapy is initiated, discontinue all opioid analgesics other than those used on an as needed basis for breakthrough pain when appropriate. There are no established conversion ratios for conversion from other opioids to Tramadol Hydrochloride Extended-Release Capsules defined by clinical trials. Initiate dosing using Tramadol Hydrochloride Extended-Release Capsules 100 mg once a day.

## **2.4 Titration and Maintenance Therapy**

Individually titrate Tramadol Hydrochloride Extended-Release Capsules by 100 mg every five days to a dose that provides adequate analgesia and minimizes adverse reactions. The maximum daily dose of Tramadol Hydrochloride Extended-Release Capsules is 300 mg per day.

Continually reevaluate patients receiving Tramadol Hydrochloride Extended-Release Capsules to assess the maintenance of pain control, signs and symptoms of opioid withdrawal, and other adverse reactions, as well as to reassess for the development of addiction, abuse, or misuse [see *Warnings and Precautions (5.1, 5.18)*]. Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration. During use of opioid therapy for an extended period of time, periodically reassess the continued need for the use of opioid analgesics.

Patients who experience breakthrough pain may require a dosage adjustment of Tramadol Hydrochloride Extended-Release Capsules or may need rescue medication with an appropriate dose of an immediate-release analgesic.

If the level of pain increases after dosage stabilization, attempt to identify the source of increased pain before increasing the Tramadol Hydrochloride Extended-Release Capsules dosage. If after increasing the dosage, unacceptable opioid-related adverse

reactions are observed (including an increase in pain after a dosage increase), consider reducing the dosage [see *Warnings and Precautions (5)*]. Adjust the dosage to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

## **2.5 Safe Reduction or Discontinuation of Tramadol Hydrochloride Extended-Release Capsules**

Do not rapidly reduce or abruptly discontinue Tramadol Hydrochloride Extended-Release Capsules in patients who may be physically dependent on opioids. Rapid reduction or abrupt discontinuation of opioid analgesics in patients who are physically dependent on opioids has resulted in serious withdrawal symptoms, uncontrolled pain, and suicide. Rapid reduction or abrupt discontinuation has also been associated with attempts to find other sources of opioid analgesics, which may be confused with drug-seeking for abuse. Patients may also attempt to treat their pain or withdrawal symptoms with illicit opioids, such as heroin, and other substances.

When a decision has been made to decrease the dose or discontinue therapy in an opioid-dependent patient taking Tramadol Hydrochloride Extended-Release Capsules, there are a variety of factors that should be considered, including the total daily dose of opioid (including Tramadol Hydrochloride Extended-Release Capsules) the patient has been taking, the duration of treatment, the type of pain being treated, and the physical and psychological attributes of the patient. It is important to ensure ongoing care of the patient and to agree on an appropriate tapering schedule and follow-up plan so that patient and provider goals and expectations are clear and realistic. When opioid analgesics are being discontinued due to a suspected substance use disorder, evaluate and treat the patient, or refer for evaluation and treatment of the substance use disorder. Treatment should include evidence-based approaches, such as medication assisted treatment of opioid use disorder. Complex patients with comorbid pain and substance use disorders may benefit from referral to a specialist.

There are no standard opioid tapering schedules that are suitable for all patients. Good clinical practice dictates a patient-specific plan to taper the dose of the opioid gradually. For patients on Tramadol Hydrochloride Extended-Release Capsules who are physically opioid-dependent, initiate the taper by a small enough increment (e.g., no greater than 10% to 25% of the total daily dose) to avoid withdrawal symptoms, and proceed with dose-lowering at an interval of every 2 to 4 weeks. Patients who have been taking opioids for briefer periods of time may tolerate a more rapid taper.

It may be necessary to provide the patient with lower dosage strengths to accomplish a successful taper. Reassess the patient frequently to manage pain and withdrawal symptoms, should they emerge. Common withdrawal symptoms include restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other signs and symptoms also may develop, including irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate. If withdrawal symptoms arise, it may be necessary to pause the taper for a period of time or raise the dose of the opioid analgesic to the previous dose, and then proceed with a slower taper. In addition, evaluate patients for any changes in mood, emergence of suicidal thoughts, or use of other substances.

When managing patients taking opioid analgesics, particularly those who have been treated for an extended period of time and/or with high doses for chronic pain, ensure

that a multimodal approach to pain management, including mental health support (if needed), is in place prior to initiating an opioid analgesic taper. A multimodal approach to pain management may optimize the treatment of chronic pain, as well as assist with the successful tapering of the opioid analgesic [see *Warnings and Precautions (5.18)*, *Drug Abuse and Dependence (9.3)*].

### **3 DOSAGE FORMS AND STRENGTHS**

Extended-release capsules are available as:

100 mg Capsules: White capsule imprinted with blue ink "G 252" on cap and "100" between lines on the body

200 mg Capsules: White capsule imprinted with violet ink "G 253" on cap and "200" between lines on the body

300 mg Capsules: White capsule imprinted with red ink "G 254" on cap and "300" between lines on the body

### **4 CONTRAINDICATIONS**

Tramadol Hydrochloride Extended-Release Capsules are contraindicated for:

- All children younger than 12 years of age [see *Warnings and Precautions (5.6)*]
- Postoperative management in children younger than 18 years of age following tonsillectomy and/or adenoidectomy [see *Warnings and Precautions (5.6)*]

Tramadol Hydrochloride Extended-Release Capsules is also contraindicated in patients with:

- Significant respiratory depression [see *Warnings and Precautions (5.12)*]
- Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment [see *Warnings and Precautions (5.12)*]
- Known or suspected gastrointestinal obstruction, including paralytic ileus [see *Warnings and Precautions (5.16)*]
- Hypersensitivity to tramadol (e.g., anaphylaxis) [see *Warnings and Precautions (5.17)*, *Adverse Reactions (6)*]
- Concurrent use of monoamine oxidase inhibitors (MAOIs) or use within the last 14 days [see *Drug Interactions (7)*]

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Addiction, Abuse, and Misuse**

Tramadol Hydrochloride Extended-Release Capsules contain tramadol, a Schedule IV controlled substance. As an opioid, Tramadol Hydrochloride Extended-Release Capsules expose users to the risks of addiction, abuse and misuse.

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed Tramadol Hydrochloride Extended-Release Capsules. Addiction can occur at recommended dosages and if the drug is misused or abused. The risk of opioid-related overdose or overdose-related death is increased with higher opioid doses,

and this risk persists over the course of therapy. In postmarketing studies, addiction, abuse, misuse, and fatal and non-fatal opioid overdose were observed in patients with long-term opioid use [see *Adverse Reactions (6.2)*].

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing Tramadol Hydrochloride Extended-Release Capsules, and reassess all patients receiving Tramadol Hydrochloride Extended-Release Capsules for the development of these behaviors and conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the proper management of pain in any given patient. Patients at increased risk may be prescribed opioids such as Tramadol Hydrochloride Extended-Release Capsules but use in such patients necessitates intensive counseling about the risks and proper use of Tramadol Hydrochloride Extended-Release Capsules along with frequent reevaluation for signs of addiction, abuse, and misuse. Consider recommending or prescribing an opioid overdose reversal agent [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.2)*].

Abuse or misuse of Tramadol Hydrochloride Extended-Release Capsules by splitting, breaking, chewing, crushing, snorting, or injecting the dissolved product will result in the uncontrolled delivery of tramadol and can result in overdose and death [see *Overdosage (10)*].

Opioids are sought for nonmedical use and are subject to diversion from legitimate prescribed use. Consider these risks when prescribing or dispensing Tramadol Hydrochloride Extended-Release Capsules. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on careful storage of the drug during the course of treatment and the proper disposal of unused drug. Contact local state professional licensing board or state-controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

## **5.2 Life-Threatening Respiratory Depression**

Serious, life-threatening, or fatal respiratory depression has been reported with the use of opioids, even when used as recommended. Respiratory depression, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid overdose reversal agents, depending on the patient's clinical status [see *Overdosage (10)*]. Carbon dioxide (CO<sub>2</sub>) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of Tramadol Hydrochloride Extended-Release Capsules, the risk is greatest during the initiation of therapy or following a dosage increase.

To reduce the risk of respiratory depression, proper dosing and titration of Tramadol Hydrochloride Extended-Release Capsules are essential [see *Dosage and Administration (2)*]. Overestimating the Tramadol Hydrochloride Extended-Release Capsules dosage when converting patients from another opioid product can result in a fatal overdose with the first dose.

Accidental ingestion of even one dose of Tramadol Hydrochloride Extended-Release

Capsules, especially by children, can result in respiratory depression and death due to an overdose of tramadol.

Educate patients and caregivers on how to recognize respiratory depression and emphasize the importance of calling 911 or getting emergency medical help right away in the event of a known or suspected overdose.

Opioids can cause sleep-related breathing disorders including central sleep apnea (CSA) and sleep-related hypoxemia. Opioid use increases the risk of CSA in a dose-dependent fashion. In patients who present with CSA, consider decreasing the opioid dosage using best practices for opioid taper [see *Dosage and Administration (2.5)*].

### Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose

Inform patients and caregivers about opioid overdose reversal agents (e.g., naloxone, nalmefene). Discuss the importance of having access to an opioid overdose reversal agent, especially if the patient has risk factors for overdose (e.g., concomitant use of CNS depressants, a history of opioid use disorder, or prior opioid overdose) or if there are household members (including children) or other close contacts at risk for accidental ingestion or opioid overdose. The presence of risk factors for overdose should not prevent the management of pain in any patient [see *Warnings and Precautions (5.1, 5.3)*].

Discuss the options for obtaining an opioid overdose reversal agent (e.g., prescription, over-the-counter, or as part of a community-based program).

There are important differences among the opioid overdose reversal agents, such as route of administration, product strength, approved patient age range, and pharmacokinetics. Be familiar with these differences, as outlined in the approved labeling for those products, prior to recommending or prescribing such an agent.

Educate patients and caregivers on how to recognize respiratory depression, and how to use an opioid overdose reversal agent for the emergency treatment of opioid overdose. Emphasize the importance of calling 911 or getting emergency medical help, even if an opioid overdose reversal agent is administered [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.1, 5.3)*, *Overdosage (10)*].

### **5.3 Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of Tramadol Hydrochloride Extended-Release Capsules with benzodiazepines and/or other CNS depressants, including alcohol (e.g., non-benzodiazepine sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, gabapentinoids [gabapentin and pregabalin], and other opioids). Because of these risks, reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioid analgesics alone. Because of similar pharmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS depressant drugs with opioid analgesics [see *Drug Interactions (7)*].

If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Inform patients and caregivers of this potential interaction and educate them on the signs and symptoms of respiratory depression (including sedation). If concomitant use is warranted, consider recommending or prescribing an opioid overdose reversal agent [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.2)*, *Overdosage (10)*].

Advise both patients and caregivers about the risks of respiratory depression and sedation when Tramadol Hydrochloride Extended-Release Capsules are used with benzodiazepines or other CNS depressants (including alcohol and illicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressants have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of additional CNS depressants including alcohol and illicit drugs [see *Drug Interactions (7)*].

#### **5.4 Neonatal Opioid Withdrawal Syndrome**

Use of Tramadol Hydrochloride Extended-Release Capsules for an extended period of time during pregnancy can result in withdrawal in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. Observe newborns for signs of neonatal opioid withdrawal syndrome and manage accordingly. Advise pregnant women using opioids for an extended period of time of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see *Use in Specific Populations (8.1)*].

#### **5.5 Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)**

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a Risk Evaluation and Mitigation Strategy (REMS) for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to do all of the following:

- Complete a REMS-compliant education program offered by an accredited provider of continuing education (CE) or another education program that includes all the elements of the FDA Education Blueprint for Healthcare Providers Involved in the Management or Support of Patients with Pain.
- Discuss the safe use, serious risks, and proper storage and disposal of opioid analgesics with patients and/or their caregivers every time these medicines are prescribed. The Patient Counseling Guide (PCG) can be obtained at this link: [www.fda.gov/OpioidAnalgesicREMSPCG](http://www.fda.gov/OpioidAnalgesicREMSPCG).
- Emphasize to patients and their caregivers the importance of reading the Medication

Guide that they will receive from their pharmacist every time an opioid analgesic is dispensed to them.

- Consider using other tools to improve patient, household, and community safety, such as patient-prescriber agreements that reinforce patient-prescriber responsibilities.

To obtain further information on the opioid analgesic REMS and for a list of accredited REMS CME/CE, call 1-800-503-0784, or log on to [www.opioidanalgesicrems.com](http://www.opioidanalgesicrems.com). The FDA Blueprint can be found at [www.fda.gov/OpioidAnalgesicREMSBlueprint](http://www.fda.gov/OpioidAnalgesicREMSBlueprint).

## **5.6 Ultra-Rapid Metabolism of Tramadol and Other Risk Factors for Life-Threatening Respiratory Depression in Children**

Life-threatening respiratory depression and death have occurred in children who received tramadol. Tramadol and codeine are subject to variability in metabolism based upon CYP2D6 genotype (described below), which can lead to increased exposure to an active metabolite. Based upon postmarketing reports with tramadol or with codeine, children younger than 12 years of age may be more susceptible to the respiratory depressant effects of tramadol. Furthermore, children with obstructive sleep apnea who are treated with opioids for post-tonsillectomy and/or adenoidectomy pain may be particularly sensitive to their respiratory depressant effect. Because of the risk of life-threatening respiratory depression and death:

- Tramadol Hydrochloride Extended-Release Capsules is contraindicated for all children younger than 12 years of age [*see Contraindications (4)*].
- Tramadol Hydrochloride Extended-Release Capsules is contraindicated for post-operative management in pediatric patients younger than 18 years of age following tonsillectomy and/or adenoidectomy [*see Contraindications (4)*].
- Avoid the use of Tramadol Hydrochloride Extended-Release Capsules in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of tramadol unless the benefits outweigh the risks. Risk factors include conditions associated with hypoventilation, such as postoperative status, obstructive sleep apnea, obesity, severe pulmonary disease, neuromuscular disease, and concomitant use of other medications that cause respiratory depression.
- As with adults, when prescribing opioids for adolescents, healthcare providers should choose the lowest effective dose for the shortest period of time and inform patients and caregivers about these risks and the signs of opioid overdose [*see Use in Specific Populations (8.4), Overdosage (10)*].

### Nursing Mothers

Tramadol is subject to the same polymorphic metabolism as codeine, with ultra-rapid metabolizers of CYP2D6 substrates being potentially exposed to life-threatening levels of O-desmethyltramadol (M1). At least one death was reported in a nursing infant who was exposed to high levels of morphine in breast milk because the mother was an ultra-rapid metabolizer of codeine. A baby nursing from an ultra-rapid metabolizer mother taking Tramadol Hydrochloride Extended-Release Capsules could potentially be exposed to high levels of M1, and experience life-threatening respiratory depression. For this reason, breastfeeding is not recommended during treatment with Tramadol Hydrochloride Extended-Release Capsules [*see Use in Specific Populations (8.2)*].

### CYP2D6 Genetic Variability: Ultra-rapid metabolizer

---

Some individuals may be ultra-rapid metabolizers because of a specific CYP2D6 genotype (gene duplications denoted as \*1/\*1×N or \*1/\*2×N). The prevalence of this CYP2D6 phenotype varies widely and has been estimated at 1 to 10% for Whites (European, North American), 3 to 4% for Blacks (African Americans), 1 to 2% for East Asians (Chinese, Japanese, Korean), and may be greater than 10% in certain racial/ethnic groups (i.e., Oceanian, Northern African, Middle Eastern, Ashkenazi Jews, Puerto Rican).

These individuals convert tramadol into its active metabolite, O-desmethyltramadol (M1), more rapidly and completely than other people. This rapid conversion results in higher than expected serum M1 levels. Even at labeled dosage regimens, individuals who are ultra-rapid metabolizers may have life-threatening or fatal respiratory depression or experience signs of overdose (such as extreme sleepiness, confusion, or shallow breathing) [see *Overdosage (10)*]. Therefore, individuals who are ultra-rapid metabolizers should not use Tramadol Hydrochloride Extended-Release Capsules.

## **5.7 Risks of Interactions with Drugs Affecting Cytochrome P450 Isoenzymes**

The effects of concomitant use or discontinuation of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors on levels of tramadol and M1 from Tramadol Hydrochloride Extended-Release Capsules are complex. Use of cytochrome P450 3A4 inducers, 3A4 inhibitors, or 2D6 inhibitors with Tramadol Hydrochloride Extended-Release Capsules requires careful consideration of the effects on the parent drug, tramadol which is a weak serotonin and norepinephrine reuptake inhibitor and mu-opioid agonist, and the active metabolite, M1, which is more potent than tramadol in mu-opioid receptor binding [see *Drug Interactions (7)*].

### Risks of Concomitant Use or Discontinuation of Cytochrome P450 2D6 Inhibitors

The concomitant use of Tramadol Hydrochloride Extended-Release Capsules with all cytochrome P450 2D6 inhibitors (e.g., amiodarone, quinidine) may result in an increase in tramadol plasma levels and a decrease in the levels of the active metabolite, M1. A decrease in M1 exposure in patients who have developed physical dependence to tramadol, may result in signs and symptoms of opioid withdrawal and reduced efficacy. The effect of increased tramadol levels may be an increased risk for serious adverse events including seizures and serotonin syndrome.

Discontinuation of a concomitantly used cytochrome P450 2D6 inhibitor may result in a decrease in tramadol plasma levels and an increase in active metabolite M1 levels, which could increase or prolong adverse reactions related to opioid toxicity and may cause potentially fatal respiratory depression.

Evaluate patients receiving Tramadol Hydrochloride Extended-Release Capsules and any CYP2D6 inhibitor at frequent intervals for the risk of serious adverse events including seizures and serotonin syndrome, signs and symptoms that may reflect opioid toxicity, and opioid withdrawal when Tramadol Hydrochloride Extended-Release Capsules are used in conjunction with inhibitors of CYP2D6 [see *Drug Interactions (7)*].

### Cytochrome P450 3A4 Interaction

The concomitant use of Tramadol Hydrochloride Extended-Release Capsules with cytochrome P450 3A4 inhibitors, such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g., ketoconazole), and protease inhibitors (e.g., ritonavir) or discontinuation of a cytochrome P450 3A4 inducer such as rifampin, carbamazepine,

and phenytoin, may result in an increase in tramadol plasma concentrations, which could increase or prolong adverse reactions, increase the risk for serious adverse events including seizures and serotonin syndrome, and may cause potentially fatal respiratory depression.

The concomitant use of Tramadol Hydrochloride Extended-Release Capsules with all cytochrome P450 3A4 inducers or discontinuation of a cytochrome P450 3A4 inhibitor may result in lower tramadol levels. This may be associated with a decrease in efficacy, and in some patients, may result in signs and symptoms of opioid withdrawal.

Evaluate patients receiving Tramadol Hydrochloride Extended-Release Capsules and any CYP3A4 inhibitor or inducer at frequent intervals for the risk for serious adverse events including seizures and serotonin syndrome, signs and symptoms that may reflect opioid toxicity and opioid withdrawal when Tramadol Hydrochloride Extended-Release Capsules is used in conjunction with inhibitors and inducers of CYP3A4 [see *Drug Interactions (7)*].

### **5.8 Opioid-Induced Hyperalgesia and Allodynia**

Opioid-Induced Hyperalgesia (OIH) occurs when an opioid analgesic paradoxically causes an increase in pain, or an increase in sensitivity to pain. This condition differs from tolerance, which is the need for increasing doses of opioids to maintain a defined effect [see *Dependence (9.3)*]. Symptoms of OIH include (but may not be limited to) increased levels of pain upon opioid dosage increase, decreased levels of pain upon opioid dosage decrease, or pain from ordinarily non-painful stimuli (allodynia). These symptoms may suggest OIH only if there is no evidence of underlying disease progression, opioid tolerance, opioid withdrawal, or addictive behavior.

Cases of OIH have been reported, both with short-term and longer-term use of opioid analgesics. Though the mechanism of OIH is not fully understood, multiple biochemical pathways have been implicated. Medical literature suggests a strong biologic plausibility between opioid analgesics and OIH and allodynia. If a patient is suspected to be experiencing OIH, carefully consider appropriately decreasing the dose of the current opioid analgesic or opioid rotation (safely switching the patient to a different opioid moiety) [see *Dosage and Administration (2.5)*, *Warnings and Precautions (5.18)*].

### **5.9 Serotonin Syndrome Risk**

Cases of serotonin syndrome, a potentially life-threatening condition, have been reported with the use of tramadol products, including Tramadol Hydrochloride Extended-Release Capsules, particularly during concomitant use with serotonergic drugs. Serotonergic drugs include selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT<sub>3</sub> receptor antagonists, drugs that affect the serotonergic neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), certain muscle relaxants (i.e., cyclobenzaprine, metaxalone), and drugs that impair metabolism of serotonin (including monoamine oxidase inhibitors, both those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue) [see *Drug Interactions (7)*]. This may occur within the recommended dosage range.

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination, rigidity), and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). The onset of

symptoms generally occurs within several hours to a few days of concomitant use but may occur later than that. Discontinue Tramadol Hydrochloride Extended-Release Capsules if serotonin syndrome is suspected.

### **5.10 Increased Risk of Seizures**

Seizures have been reported in patients receiving tramadol within the recommended dosage range. Spontaneous post-marketing reports indicate that seizure risk is increased with doses of tramadol above the recommended range.

Concomitant use of tramadol increases the seizure risk in patients taking: *[see Drug Interactions (7)]*

- Selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) antidepressants or anorectics,
- Tricyclic antidepressants (TCAs), and other tricyclic compounds (e.g., cyclobenzaprine, promethazine, etc.),
- Other opioids,
- Monoamine oxidase inhibitors (MAOIs) *[see Warnings and Precautions (5.9), Drug Interactions (7)]*,
- Neuroleptics, or
- Other drugs that reduce the seizure threshold.

Risk of seizures may also increase in patients with epilepsy, those with a history of seizures, or in patients with a recognized risk for seizure (such as head trauma, metabolic disorders, alcohol and drug withdrawal, CNS infections).

In tramadol overdose, administration of an opioid overdose reversal agent (e.g., naloxone or nalmefene) may increase the risk of seizure.

### **5.11 Suicide Risk**

- Do not prescribe Tramadol Hydrochloride Extended-Release Capsules for patients who are suicidal or addiction-prone. Consideration should be given to the use of non-narcotic analgesics in patients who are suicidal or depressed *[see Drug Abuse and Dependence (9.2)]*.
- Prescribe Tramadol Hydrochloride Extended-Release Capsules with caution for patients with a history of misuse and/or who are currently taking CNS-active drugs including tranquilizers or antidepressant drugs, or alcohol in excess, and patients who suffer from emotional disturbance or depression *[see Drug Interactions (7)]*.
- Inform patients not to exceed the recommended dose and to limit their intake of alcohol *[see Dosage and Administration (2.1), Warnings and Precautions (5.3)]*.

### **5.12 Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients**

The use of Tramadol Hydrochloride Extended-Release Capsules in patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment is contraindicated.

Patients with Chronic Pulmonary Disease: Tramadol Hydrochloride Extended-Release Capsules treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased

respiratory drive including apnea, even at recommended dosages of Tramadol Hydrochloride Extended-Release Capsules [see *Warnings and Precautions (5.2)*].

Elderly, Cachectic, or Debilitated Patients: Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics or altered clearance compared to younger, healthier patients [see *Warnings and Precautions (5.2)*].

Regularly evaluate patients, particularly when initiating and titrating Tramadol Hydrochloride Extended-Release Capsules and when Tramadol Hydrochloride Extended-Release Capsules are given concomitantly with other drugs that depress respiration [see *Warnings and Precautions (5.2, 5.7)*]. Alternatively, consider the use of non-opioid analgesics in these patients.

### **5.13 Adrenal Insufficiency**

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

### **5.14 Severe Hypotension**

Tramadol Hydrochloride Extended-Release Capsules may cause severe hypotension including orthostatic hypotension and syncope in ambulatory patients. There is increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume or concurrent administration of certain CNS depressant drugs (e.g., phenothiazines or general anesthetics) [see *Drug Interactions (7)*]. Regularly evaluate these patients for signs of hypotension after initiating or titrating the dosage of Tramadol Hydrochloride Extended-Release Capsules. In patients with circulatory shock, Tramadol Hydrochloride Extended-Release Capsules may cause vasodilation that can further reduce cardiac output and blood pressure. Avoid the use of Tramadol Hydrochloride Extended-Release Capsules in patients with circulatory shock.

### **5.15 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness**

In patients who may be susceptible to the intracranial effects of CO<sub>2</sub> retention (e.g., those with evidence of increased intracranial pressure or brain tumors), Tramadol Hydrochloride Extended-Release Capsules may reduce respiratory drive, and the resultant CO<sub>2</sub> retention can further increase intracranial pressure. Monitor such patients for signs of sedation and respiratory depression, particularly when initiating therapy with Tramadol Hydrochloride Extended-Release Capsules.

Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of Tramadol Hydrochloride Extended-Release Capsules in patients with impaired

consciousness or coma.

### **5.16 Risks of Gastrointestinal Complications**

Tramadol Hydrochloride Extended-Release Capsules is contraindicated in patients with known or suspected gastrointestinal obstruction, including paralytic ileus.

The tramadol in Tramadol Hydrochloride Extended-Release Capsules may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Regularly evaluate patients with biliary tract disease, including acute pancreatitis, for worsening symptoms.

Cases of opioid-induced esophageal dysfunction (OIED) have been reported in patients taking opioids. The risk of OIED may increase as the dose and/or duration of opioids increases. Regularly evaluate patients for signs and symptoms of OIED (e.g., dysphagia, regurgitation, non-cardiac chest pain), and if necessary, adjust opioid therapy as clinically appropriate. [see *Clinical Pharmacology (12.2)*].

### **5.17 Anaphylaxis and Other Hypersensitivity Reactions**

Serious and rarely fatal hypersensitive reactions have been reported in patients receiving therapy with tramadol. When these events do occur, it is often following the first dose. Other reported hypersensitivity reactions include pruritus, hives, bronchospasm, angioedema, toxic epidermal necrolysis and Stevens-Johnson syndrome. Patients with a history of hypersensitivity reactions to tramadol and other opioids may be at increased risk and therefore should not receive Tramadol Hydrochloride Extended-Release Capsules. If anaphylaxis or other hypersensitivity occurs, stop administration of Tramadol Hydrochloride Extended-Release Capsules immediately, discontinue Tramadol Hydrochloride Extended-Release Capsules permanently, and do not rechallenge with any formulation of tramadol. Advise patients to seek immediate medical attention if they experience any symptoms of a hypersensitivity reaction [see *Contraindications (4)*].

### **5.18 Withdrawal**

Do not rapidly reduce or abruptly discontinue Tramadol Hydrochloride Extended-Release Capsules in a patient physically-dependent on opioids. When discontinuing Tramadol Hydrochloride Extended-Release Capsules in a physically dependent patient, gradually taper the dosage. Rapid tapering of tramadol in a patient physically dependent on opioids may lead to a withdrawal syndrome and return of pain [see *Dosage and Administration (2.5)*, *Drug Abuse and Dependence (9.3)*].

Additionally, avoid the use of mixed agonist/antagonist (e.g., pentazocine, nalbuphine, and butorphanol) or partial agonist (e.g., buprenorphine) analgesics in patients who are receiving a full opioid agonist analgesic, including Tramadol Hydrochloride Extended-Release Capsules. In these patients, mixed agonist/antagonist and partial agonist analgesics may reduce the analgesic effect and/or may precipitate withdrawal symptoms [see *Drug Interactions (7)*].

When discontinuing Tramadol Hydrochloride Extended-Release Capsules, gradually taper the dosage [see *Dosage and Administration (2.5)*]. Do not abruptly discontinue Tramadol Hydrochloride Extended-Release Capsules [see *Drug Abuse and Dependence (9.3)*].

### **5.19 Risks of Driving and Operating Machinery**

Tramadol Hydrochloride Extended-Release Capsules may impair the mental or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to the effects of Tramadol Hydrochloride Extended-Release Capsules and know how they will react to the medication.

## 5.20 Hyponatremia

Hyponatremia (serum sodium <135 mmol/L) has been reported with the use of tramadol, and many cases are severe (sodium level <120 mmol/L). Most cases of hyponatremia occurred in females over the age of 65 and within the first week of therapy. In some reports, hyponatremia resulted from the syndrome of inappropriate antidiuretic hormone secretion (SIADH). Assess patients for signs and symptoms of hyponatremia (e.g., confusion, disorientation) during treatment with Tramadol Hydrochloride Extended-Release Capsules, especially during initiation of therapy. If signs and symptoms of hyponatremia are present, initiate appropriate treatment (e.g., fluid restriction) and discontinue Tramadol Hydrochloride Extended-Release Capsules [see *Dosage and Administration (2.5)*].

## 5.21 Hypoglycemia

Cases of tramadol-associated hypoglycemia have been reported, some resulting in hospitalization. In most cases, patients had predisposing risk factors (e.g., diabetes). If hypoglycemia is suspected, monitor blood glucose levels and consider drug discontinuation as appropriate [see *Dosage and Administration (2.5)*].

## 6 ADVERSE REACTIONS

The following serious or otherwise important adverse reactions are described in greater detail, in other sections:

- Addiction, Abuse, and Misuse [see *Warnings and Precautions (5.1)*]
- Life-Threatening Respiratory Depression [see *Warnings and Precautions (5.2)*]
- Interactions with Benzodiazepines and Other CNS Depressants [see *Warnings and Precautions (5.3)*]
- Neonatal Opioid Withdrawal Syndrome [see *Warnings and Precautions (5.4)*]
- Ultra-Rapid Metabolism of Tramadol and Other Risk Factors for Life-Threatening Respiratory Depression in Children [see *Warnings and Precautions (5.6)*]
- Opioid-Induced Hyperalgesia and Allodynia [see *Warnings and Precautions (5.8)*]
- Serotonin Syndrome [see *Warnings and Precautions (5.9)*]
- Seizures [see *Warnings and Precautions (5.10)*]
- Suicide [see *Warnings and Precautions (5.11)*]
- Adrenal Insufficiency [see *Warnings and Precautions (5.13)*]
- Severe Hypotension [see *Warnings and Precautions (5.14)*]
- Gastrointestinal Adverse Reactions [see *Warnings and Precautions (5.16)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.17)*]
- Withdrawal [see *Warnings and Precautions (5.18)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction

rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Tramadol Hydrochloride Extended-Release Capsules were administered to a total of 1987 patients in clinical trials. These included four double-blind and one long-term, open-label study in patients with osteoarthritis of the hip and knee. A total of 812 patients were 65 years or older. Adverse reactions with doses from 100 mg to 300 mg in the four pooled, randomized, double-blind, placebo-controlled studies in patients with chronic non-malignant pain are presented in the following table (see Table 1).

**Table 1: Incidence (%) of Patients with Adverse Reaction Rates  $\geq 5\%$  from Four Double-Blind, Placebo-Controlled Studies in Patients with Moderate to Moderately Severe Chronic Pain by Dose (N=1917)**

Preferred Term	Tramadol Hydrochloride Extended-Release Capsules			Placebo (N=646) n (%)
	100 mg (N=429) n (%)	200 mg (N=434) n (%)	300 mg (N=1054) n (%)	
Headache	99 (23.1)	96 (22.1)	200 (19.0)	128 (19.8)
Nausea	69 (16.1)	93 (21.4)	265 (25.1)	37 (5.7)
Somnolence	50 (11.7)	60 (13.8)	170 (16.1)	26 (4.0)
Dizziness	41 (9.6)	54 (12.4)	143 (13.6)	31 (4.8)
Constipation	40 (9.3)	59 (13.6)	225 (21.3)	27 (4.2)
Vomiting	28 (6.5)	45 (10.4)	98 (9.3)	12 (1.9)
Arthralgia	23 (5.4)	20 (4.6)	53 (5.0)	33 (5.1)
Dry Mouth	20 (4.7)	36 (8.3)	138 (13.1)	22 (3.4)
Sweating	18 (4.2)	23 (5.3)	71 (6.7)	4 (0.6)
Asthenia	15 (3.5)	26 (6.0)	91 (8.6)	17 (2.6)
Pruritus	13 (3.0)	25 (5.8)	77 (7.3)	12 (1.9)
Anorexia	9 (2.1)	23 (5.3)	60 (5.7)	1 (0.2)
Insomnia	9 (2.1)	9 (2.1)	53 (5.0)	11 (1.7)

The following adverse reactions were reported from all chronic pain studies (N=1917). The lists below include adverse reactions not otherwise noted in Table 1.

Adverse reactions with incidence rates of 1.0% to <5.0%

*Cardiac disorders:* hypertension

*Gastrointestinal disorders:* dyspepsia, flatulence

*General disorders:* abdominal pain, accidental injury, chills, fever, flu syndrome, neck pain, pelvic pain

*Investigations:* hyperglycemia, urine abnormality

*Metabolism and nutrition disorders:* peripheral edema, weight loss

*Musculoskeletal, connective tissue and bone disorders:* myalgia

*Nervous system disorders:* paresthesia, tremor, withdrawal syndrome

*Psychiatric disorders:* agitation, anxiety, apathy, confusion, depersonalization, depression, euphoria, nervousness

*Respiratory, thoracic and mediastinal disorders:* bronchitis, pharyngitis, rhinitis, sinusitis

*Skin and subcutaneous tissue disorders:* rash

*Urogenital disorders:* prostatic disorder, urinary tract infection

*Vascular disorders:* vasodilatation

Adverse reactions with incidence rates of 0.5% to <1.0% at any dose and serious adverse reactions reported in at least two patients

*Cardiac disorders:* EKG abnormal, hypotension, tachycardia

*Gastrointestinal disorders:* gastroenteritis

*General disorders:* neck rigidity, viral infection

*Hematologic/Lymphatic disorders:* anemia, ecchymoses

*Metabolism and nutrition disorders:* blood urea nitrogen increased, GGT increased, gout, SGPT increased

*Musculoskeletal disorders:* arthritis, arthrosis, joint disorder, leg cramps

*Nervous system disorders:* emotional lability, hyperkinesia, hypertonia, thinking abnormal, twitching, vertigo

*Respiratory disorders:* pneumonia

*Skin and subcutaneous tissue disorders:* hair disorder, skin disorder, urticaria

*Special Senses:* eye disorder, lacrimation disorder

*Urogenital disorders:* cystitis, dysuria, sexual function abnormality, urinary retention

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during post approval use of tramadol. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Serotonin syndrome: Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs.

Adrenal insufficiency: Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use.

Anaphylaxis: Anaphylaxis has been reported with ingredients contained in Tramadol Hydrochloride Extended-Release Capsules.

Androgen deficiency: Cases of androgen deficiency have occurred with use of opioids for an extended period of time [see *Clinical Pharmacology (12.2)*].

Hyperalgesia and Allodynia: Cases of hyperalgesia and allodynia have been reported with

opioid therapy of any duration [see *Warnings and Precautions (5.8)*].

QT prolongation/torsade de pointes: Cases of QT prolongation and/or torsade de pointes have been reported with tramadol use. Many of these cases were reported in patients taking another drug labeled for QT prolongation, in patients with a risk factor for QT prolongation (e.g., hypokalemia), or in overdose setting.

#### Metabolism and nutrition disorders

Hyponatremia: Cases of severe hyponatremia and/or SIADH have been reported in patients taking tramadol, most often in females over the age of 65, and within the first week of therapy [see *Warnings and Precautions (5.20)*].

Hypoglycemia: Cases of hypoglycemia have been reported in patients taking tramadol. Most reports were in patients with predisposing risk factors, including diabetes or renal insufficiency, or in elderly patients [see *Warnings and Precautions (5.21)*].

Opioid-induced esophageal dysfunction (OIED): Cases of OIED have been reported in patients taking opioids, and may occur more frequently in patients taking higher doses of opioid, and/or in patients taking opioids longer term [see *Warnings and Precautions (5.16)*].

#### Adverse Reactions from Observational Studies

A prospective, observational cohort study estimated the risks of addiction, abuse, and misuse in patients initiating long-term use of Schedule II opioid analgesics between 2017 and 2021. Study participants included in one or more analyses had been enrolled in selected insurance plans or health systems for at least one year, were free of at least one outcome at baseline, completed a minimum number of follow-up assessments, and either: 1) filled multiple extended-release/long-acting opioid analgesic prescriptions during a 90 day period (n=978); or 2) filled any Schedule II opioid analgesic prescriptions covering at least 70 of 90 days (n=1,244). Those included also had no dispensing of the qualifying opioids in the previous 6 months.

Over 12 months:

- approximately 1% to 6% of participants across the two cohorts newly met criteria for addiction, as assessed with two validated interview-based measures of moderate-to-severe opioid use disorder based on Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5) criteria, and
- approximately 9% and 22% of participants across the two cohorts newly met criteria for prescription opioid abuse and misuse [defined in *Drug Abuse and Dependence (9.2)*], respectively, as measured with a validated self-reported instrument.

A retrospective, observational cohort study estimated the risk of opioid-involved overdose or opioid overdose-related death in patients with new long-term use of Schedule II opioid analgesics from 2006 through 2016 (n=220,249). Included patients had been enrolled in either one of two commercial insurance programs, one managed care program, or one Medicaid program for at least 9 months. New long-term use was defined as having Schedule II opioid analgesic prescriptions covering at least 70 days' supply over the 3 months prior to study entry and none during the preceding 6 months. Patients were excluded if they had an opioid-involved overdose in the 9 months prior to study entry. Overdose was measured using a validated medical code-based algorithm with linkage to the National Death Index database. The 5-year cumulative incidence estimates for opioid-involved overdose or opioid overdose-related death ranged from

approximately 1.5% to 4% across study sites, counting only the first event during follow-up. Approximately 17% of first opioid overdoses observed over the entire study period (5-11 years, depending on the study site) were fatal. Higher baseline opioid dose was the strongest and most consistent predictor of opioid-involved overdose or opioid overdose-related death. Study exclusion criteria may have selected patients at lower risk of overdose, and substantial loss to follow-up (approximately 80%) also may have biased estimates.

The risk estimates from the studies described above may not be generalizable to all patients receiving opioid analgesics, such as those with exposures shorter or longer than the duration evaluated in the studies.

## 7 DRUG INTERACTIONS

Table 2 includes clinically significant drug interactions with Tramadol Hydrochloride Extended-Release Capsules.

**Table 2: Clinically Significant Drug Interactions with Tramadol Hydrochloride Extended-Release Capsules**

<b>Inhibitors of CYP2D6</b>	
Clinical Impact:	<p>The concomitant use of Tramadol Hydrochloride Extended-Release Capsules and CYP2D6 inhibitors may result in an increase in the plasma concentration of tramadol and a decrease in the plasma concentration of M1, particularly when an inhibitor is added after a stable dose of Tramadol Hydrochloride Extended-Release Capsules is achieved. Since M1 is a more potent mu-opioid agonist, decreased M1 exposure could result in decreased therapeutic effects, and may result in signs and symptoms of opioid withdrawal in patients who had developed physical dependence to tramadol. Increased tramadol exposure can result in increased or prolonged therapeutic effects and increased risk for serious adverse events including seizures and serotonin syndrome.</p> <p>After stopping a CYP2D6 inhibitor, as the effects of the inhibitor decline, the tramadol plasma concentration will decrease and the M1 plasma concentration will increase which could increase or prolong therapeutic effects but also increase adverse reactions related to opioid toxicity, and may cause potentially fatal respiratory depression [see <i>Clinical Pharmacology (12.3)</i>].</p>
Intervention:	<p>If concomitant use of a CYP2D6 inhibitor is necessary, evaluate patients at frequent intervals for adverse reactions including opioid withdrawal, seizures, and serotonin syndrome.</p> <p>If a CYP2D6 inhibitor is discontinued, consider lowering Tramadol Hydrochloride Extended-Release Capsules dosage until stable drug effects are achieved. Evaluate</p>

	patients at frequent intervals for adverse events including respiratory depression and sedation.
Examples:	Quinidine, fluoxetine, paroxetine, and bupropion
<b>Inhibitors of CYP3A4</b>	
Clinical Impact:	The concomitant use of Tramadol Hydrochloride Extended-Release Capsules and CYP3A4 inhibitors can increase the plasma concentration of tramadol and may result in a greater amount of metabolism via CYP2D6 and greater levels of M1. Follow patients closely for increased risk of serious adverse events including seizures and serotonin syndrome, and adverse reactions related to opioid toxicity including potentially fatal respiratory depression, particularly when an inhibitor is added after a stable dose of Tramadol Hydrochloride Extended-Release Capsules is achieved. After stopping a CYP3A4 inhibitor, as the effects of the inhibitor decline, the tramadol plasma concentration will decrease [see <i>Clinical Pharmacology (12.3)</i> ], resulting in decreased opioid efficacy and possibly signs and symptoms of opioid withdrawal in patients who had developed physical dependence to tramadol.
Intervention:	If concomitant use is necessary, consider dosage reduction of Tramadol Hydrochloride Extended-Release Capsules until stable drug effects are achieved. Evaluate patients at frequent intervals for seizures and serotonin syndrome, and signs of respiratory depression and sedation. If a CYP3A4 inhibitor is discontinued, consider increasing the Tramadol Hydrochloride Extended-Release Capsules dosage until stable drug effects are achieved and evaluate patients for signs and symptoms of opioid withdrawal.
Examples:	Macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g., ketoconazole), protease inhibitors (e.g., ritonavir)
<b>CYP3A4 Inducers</b>	
Clinical Impact:	The concomitant use of Tramadol Hydrochloride Extended-Release Capsules and CYP3A4 inducers can decrease the plasma concentration of tramadol [see <i>Clinical Pharmacology (12.3)</i> ], resulting in decreased efficacy or onset of a withdrawal syndrome in patients who have developed physical dependence to tramadol [see <i>Warnings and Precautions (5.7)</i> ]. After stopping a CYP3A4 inducer, as the effects of the inducer decline, the tramadol plasma concentration will increase [see <i>Clinical Pharmacology (12.3)</i> ], which could increase or prolong both the therapeutic effects and adverse reactions and may cause seizures and serotonin syndrome, and potentially fatal respiratory

	depression.
Intervention:	<p>If concomitant use is necessary, consider increasing the Tramadol Hydrochloride Extended-Release Capsules dosage until stable drug effects are achieved. Evaluate patients for signs of opioid withdrawal.</p> <p>If a CYP3A4 inducer is discontinued, consider Tramadol Hydrochloride Extended-Release Capsules dosage reduction and evaluate patients at frequent intervals for seizures and serotonin syndrome, and signs of respiratory depression and sedation.</p> <p>Patients taking carbamazepine, a CYP3A4 inducer, may have a significantly reduced analgesic effect of tramadol. Because carbamazepine increases tramadol metabolism and because of the seizure risk associated with tramadol, concomitant administration of Tramadol Hydrochloride Extended-Release Capsules and carbamazepine is not recommended.</p>
Examples:	Rifampin, carbamazepine, phenytoin
<b>Benzodiazepines and Other Central Nervous System (CNS) Depressants</b>	
Clinical Impact:	Due to additive pharmacologic effect, the concomitant use of benzodiazepines or other CNS depressants, including alcohol, can increase the risk of hypotension, respiratory depression, profound sedation, coma, and death [ <i>see Warnings and Precautions (5.3)</i> ].
Intervention:	Reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate. Limit dosages and durations to the minimum required. Inform patients and caregivers of this potential interaction and educate them on the signs and symptoms of respiratory depression (including sedation). If concomitant use is warranted, consider recommending or prescribing an opioid overdose reversal agent [ <i>see Dosage and Administration (2.2), Warnings and Precautions (5.1, 5.2, 5.3)</i> ].
Examples:	Benzodiazepines and other sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, gabapentinoids (gabapentin or pregabalin), other opioids, alcohol
<b>Serotonergic Drugs</b>	
Clinical Impact:	The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system has resulted in serotonin syndrome [ <i>see Warnings and Precautions (5.9)</i> ].
Intervention:	If concomitant use is warranted, frequently evaluate the patient, particularly during treatment initiation and dose adjustment. Discontinue Tramadol Hydrochloride Extended-Release Capsules if serotonin syndrome is suspected.

Examples:	Selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT <sub>3</sub> receptor antagonists, drugs that affect the serotonin neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), certain muscle relaxants (i.e., cyclobenzaprine, metaxalone), monoamine oxidase inhibitors (those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue)
-----------	---

### **Monoamine Oxidase Inhibitors (MAOIs)**

Clinical Impact:	MAOI interactions with opioids may manifest as serotonin syndrome [see <i>Warnings and Precautions (5.9)</i> ] or opioid toxicity (e.g., respiratory depression, coma) [see <i>Warnings and Precautions (5.2)</i> ].
------------------	--

Intervention:	Do not use Tramadol Hydrochloride Extended-Release Capsules in patients taking MAOIs or within 14 days of stopping such treatment.
---------------	--

Examples:	Phenelzine, tranylcypromine, linezolid
-----------	--

### **Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics**

Clinical Impact:	May reduce the analgesic effect of Tramadol Hydrochloride Extended-Release Capsules and/or precipitate withdrawal symptoms.
------------------	---

Intervention:	Avoid concomitant use.
---------------	------------------------

Examples:	Butorphanol, nalbuphine, pentazocine, buprenorphine
-----------	---

### **Muscle Relaxants**

Clinical Impact:	Tramadol may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression.
------------------	--

Intervention:	Because respiratory depression may be greater than otherwise expected, decrease the dosage of Tramadol Hydrochloride Extended-Release Capsules and/or the muscle relaxant as necessary. Due to the risk of respiratory depression with concomitant use of skeletal muscle relaxants and opioids, consider recommending or prescribing an opioid overdose reversal agent [see <i>Dosage and Administration (2.2)</i> , <i>Warnings and Precautions (5.2)</i> ].
---------------	--

Examples:	Cyclobenzaprine, metaxalone.
-----------	------------------------------

### **Diuretics**

Clinical Impact:	Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone.
------------------	---

Intervention:	Evaluate patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage of the diuretic as needed.
---------------	--

### **Anticholinergic Drugs**

Clinical Impact:	The concomitant use of anticholinergic drugs may increase risk of urinary retention and/or severe
------------------	---

	constipation, which may lead to paralytic ileus.
Intervention:	Evaluate patients for signs of urinary retention or reduced gastric motility when Tramadol Hydrochloride Extended-Release Capsules is used concomitantly with anticholinergic drugs.
<b>Digoxin</b>	
Clinical Impact:	Post-marketing surveillance of tramadol has revealed rare reports of digoxin toxicity.
Intervention:	Evaluate patients at frequent intervals for signs of digoxin toxicity and adjust dosage of digoxin as needed.
<b>Warfarin</b>	
Clinical Impact:	Post-marketing surveillance of tramadol has revealed rare reports of alteration of warfarin effect, including elevation of prothrombin times.
Intervention:	Frequently reevaluate the prothrombin time of patients on warfarin for signs of an interaction and adjust the dosage of warfarin as needed.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Use of opioid analgesics for an extended period of time during pregnancy may cause neonatal opioid withdrawal syndrome [see *Warnings and Precautions (5.4)*]. Available data with Tramadol Hydrochloride Extended-Release Capsules in pregnant women are insufficient to inform a drug-associated risk for major birth defects and miscarriage.

In animal reproduction studies, tramadol administration during organogenesis decreased fetal weights and reduced ossification in mice, rats, and rabbits at 1.4, 0.6, and 3.6 times the maximum recommended human daily dosage (MRHD). Tramadol decreased pup body weight and increased pup mortality at 1.2 and 1.9 times the MRHD [see *Data*]. Based on animal data, advise pregnant women of the potential risk to a fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

#### Clinical Considerations

##### *Fetal/Neonatal Adverse Reactions*

Use of opioid analgesics for an extended period of time during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth. Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea, and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by

the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome and manage accordingly [*see Warnings and Precautions (5.4)*].

Neonatal seizures, neonatal withdrawal syndrome, fetal death and still birth have been reported with tramadol during post-approval use of tramadol immediate-release products.

### *Labor or Delivery*

Opioids cross the placenta and may produce respiratory depression and psychophysiological effects in neonates. An opioid overdose reversal agent, such as naloxone or nalmefene, must be available for reversal of opioid-induced respiratory depression in the neonate. Tramadol Hydrochloride Extended-Release Capsules is not recommended for use in pregnant women during or immediately prior to labor, when use of shorter-acting analgesics or other analgesic techniques are more appropriate. Opioid analgesics, including Tramadol Hydrochloride Extended-Release Capsules can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression.

Tramadol has been shown to cross the placenta. The mean ratio of serum tramadol in the umbilical veins compared to maternal veins was 0.83 for 40 women given tramadol during labor.

The effect of Tramadol Hydrochloride Extended-Release Capsules, if any, on the later growth, development, and functional maturation of the child is unknown.

### Data

#### *Animal Data*

Tramadol has been shown to be embryotoxic and fetotoxic in mice, (120 mg/kg), rats (25 mg/kg) and rabbits (75 mg/kg) at maternally toxic dosages but was not teratogenic at these dose levels. These doses on a mg/m<sup>2</sup> basis are 1.9, 0.8, and 4.9 times the maximum recommended human daily dosage (MRHD) for mouse, rat and rabbit, respectively.

No drug-related teratogenic effects were observed in progeny of mice (up to 140 mg/kg), rats (up to 80 mg/kg) or rabbits (up to 300 mg/kg) treated with tramadol by various routes. Embryo and fetal toxicity consisted primarily of decreased fetal weights, decreased skeletal ossification, and increased supernumerary ribs at maternally toxic dose levels. Transient delays in developmental or behavioral parameters were also seen in pups from rat dams allowed to deliver. Embryo and fetal lethality were reported only in one rabbit study at 300 mg/kg, a dose that would cause extreme maternal toxicity in the rabbit. The dosages listed for mouse, rat, and rabbit are 2.3, 2.6, and 19 times the MRHD, respectively.

Tramadol was evaluated in pre- and post-natal studies in rats. Progeny of dams receiving oral (gavage) dose levels of 50 mg/kg (1.6 times the MRHD) or greater had decreased weights, and pup survival was decreased early in lactation at 80 mg/kg (2.6 times the MRHD).

## **8.2 Lactation**

## Risk Summary

Tramadol Hydrochloride Extended-Release Capsules are not recommended for obstetrical preoperative medication or for post-delivery analgesia in nursing mothers because its safety in infants and newborns has not been studied.

Tramadol and its metabolite, O-desmethyltramadol (M1), are present in human milk. There is no information on the effects of the drug on the breastfed infant or the effects of the drug on milk production. The M1 metabolite is more potent than tramadol in mu-opioid receptor binding [see *Clinical Pharmacology (12.1)*]. Published studies have reported tramadol and M1 in colostrum with administration of tramadol to nursing mothers in the early post-partum period. Women who are ultra-rapid metabolizers of tramadol may have higher than expected serum levels of M1, potentially leading to higher levels of M1 in breast milk that can be dangerous in their breastfed infants. In women with normal tramadol metabolism, the amount of tramadol secreted into human milk is low and dose dependent. Because of the potential for serious adverse reactions, including excess sedation and respiratory depression in a breastfed infant, advise patients that breastfeeding is not recommended during treatment with Tramadol Hydrochloride Extended-Release Capsules.

## Clinical Considerations

Monitor infants exposed to Tramadol Hydrochloride Extended-Release Capsules through breast milk for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

## Data

Following a single IV 100 mg dose of tramadol, the cumulative excretion in breast milk within 16 hours post dose was 100 mcg of tramadol (0.1% of the maternal dose) and 27 mcg of M1.

## **8.3 Females and Males of Reproductive Potential**

### Infertility

Use of opioids for an extended period of time may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [see *Adverse Reactions (6.2)*, *Clinical Pharmacology (12.2)*, *Nonclinical Toxicology (13.1)*].

## **8.4 Pediatric Use**

The safety and effectiveness of Tramadol Hydrochloride Extended-Release Capsules in pediatric patients have not been established.

Life-threatening respiratory depression and death have occurred in children who received tramadol [see *Warnings and Precautions (5.6)*]. In some of the reported cases, these events followed tonsillectomy and/or adenoidectomy, and one of the children had evidence of being an ultra-rapid metabolizer of tramadol (i.e., multiple copies of the gene for cytochrome P450 isoenzyme 2D6). Children with sleep apnea may be particularly sensitive to the respiratory depressant effects of tramadol. Because of the risk of life-threatening respiratory depression and death:

- Tramadol Hydrochloride Extended-Release Capsules is contraindicated for all children

younger than 12 years of age [see *Contraindications (4)*].

- Tramadol Hydrochloride Extended-Release Capsules is contraindicated for post-operative management in pediatric patients younger than 18 years of age following tonsillectomy and/or adenoidectomy [see *Contraindications (4)*].
- Avoid the use of Tramadol Hydrochloride Extended-Release Capsules in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of tramadol unless the benefits outweigh the risks. Risk factors include conditions associated with hypoventilation, such as postoperative status, obstructive sleep apnea, obesity, severe pulmonary disease, neuromuscular disease, and concomitant use of other medications that cause respiratory depression [see *Warnings and Precautions (5.6)*].

## 8.5 Geriatric Use

Eight hundred and twelve elderly (65 years of age or older) subjects were exposed to Tramadol Hydrochloride Extended-Release Capsules in clinical trials. Of those subjects, two hundred and forty were 75 years of age and older. In general, higher incidence rates of adverse events were observed for patients older than 65 years of age compared with patients 65 years and younger, particularly for the following adverse events: nausea, constipation, somnolence, dizziness, dry mouth, vomiting, asthenia, pruritus, anorexia, sweating, fatigue, weakness, postural hypotension and dyspepsia. For this reason, Tramadol Hydrochloride Extended-Release Capsules should be used with great caution in patients older than 75 years of age [see *Dosage and Administration (2.3)*].

Respiratory depression is the chief risk for elderly patients treated with opioids and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were co-administered with other agents that depress respiration. Titrate the dosage of Tramadol Hydrochloride Extended-Release Capsules slowly in geriatric patients and frequently reevaluate the patient for signs of central nervous system and respiratory depression [see *Warnings and Precautions (5.12)*].

Tramadol is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to regularly evaluate renal function.

## 8.6 Hepatic Impairment

Metabolism of tramadol and M1 is reduced in patients with advanced cirrhosis of the liver. Tramadol Hydrochloride Extended-Release Capsules has not been studied in patients with hepatic impairment. The limited availability of dose strengths of Tramadol Hydrochloride Extended-Release Capsules does not permit the dosing flexibility required for safe use in patients with severe hepatic impairment (Child-Pugh Class C). Therefore, Tramadol Hydrochloride Extended-Release Capsules should not be used in patients with severe hepatic impairment [see *Clinical Pharmacology (12.3)*].

## 8.7 Renal Impairment

Tramadol Hydrochloride Extended-Release Capsules has not been studied in patients with renal impairment. Impaired renal function results in a decreased rate and extent of excretion of tramadol and its active metabolite, M1. The limited availability of dose

strengths of Tramadol Hydrochloride Extended-Release Capsules does not permit the dosing flexibility required for safe use in patients with severe renal impairment (Child-Pugh Class C). Therefore, Tramadol Hydrochloride Extended-Release Capsules should not be used in patients with severe renal impairment [see *Clinical Pharmacology (12.3)*].

## **9 DRUG ABUSE AND DEPENDENCE**

### **9.1 Controlled Substance**

Tramadol Hydrochloride Extended-Release Capsules contain tramadol, a Schedule IV controlled substance.

### **9.2 Abuse**

Tramadol Hydrochloride Extended-Release Capsules contain tramadol, a substance with potential for misuse and abuse, which can lead to the development of substance use disorder, including addiction [see *Warnings and Precautions (5.1)*].

Misuse is the intentional use, for therapeutic purposes, of a drug by an individual in a way other than prescribed by a healthcare provider or for whom it was not prescribed.

Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that may include a strong desire to take the drug, difficulties in controlling drug use (e.g., continuing drug use despite harmful consequences, giving a higher priority to drug use than other activities and obligations), and possible tolerance or physical dependence.

Misuse and abuse of Tramadol Hydrochloride Extended-Release Capsules increases risk of overdose, which may lead to central nervous system and respiratory depression, hypotension, seizures, and death. The risk is increased with concurrent abuse of Tramadol Hydrochloride Extended-Release Capsules with alcohol and/or other CNS depressants. Abuse of and addiction to opioids in some individuals may not be accompanied by concurrent tolerance and symptoms of physical dependence. In addition, abuse of opioids can occur in the absence of addiction.

All patients treated with opioids require careful and frequent reevaluation for signs of misuse, abuse, and addiction, because use of opioid analgesic products carries the risk of addiction even under appropriate medical use. Patients at high risk of Tramadol Hydrochloride Extended-Release Capsules abuse include those with a history of prolonged use of any opioid, including products containing tramadol, those with a history of drug or alcohol abuse, or those who use Tramadol Hydrochloride Extended-Release Capsules in combination with other abused drugs.

"Drug-seeking" behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing, or referral, repeated "loss" of prescriptions, tampering with prescriptions, and reluctance to provide prior medical records or contact information for other treating healthcare provider(s). "Doctor shopping" (visiting multiple prescribers to obtain additional prescriptions) is common among people who abuse drugs and people with substance use disorder. Preoccupation with achieving adequate pain relief can be appropriate behavior in a patient with

inadequate pain control.

Tramadol Hydrochloride Extended-Release Capsules, like other opioids, can be diverted for nonmedical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic reevaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

### Risks Specific to Abuse of Tramadol Hydrochloride Extended-Release Capsules

Abuse of Tramadol Hydrochloride Extended-Release Capsules poses a risk of overdose and death. This risk is increased with concurrent use of Tramadol Hydrochloride Extended-Release Capsules with alcohol and/or other CNS depressants [see *Warnings and Precautions (5.1, 5.3), Drug Interactions (7)*].

Tramadol Hydrochloride Extended-Release Capsules is approved for oral use only. With parental abuse, the inactive ingredients in Tramadol Hydrochloride Extended-Release Capsules can result in local tissue necrosis, infection, pulmonary granulomas, increased risk of endocarditis and valvular heart injury, embolism, and death.

Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

## **9.3 Dependence**

Both tolerance and physical dependence can develop during use of opioid therapy.

Tolerance is a physiological state characterized by a reduced response to a drug after repeated administration (i.e., a higher dose of a drug is required to produce the same effect that was once obtained at a lower dose).

Physical dependence is a state that develops as a result of a physiological adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discontinuation or a significant dose reduction of a drug.

Withdrawal may be precipitated through the administration of drugs with opioid antagonist activity (e.g., naloxone, nalmefene), mixed agonist/antagonist analgesics (e.g., pentazocine, butorphanol, nalbuphine), or partial agonists (e.g., buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued use.

Do not rapidly reduce or abruptly discontinue Tramadol Hydrochloride Extended-Release Capsules in a patient physically dependent on opioids. Rapid tapering of Tramadol Hydrochloride Extended-Release Capsules in a patient physically dependent on opioids may lead to serious withdrawal symptoms, uncontrolled pain, and suicide. Rapid discontinuation has also been associated with attempts to find other sources of opioid analgesics, which may be confused with drug-seeking for abuse.

When discontinuing Tramadol Hydrochloride Extended-Release Capsules, gradually taper the dosage using a patient-specific plan that considers the following: the dose of Tramadol Hydrochloride Extended-Release Capsules the patient has been taking, the duration of treatment, and the physical and psychological attributes of the patient. To improve the likelihood of a successful taper and minimize withdrawal symptoms, it is

important that the opioid tapering schedule is agreed upon by the patient. In patients taking opioids for an extended period of time at high doses, ensure that a multimodal approach to pain management, including mental health support (if needed), is in place prior to initiating an opioid analgesic taper [see *Dosage and Administration (2.5)*, *Warnings and Precautions (5.18)*].

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [see *Use in Specific Populations (8.1)*].

## **10 OVERDOSAGE**

### Clinical Presentation

Acute overdosage with tramadol can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, QT prolongation, hypotension, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations. [see *Clinical Pharmacology (12.2)*]. Toxic leukoencephalopathy has been reported after opioid overdose and can present hours, days, or weeks after apparent recovery from the initial intoxication.

### Treatment of Overdose

In case of overdosage, priorities are the reestablishment of a patent and protected airway and institution of assisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life-support measures.

For clinically significant respiratory or circulatory depression secondary to opioid overdose, administer an opioid overdose reversal agent such as naloxone or nalmefene.

While an opioid overdose reversal agent will reverse some, but not all, symptoms caused by overdosage with tramadol, the risk of seizures is also increased with reversal agent administration. In animals, convulsions following the administration of toxic doses of Tramadol Hydrochloride Extended-Release Capsules could be suppressed with barbiturates or benzodiazepines but were increased with naloxone. Naloxone administration did not change the lethality of an overdose in mice. Hemodialysis is not expected to be helpful in an overdose because it removes less than 7% of the administered dose in a 4-hour dialysis period.

Because the duration of opioid reversal is expected to be less than the duration of action of tramadol in Tramadol Hydrochloride Extended-Release Capsules, carefully monitor the patient until spontaneous respiration is reliably reestablished. Tramadol Hydrochloride Extended-Release Capsules will continue to release tramadol and add to the tramadol load for 24 to 48 hours or longer following ingestion, necessitating prolonged monitoring. If the response to an opioid overdose reversal agent is suboptimal or only brief in nature, administer additional reversal agent as directed by the product's prescribing information.


In an individual physically dependent on opioids, administration of the recommended

usual dosage of the opioid overdose reversal agent will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms experienced will depend on the degree of physical dependence and the dose of the reversal agent administered. If a decision is made to treat serious respiratory depression in the physically dependent patient, administration of the reversal agent should be initiated with care and by titration with smaller than usual doses of the reversal agent.

## 11 DESCRIPTION

Tramadol Hydrochloride Extended-Release Capsules are an opioid agonist in an extended-release oral formulation. The chemical name for tramadol hydrochloride USP is ( $\pm$ )*cis*-2-[(dimethylamino)methyl]-1-(3-methoxyphenyl) cyclohexanol hydrochloride. Its structural formula is:

**Figure 1**



The molecular weight of tramadol hydrochloride USP is 299.8. It is a white, bitter, crystalline and odorless powder that is readily soluble in water and ethanol and has a pKa of 9.41. The n-octanol/water log partition coefficient (logP) is 1.35 at pH 7.

Tramadol Hydrochloride Extended-Release Capsules contain a total dose of tramadol hydrochloride 100, 200, and 300 mg in a combination of immediate-release and extended-release components.

<b>Dosage</b>	<b>Immediate-release</b>	<b>Extended-release</b>
100 mg	25 mg	75 mg
200 mg	50 mg	150 mg
300 mg	50 mg	250 mg

Tramadol Hydrochloride Extended-Release Capsules are white in color. Inactive ingredients include gelatin, titanium dioxide, shellac, FD & C Blue #2 aluminum lake (E132) (100 and 200 mg capsules), D & C Red #7 calcium lake (E180) (200 and 300 mg capsules), D & C Yellow #10 aluminum lake (300 mg capsule), lactose monohydrate 200 mesh, microcrystalline cellulose, povidone K30, corn starch, sodium starch glycolate, magnesium stearate, sucrose stearate, hypromellose, talc, polysorbate 80, Eudragit NE

30D, and simethicone emulsion.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Tramadol Hydrochloride Extended-Release Capsules contain tramadol, an opioid agonist, and an inhibitor of reuptake of norepinephrine and serotonin. Although its mode of action is not completely understood, from animal tests, at least two complementary mechanisms appear applicable: binding of parent and M1 metabolite to mu-opioid receptors and weak inhibition of reuptake of norepinephrine and serotonin.

Opioid activity of tramadol is due to both low affinity binding of the parent compound and higher affinity binding of the O-demethylated metabolite M1 to mu-opioid receptors. In animal models, M1 is up to 6 times more potent than tramadol in producing analgesia and 200 times more potent in mu-opioid binding. Tramadol-induced analgesia is only partially antagonized by the opioid antagonist naloxone in several animal tests. The relative contribution of both tramadol and M1 to human analgesia is dependent upon the plasma concentrations of each compound.

Tramadol has been shown to inhibit reuptake of norepinephrine and serotonin in vitro, as have some other opioid analgesics. These mechanisms may contribute independently to the overall analgesic profile of tramadol. The relationship between exposure of tramadol and M1 and efficacy has not been evaluated in clinical studies.

Apart from analgesia, tramadol administration may produce a constellation of symptoms (including dizziness, somnolence, nausea, constipation, sweating and pruritus) similar to that of other opioids. In contrast to morphine, tramadol has not been shown to cause histamine release. At therapeutic doses, tramadol has no effect on heart rate, left ventricular function or cardiac index. Orthostatic hypotension has been observed.

### **12.2 Pharmacodynamics**

#### Effects on the Central Nervous System

Tramadol produces respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and electrical stimulation.

Tramadol causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar findings). Marked mydriasis rather than miosis may be seen due to hypoxia in overdose situations.

#### Effects on the Gastrointestinal Tract and Other Smooth Muscle

Tramadol causes a reduction in motility associated with an increase in smooth muscle tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone is increased to the point of spasm, resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, transient elevations in serum

amylase, and opioid-induced esophageal dysfunction (OIED).

### Effects on the Cardiovascular System

Tramadol produces peripheral vasodilation, which may result in orthostatic hypotension or syncope. Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes, sweating, and/or orthostatic hypotension.

The effect of oral tramadol on the QTcF interval was evaluated in a double-blind, randomized, four-way crossover, placebo- and positive- (moxifloxacin) controlled study in 68 adult male and female healthy subjects. At a 600 mg/day dose (1.5-fold the maximum immediate-release daily dose), the study demonstrated no significant effect on the QTcF interval.

### Effects on the Endocrine System

Opioids inhibit the secretion of adrenocorticotropic hormone (ACTH), cortisol, and luteinizing hormone (LH) in humans [see *Adverse Reactions (6.2)*]. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Use of opioids for an extended period of time may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see *Adverse Reactions (6.2)*].

### Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system in in vitro and animal models. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

### Concentration-Efficacy Relationships

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with opioid agonists. The minimum effective analgesic concentration of tramadol for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome, and/or the development of analgesic tolerance [see *Dosage and Administration (2.1, 2.4)*].

### Concentration-Adverse Reaction Relationships

There is a relationship between increasing tramadol plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions [see *Dosage and Administration (2.1, 2.3, 2.4)*].

## **12.3 Pharmacokinetics**

The analgesic activity of tramadol is due to both parent drug and the M1 metabolite. Tramadol Hydrochloride Extended-Release Capsules is administered as a racemate and both tramadol and M1 are detected in the circulation. The  $C_{max}$  and AUC of Tramadol Hydrochloride Extended-Release Capsules have been observed to be dose-proportional

over an oral dose range of 100 to 300 mg in healthy subjects.

### Absorption

After a single dose administration of Tramadol Hydrochloride Extended-Release Capsules,  $T_{max}$  occurs around 10-12 hours.

The mean  $C_{max}$  and AUC of Tramadol Hydrochloride Extended-Release Capsules after a 300 mg single dose was 308 ng/mL and 6777 ng.hr/mL, respectively under fasting conditions. Tramadol Hydrochloride Extended-Release Capsules are bioequivalent to a reference extended-release tramadol product following a single 300 mg dose under fasting conditions.

At steady-state, Tramadol Hydrochloride Extended-Release Capsules at 200 mg have been observed to be bioequivalent to a reference extended-release tramadol product at 200 mg under fasting conditions (Table 3). Following administration of Tramadol Hydrochloride Extended-Release Capsules 200 mg capsules, steady-state plasma concentrations of both tramadol and M1 are achieved within four days of once daily dosing.

**Table 3**

<b>Mean (%CV) Steady-State Pharmacokinetic Parameter Values (N= 38)</b>				
<b>Parameter</b>	<b>Tramadol</b>		<b>O-Desmethyltramadol (M1 Metabolite)</b>	
	<b>Tramadol Hydrochloride Extended-Release Capsules 200 mg</b>	<b>A Reference Extended-Release Tramadol Product 200 mg</b>	<b>Tramadol Hydrochloride Extended-Release Capsules 200 mg</b>	<b>A Reference Extended-Release Tramadol Product 200 mg</b>
AUC <sub>0-24</sub> (ng.hr/mL)	5678 (27%)	5563 (32%)	1319 (34%)	1302 (40%)
$C_{max}$ (ng/mL)	332 (25%)	350 (31%)	70 (34%)	74 (41%)
$C_{min}$ (ng/mL)	128 (39%)	125 (45%)	35 (34%)	33 (42%)
$T_{max}$	5.9 (66%)	10 (30%)	11 (37%)	13 (29%)
% Fluctuation	88 (19%)	101 (30%)	64 (22%)	76 (30%)

AUC<sub>0-24</sub>: Area Under the Curve in a 24-hour dosing interval

$C_{max}$ : Peak Concentration in a 24-hour dosing interval

$C_{min}$ : Trough Concentration in a 24-hour dosing interval

$T_{max}$ : Time to Peak Concentration

### *Food Effect*

The rate and extent of absorption of Tramadol Hydrochloride Extended-Release Capsules (300 mg) are similar following oral administration with or without food. Therefore, Tramadol Hydrochloride Extended-Release Capsules can be administered without regard to meals.

### Distribution

The volume of distribution of tramadol was 2.6 and 2.9 liters/kg in male and female subjects, respectively, following a 100 mg intravenous tramadol dose. The binding of tramadol to human plasma proteins is approximately 20% and binding also appears to be independent of concentration up to 10 mcg/mL. Saturation of plasma protein binding occurs only at concentrations outside the clinically relevant range.

### Elimination

Tramadol is eliminated primarily through metabolism by the liver and the metabolites are eliminated primarily by the kidneys. The mean plasma elimination half-lives of racemic tramadol and racemic M1 after administration of Tramadol Hydrochloride Extended-Release Capsules are approximately 10 and 11 hours, respectively.

### *Metabolism*

Tramadol is extensively metabolized after oral administration. The major metabolic pathways appear to be N- (mediated by CYP3A4 and CYP2B6) and O- (mediated by CYP2D6) demethylation and glucuronidation or sulfation in the liver. One metabolite (O-desmethyltramadol, denoted M1) is pharmacologically active in animal models. Formation of M1 is dependent on CYP2D6 and as such is subject to inhibition and polymorphism, which may affect the therapeutic response [see *Drug Interactions (7)*].

### *Excretion*

Approximately 30% of the dose is excreted in the urine as unchanged drug, whereas 60% of the dose is excreted as metabolites. The remainder is excreted either as unidentified or as unextractable metabolites.

### Special Populations

#### *Hepatic Impairment*

Pharmacokinetics of tramadol was studied in patients with mild or moderate hepatic impairment after receiving multiple doses of an extended-release tramadol product at 100 mg. The exposure of (+)- and (-)-tramadol was similar in mild and moderate hepatic impairment patients in comparison to patients with normal hepatic function. However, exposure of (+)- and (-)-M1 decreased ~50% with increased severity of the hepatic impairment (from normal to mild and moderate). The pharmacokinetics of tramadol has not been studied in patients with severe hepatic impairment. After the administration of tramadol immediate-release tablets to patients with advanced cirrhosis of the liver, tramadol area under the plasma concentration time curve was larger and the tramadol and M1 half-lives were longer than subjects with normal hepatic function. The limited availability of dose strengths of Tramadol Hydrochloride Extended-Release Capsules does not permit the dosing flexibility required for safe use in patients with severe hepatic impairment. Therefore, Tramadol Hydrochloride Extended-Release Capsules should not be used in patients with severe hepatic impairment [see *Use in Specific Populations (8.6)*].

#### *Renal Impairment*

Impaired renal function results in a decreased rate and extent of excretion of tramadol and its active metabolite, M1. The pharmacokinetics of tramadol was studied in patients with mild or moderate renal impairment after receiving multiple doses of an extended-release tramadol product at 100 mg. There is no consistent trend observed for tramadol exposure related to renal function in patients with mild ( $CL_{Cr}$ : 50-80 mL/min) or

moderate ( $CL_{Cr}$ : 30-50 mL/min) renal impairment in comparison to patients with normal renal function ( $CL_{Cr}$  > 80 mL/min). However, exposure of M1 increased 20-40% with increased severity of the renal impairment (from normal to mild and moderate). The pharmacokinetics of tramadol has not been studied in patients with severe renal impairment ( $CL_{Cr}$  < 30 mL/min). The limited availability of dose strengths of Tramadol Hydrochloride Extended-Release Capsules does not permit the dosing flexibility required for safe use in patients with severe renal impairment. Therefore, Tramadol Hydrochloride Extended-Release Capsules should not be used in patients with severe renal impairment. The total amount of tramadol and M1 removed during a 4-hour dialysis period is less than 7% of the administered dose [see *Use in Specific Populations (8.6)*].

## Sex

Based on pooled multiple-dose pharmacokinetics studies for an extended-release tramadol product in 166 healthy subjects (111 males and 55 females), the dose-normalized AUC values for tramadol were somewhat higher in females than in males. There was a considerable degree of overlap in values between male and female groups. Dosage adjustment based on sex is not recommended.

## Age: Geriatric Population

The effect of age on pharmacokinetics of Tramadol Hydrochloride Extended-Release Capsules has not been studied. Healthy elderly subjects aged 65 to 75 years administered an immediate-release formulation of tramadol, have plasma concentrations and elimination half-lives comparable to those observed in healthy subjects younger than 65 years of age. In subjects over 75 years, mean maximum plasma concentrations are elevated (208 vs. 162 ng/mL) and the mean elimination half-life is prolonged (7 vs. 6 hours) compared to subjects 65 to 75 years of age. Adjustment of the daily dose is recommended for patients older than 75 years [see *Dosage and Administration (2.3)*].

## Drug Interaction Studies

### *Potential for Tramadol to Affect Other Drugs*

In vitro studies indicate that tramadol is unlikely to inhibit the CYP3A4-mediated metabolism of other drugs when tramadol is administered concomitantly at therapeutic doses. Tramadol does not appear to induce its own metabolism in humans, since observed maximal plasma concentrations after multiple oral doses are higher than expected based on single-dose data.

### *Poor / Extensive Metabolizers, CYP2D6*

The formation of the active metabolite, M1, is mediated by CYP2D6, a polymorphic enzyme. Approximately 7% of the population has reduced activity of the CYP2D6 isoenzyme of cytochrome P-450 metabolizing enzyme system. These individuals are "poor metabolizers" of debrisoquine, dextromethorphan and tricyclic antidepressants, among other drugs. Based on a population PK analysis of Phase 1 studies with IR tablets in healthy subjects, concentrations of tramadol were approximately 20% higher in "poor metabolizers" versus "extensive metabolizers", while M1 concentrations were 40% lower.

### *CYP2D6 Inhibitors*

In vitro drug interaction studies in human liver microsomes indicate that concomitant administration with inhibitors of CYP2D6 such as fluoxetine, paroxetine, and amitriptyline could result in some inhibition of the metabolism of tramadol.

## *Quinidine*

Tramadol is metabolized to active metabolite M1 by CYP2D6. Coadministration of quinidine, a selective inhibitor of CYP2D6, with tramadol ER resulted in a 50-60% increase in tramadol exposure and a 50-60% decrease in M1 exposure. The clinical consequences of these findings are unknown.

To evaluate the effect of tramadol, a CYP2D6 substrate on quinidine, an in vitro drug interaction study in human liver microsomes was conducted. The results from this study indicate that tramadol has no effect on quinidine metabolism [see *Warnings and Precautions (5.1, 5.7), Drug Interactions (7)*].

## *CYP3A4 Inhibitors and Inducers*

Since tramadol is also metabolized by CYP3A4, administration of CYP3A4 inhibitors, such as ketoconazole and erythromycin, or CYP3A4 inducers, such as rifampin and St. John's Wort, with Tramadol Hydrochloride Extended-Release Capsules may affect the metabolism of tramadol leading to altered tramadol exposure [see *Warnings and Precautions (5.1, 5.7), Drug Interactions (7)*].

## *Cimetidine*

Concomitant administration of tramadol immediate-release tablets with cimetidine, a weak CYP3A4 inhibitor, does not result in clinically significant changes in tramadol pharmacokinetics. No alteration of the Tramadol Hydrochloride Extended-Release Capsules dosage regimen with cimetidine is recommended.

## *Carbamazepine*

Carbamazepine, a CYP3A4 inducer, increases tramadol metabolism. Patients taking carbamazepine may have a significantly reduced analgesic effect of tramadol. Concomitant administration of Tramadol Hydrochloride Extended-Release Capsules and carbamazepine is not recommended.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

Carcinogenicity assessment has been conducted in mice, rats and p53(+/-) heterozygous mice. A slight but statistically significant increase in two common murine tumors, pulmonary and hepatic, was observed in an NMRI mouse carcinogenicity study, particularly in aged mice. Mice were dosed orally up to 30 mg/kg in the drinking water (0.5 times the maximum recommended daily human dosage or MRHD) for approximately two years, although the study was not done with the Maximum Tolerated Dose. This finding is not believed to suggest risk in humans.

No evidence of carcinogenicity was noted in a rat 2-year carcinogenicity study testing oral doses of up to 30 mg/kg in the drinking water (1 times the MRHD). In a second rat study, no evidence of carcinogenicity was noted in rats at oral doses up to 75 mg/kg/day for males and 100 mg/kg/day for females (approximately 2 fold the maximum recommended human daily dose MRHD) for two years. However, the excessive decrease in body weight gain observed in the rat study might have reduced their

sensitivity to any potential carcinogenic effect of the drug. No carcinogenic effect of tramadol was observed in p53(+/-)-heterozygous mice at oral doses up to 150 mg/kg/day for 26 weeks.

### Mutagenesis

Tramadol was mutagenic in the presence of metabolic activation in the mouse lymphoma assay. Tramadol was not mutagenic in the in vitro bacterial reverse mutation assay using Salmonella and E. coli (Ames), the mouse lymphoma assay in the absence of metabolic activation, the in vitro chromosomal aberration assay, or the in vivo micronucleus assay in bone marrow.

### Impairment of Fertility

No effects on fertility were observed for tramadol at oral dose levels up to 50 mg/kg in male rats and 75 mg/kg in female rats. These dosages are 1.2 and 1.8 times the maximum recommended human daily dose based on body surface area, respectively.

## **14 CLINICAL STUDIES**


Tramadol Hydrochloride Extended-Release Capsules are bioequivalent under fasting conditions to another extended-release tramadol product [see *Clinical Pharmacology (12.3)*] which demonstrated efficacy in two of four clinical trials of patients with chronic pain. To qualify for inclusion into these studies, patients were required to have moderate to moderately severe pain as defined by a pain intensity score of  $\geq 40$  mm, off previous medications, on a 0 - 100 mm visual analog scale (VAS).

In one 12-week randomized, double-blind, placebo-controlled study, patients with moderate to moderately severe pain due to osteoarthritis of the knee and/or hip were administered doses from 100 mg to 400 mg daily. Treatment with the extended-release tramadol product was initiated at 100 mg once daily for four days then increased by 100 mg per day increments every five days to the randomized fixed dose. Between 51% and 59% of patients in active treatment groups completed the study and 56% of patients in the placebo group completed the study. Discontinuations due to adverse events were more common in the extended-release tramadol product 200 mg, 300 mg and 400 mg treatment groups (20%, 27%, and 30% of discontinuations, respectively) compared to 14% of the patients treated with the extended-release tramadol product 100 mg and 10% of patients treated with placebo.

Pain, as assessed by the WOMAC Pain subscale, was measured at 1, 2, 3, 6, 9, and 12 weeks and changed from baseline assessed. A responder analysis based on the percent change in WOMAC Pain subscale demonstrated a statistically significant improvement in pain for the 100 mg and 200 mg treatment groups compared to placebo (see Figure 2).

### **Figure 2**


**Tramadol ER Tablets Study 023 WOMAC Pain Responder Analysis**  
 Patients Achieving Various Levels of Response Threshold



In one 12-week randomized, double-blind, placebo-controlled flexible-dosing trial of the extended-release tramadol product in patients with osteoarthritis of the knee, patients titrated to an average daily dose of approximately 270 mg/day. Forty-nine percent of patients randomized to the active treatment group completed the study, while 52% of patients randomized to placebo completed the study. Most of the early discontinuations in the active treatment group were due to adverse events, accounting for 27% of the early discontinuations in contrast to 7% of the discontinuations from the placebo group. Thirty-seven percent of the placebo-treated patients discontinued the study due to lack of efficacy compared to 15% of active-treated patients. The active treatment group demonstrated a statistically significant decrease in the mean Visual Analog Scale (VAS) score, and a statistically significant difference in the responder rate, based on the percent change from baseline in the VAS score, measured at 1, 2, 4, 8, and 12 weeks, between patients receiving the extended-release tramadol product and placebo (see Figure 3).

**Figure 3**

**Tramadol ER Tablets Study 015 Arthritis Pain Intensity VAS Responder Analysis**  
 Patients Achieving Various Levels of Response Threshold



Four randomized, placebo-controlled clinical trials of Tramadol Hydrochloride Extended-Release Capsules were conducted, none of which demonstrated efficacy but which differed in design from the preceding clinical studies described. Two trials were 12-week randomized placebo-controlled trials of Tramadol Hydrochloride Extended-Release Capsules 100 mg/day, 200 mg/day, and 300 mg/day versus placebo in patients with moderate to moderately severe osteoarthritis pain of the hip and knee. The other two 12 week trials were similar in design, but only studied Tramadol Hydrochloride Extended-Release Capsules 300 mg/day. In this fixed-dose design, subjects were required to titrate to a fixed dose, even if their pain responded to a lower titration dose.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Tramadol Hydrochloride Extended-Release Capsules are supplied as opaque white hard gelatin capsules, imprinted as follows.

100 mg Capsules: Bottle of 30 capsules:	White capsule imprinted with blue ink "G 252" on cap and "100" between lines on the body  NDC 13811-689-30
200 mg Capsules: Bottle of 30 capsules:	White capsule imprinted with violet ink "G 253" on cap and "200" between lines on the body  NDC 13811-690-30
300 mg Capsules: Bottle of 30 capsules:	White capsule imprinted with red ink "G 254" on cap and "300" between lines on the body  NDC 13811-691-30

Dispense in a tight container. Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [*see USP Controlled Room Temperature*]. Keep out of reach of children.

Store Tramadol Hydrochloride Extended-Release Capsules securely and dispose of properly.

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide)

### Storage and Disposal

Because of the risks associated with accidental ingestion, misuse, and abuse, advise patients to store Tramadol Hydrochloride Extended-Release Capsules securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home. Inform patients that leaving Tramadol Hydrochloride Extended-Release Capsules unsecured can pose a deadly risk to others in the home [*see Warnings and Precautions (5.1, 5.2), Drug Abuse and Dependence (9.2)*].

Advise patients and caregivers that when medicines are no longer needed, they should be disposed of promptly. Inform patients that medicine take-back options are the preferred way to safely dispose of most types of unneeded medicines. If no take back programs or DEA-registered collectors are available, instruct patients to dispose of Tramadol Hydrochloride Extended-Release Capsules by following these four steps:

- Mix Tramadol Hydrochloride Extended-Release Capsules (do not crush) with an unpalatable substance such as dirt, cat litter, or used coffee grounds;
- Place the mixture in a container such as a sealed plastic bag;
- Throw the container in the household trash;
- Remove all personal information on the prescription label of the empty bottle

Inform patients that they can visit [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal) for additional information on disposal of unused medicines.

### Addiction, Abuse, and Misuse

Inform patients that the use of Tramadol Hydrochloride Extended-Release Capsules even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose and death [see *Warnings and Precautions (5.1)*]. Instruct patients not to share Tramadol Hydrochloride Extended-Release Capsules with others and to take steps to protect Tramadol Hydrochloride Extended-Release Capsules from theft or misuse.

### Life-Threatening Respiratory Depression

Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting Tramadol Hydrochloride Extended-Release Capsules or when the dosage is increased, and that it can occur even at recommended dosages.

Educate patients and caregivers on how to recognize respiratory depression and emphasize the importance of calling 911 or getting emergency medical help right away in the event of a known or suspected overdose [see *Warnings and Precautions (5.2), Overdosage (10)*].

### Accidental Ingestion

Inform patients that accidental ingestion, especially by children, may result in respiratory depression or death [see *Warnings and Precautions (5.2)*]. Instruct patients to take steps to store Tramadol Hydrochloride Extended-Release Capsules securely and to dispose of unused Tramadol Hydrochloride Extended-Release Capsules in accordance with the local state guidelines and/or regulations.

### Interactions with Benzodiazepines and Other CNS Depressants

Inform patients and caregivers that potentially fatal additive effects may occur if Tramadol Hydrochloride Extended-Release Capsules are used with benzodiazepines or other CNS depressants, including alcohol (e.g., non-benzodiazepine sedative/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, gabapentinoids [gabapentin or pregabalin], and other opioids), and not to use these concomitantly unless supervised by a healthcare provider [see *Warnings and Precautions (5.3), Drug Interactions (7)*].

### Patient Access to an Opioid Overdose Reversal Agent for the Emergency Treatment of Opioid Overdose

Inform patients and caregivers about opioid overdose reversal agents (e.g., naloxone, nalmefene). Discuss the importance of having access to an opioid overdose reversal agent, especially if the patient has risk factors for overdose (e.g., concomitant use of CNS depressants, a history of opioid use disorder, or prior opioid overdose) or if there are household members (including children) or other close contacts at risk for accidental ingestion or opioid overdose.

Discuss with the patient the options for obtaining an opioid overdose reversal agent (e.g., prescription, over-the-counter, or as part of a community-based program) [see *Dosage and Administration (2.2)*, *Warnings and Precautions (5.2)*].

Educate patients and caregivers on how to recognize the signs and symptoms of an overdose.

Explain to patients and caregivers that effects of opioid overdose reversal agents like naloxone and nalmefene are temporary, and that they must call 911 or get emergency medical help right away in all cases of known or suspected opioid overdose, even if an opioid overdose reversal agent is administered [see *Overdosage (10)*].

Advise patients and caregivers:

- how to treat with the overdose reversal agent in the event of an opioid overdose.
- to tell family and friends about the opioid overdose reversal agent and to keep it in a place where family and friends can access it in an emergency.
- to read the Patient Information (or other educational material) that will come with the opioid overdose reversal agent. Emphasize the importance of doing this before an opioid emergency happens, so the patient and caregiver will know what to do.

### Ultra-Rapid Metabolism of Tramadol and Other Risk Factors for Life-Threatening Respiratory Depression in Children

Advise caregivers that Tramadol Hydrochloride Extended-Release Capsules is contraindicated in all children younger than 12 years of age and in children younger than 18 years of age following tonsillectomy and/or adenoidectomy. Advise caregivers of children ages 12 to 18 years of age receiving Tramadol Hydrochloride Extended-Release Capsules to watch for signs of respiratory depression [see *Warnings and Precautions (5.6)*].

### Hyperalgesia and Allodynia

Inform patients and caregivers not to increase opioid dosage without first consulting a clinician. Advise patients to seek medical attention if they experience symptoms of hyperalgesia, including worsening pain, increased sensitivity to pain, or new pain [see *Warnings and Precautions (5.8)*, *Adverse Reactions (6.2)*].

### Serotonin Syndrome

Inform patients that tramadol could cause a rare but potentially life-threatening condition called serotonin syndrome, particularly during concomitant use with serotonergic drugs. Warn patients of the symptoms of serotonin syndrome and to seek medical attention right away if symptoms develop. Instruct patients to inform their healthcare provider if they are taking, or plan to take serotonergic medications [see *Warnings and Precautions (5.9)*, *Drug Interactions (7)*].

### Seizures

Inform patients that Tramadol Hydrochloride Extended-Release Capsules may cause seizures with concomitant use of serotonergic agents (including SSRIs, SNRIs, and triptans) or drugs that significantly reduce the metabolic clearance of tramadol [see *Warnings and Precautions (5.10)*].

#### MAOI Interaction

Inform patients not to take Tramadol Hydrochloride Extended-Release Capsules while using any drugs that inhibit monoamine oxidase. Patients should not start MAOIs while taking Tramadol Hydrochloride Extended-Release Capsules [see *Drug Interactions (7)*].

#### Important Administration Instructions

Instruct patients how to properly take Tramadol Hydrochloride Extended-Release Capsules, including the following:

- Tramadol Hydrochloride Extended-Release Capsules are designed to work properly only if swallowed intact. Taking cut, broken, chewed, crushed, or dissolved Tramadol Hydrochloride Extended-Release Capsules can result in a fatal overdose [see *Dosage and Administration (2.1)*].
- Advise patients not to exceed the single-dose and 24-hour dose limit and the time interval between doses, since exceeding these recommendations can result in respiratory depression, seizures, hepatic toxicity, and death.
- Tramadol Hydrochloride Extended-Release Capsules should not be taken with alcohol-containing beverages.

#### Important Discontinuation Instructions

In order to avoid developing withdrawal symptoms, instruct patients not to discontinue Tramadol Hydrochloride Extended-Release Capsules without first discussing a tapering plan with the prescriber [see *Dosage and Administration (2.5)*].

#### Driving or Operating Heavy Machinery

Inform patients that Tramadol Hydrochloride Extended-Release Capsules may impair the ability to perform potentially hazardous activities such as driving a car or operating heavy machinery. Advise patients not to perform such tasks until they know how they will react to the medication [see *Warnings and Precautions (5.19)*].

#### Constipation

Advise patients of the potential for severe constipation, including management instructions and when to seek medical attention [see *Adverse Reactions (6)*, *Clinical Pharmacology (12.1)*].

#### Adrenal Insufficiency

Inform patients that Tramadol Hydrochloride Extended-Release Capsules could cause adrenal insufficiency, a potentially life-threatening condition. Adrenal insufficiency may present with non-specific symptoms and signs such as nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. Advise patients to seek medical attention if they experience a constellation of these symptoms [see *Warnings and Precautions (5.13)*].

#### Hypotension

Inform patients that Tramadol Hydrochloride Extended-Release Capsules may cause

orthostatic hypotension and syncope. Instruct patients how to recognize symptoms of low blood pressure and how to reduce the risk of serious consequences should hypotension occur (e.g., sit or lie down, carefully rise from a sitting or lying position) *[see Warnings and Precautions (5.14)]*.

### Anaphylaxis

Inform patients that anaphylaxis has been reported with ingredients contained in Tramadol Hydrochloride Extended-Release Capsules. Advise patients how to recognize such a reaction and when to seek medical attention *[see Contraindications (4), Adverse Reactions (6)]*.

### Pregnancy

#### *Neonatal Opioid Withdrawal Syndrome*

Inform female patients of reproductive potential that use of Tramadol Hydrochloride Extended-Release Capsules for an extended period of time during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated *[see Warnings and Precautions (5.5), Use in Specific Populations (8.1)]*.

#### *Embryo-Fetal Toxicity*

Inform female patients of reproductive potential that Tramadol Hydrochloride Extended-Release Capsules can cause fetal harm and to inform their healthcare provider of a known or suspected pregnancy *[see Use in Specific Populations (8.1)]*.

### Lactation

Advise women that breastfeeding is not recommended during treatment with Tramadol Hydrochloride Extended-Release Capsules *[see Use in Specific Populations (8.2)]*.

### Infertility

Inform patients that use of opioids for an extended period of time may cause reduced fertility. It is not known whether these effects on fertility are reversible *[see Adverse Reactions (6.2), Use in Specific Populations (8.3)]*.

Manufactured by: Galephar P.R., Inc.  
Juncos, Puerto Rico 00777

Manufactured for: Trigen Laboratories, LLC  
Alpharetta, GA 30005, USA

Rev. 12/2025

## **Medication Guide**

### **Tramadol Hydrochloride Extended-Release Capsules, CIV**

#### **Tramadol Hydrochloride Extended-Release Capsules are:**

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage severe and persistent pain that requires an extended treatment period with a daily opioid pain medicine when other pain medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.

- Not to be taken on an "as needed" basis.

### **Important information about Tramadol Hydrochloride Extended-Release Capsules:**

- **Get emergency help right away or call 911 right away if you take too much Tramadol Hydrochloride Extended-Release Capsules (overdose).** When you first start taking Tramadol Hydrochloride Extended-Release Capsules, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur. Ask your healthcare provider about medicines like naloxone or nalmefene that can be used in an emergency to reverse an opioid overdose.
- Taking Tramadol Hydrochloride Extended-Release Capsules with other opioid medicines, benzodiazepines, gabapentinoids (gabapentin or pregabalin), alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Tramadol Hydrochloride Extended-Release Capsules. They could die from taking them. Selling or giving away Tramadol Hydrochloride Extended-Release Capsules is against the law.
- Store Tramadol Hydrochloride Extended-Release Capsules securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

### **Important Information Guiding Use in Pediatric Patients:**

- Do not give Tramadol Hydrochloride Extended-Release Capsules to a child younger than 12 years of age.
- Do not give Tramadol Hydrochloride Extended-Release Capsules to a child younger than 18 years of age after surgery to remove the tonsils and/or adenoids.
- Avoid giving Tramadol Hydrochloride Extended-Release Capsules to children between 12 to 18 years of age who have risk factors for breathing problems such as obstructive sleep apnea, obesity, or underlying lung problems.

### **Do not take Tramadol Hydrochloride Extended-Release Capsules if you have:**

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

### **Before taking Tramadol Hydrochloride Extended-Release Capsules, tell your healthcare provider if you have a history of:**

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, opioid overdose, or mental health problems.

### **Tell your healthcare provider if you are:**

- **noticing your pain getting worse.** If your pain gets worse after you take Tramadol Hydrochloride Extended-Release Capsules, do not take more of Tramadol Hydrochloride Extended-Release Capsules without first talking to your healthcare provider. Talk to your healthcare provider if the pain that you have increases, if you feel more sensitive to pain, or if you have new pain after taking Tramadol Hydrochloride Extended-Release Capsules.

- **pregnant or planning to become pregnant.** Use of Tramadol Hydrochloride Extended-Release Capsules for an extended period of time during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** Not recommended during treatment with Tramadol Hydrochloride Extended-Release Capsules; it may harm your baby.
- living in a household where there are small children or someone who has abused street or prescription drugs.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking Tramadol Hydrochloride Extended-Release Capsules with certain other medicines can cause serious side effects that could lead to death.

#### **When taking Tramadol Hydrochloride Extended-Release Capsules:**

- Do not change your dose. Take Tramadol Hydrochloride Extended-Release Capsules exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- Take your prescribed dose once a day at the same time every day. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Swallow Tramadol Hydrochloride Extended-Release Capsules whole. Do not split, break, chew, crush, dissolve, snort, or inject Tramadol Hydrochloride Extended-Release Capsules because this may cause you to overdose and die.
- **Call your healthcare provider if the dose you are taking does not control your pain.**
- **Do not stop taking Tramadol Hydrochloride Extended-Release Capsules without talking to your healthcare provider.**
- Dispose of expired, unwanted, or unused Tramadol Hydrochloride Extended-Release Capsules by taking your drug to an authorized DEA-registered collector or drug take-back program. If one is not available, you can dispose of Tramadol Hydrochloride Extended-Release Capsules by mixing the product with dirt, cat litter, or coffee grounds; placing the mixture in a sealed plastic bag and throwing the bag in your trash. Visit [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal) for additional information on disposal of unused medicines.

#### **While taking Tramadol Hydrochloride Extended-Release Capsules DO NOT:**

- Drive or operate heavy machinery, until you know how Tramadol Hydrochloride Extended-Release Capsules affect you. Tramadol Hydrochloride Extended-Release Capsules can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Tramadol Hydrochloride Extended-Release Capsules may cause you to overdose and die.

#### **The possible side effects of Tramadol Hydrochloride Extended-Release Capsules:**

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain, seizure. Call your healthcare provider if you have any of these symptoms and they are severe.

#### **Get emergency medical help or call 911 right away if you have:**

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your

face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of Tramadol Hydrochloride Extended-Release Capsules. Call your healthcare provider for medical advice about side effects. You may report side effects to Trigen Laboratories, LLC at 1-800-541-4802 or FDA at 1-800-FDA-1088. **For more information go to [dailymed.nlm.nih.gov](https://dailymed.nlm.nih.gov)**

Manufactured for: Trigen Laboratories, LLC, Alpharetta, GA 30005, USA, call 1-800-541-4802.

**This Medication Guide has been approved by the U.S. Food and Drug Administration.**

Revised:  
December  
2025

### Principal Display Panel - 100 mg Capsule Bottle Label

NDC 13811-689-30

Tramadol Hydrochloride CIV  
Extended-Release Capsules

100 mg per capsule

Once Daily  
Rx ONLY

TRIGEN  
LABORATORIES

30 Capsules

The image shows a detailed view of a bottle label for Tramadol Hydrochloride CIV Extended-Release Capsules. The label is rectangular with a dashed border. On the left side, it features the NDC number 13811-689-30, the product name in bold, and a blue box indicating '100 mg per capsule'. Below this, it says 'Once Daily Rx ONLY' and the Trigen Laboratories logo. On the right side, there is a 'Usual Adult Dose' section with three bullet points: 'Refer to package insert.', 'Swallow capsule whole.', and 'Do not crush or chew.' Below that is a 'Dispense the accompanying Medication Guide to each patient.' warning: 'KEEP OUT OF REACH OF CHILDREN.' and storage instructions: 'Each extended-release capsule contains: 100 mg of tramadol hydrochloride, USP. Store at 25°C; excursions permitted to 15°C - 30°C (59°F - 86°F). [See USP Controlled Room Temperature].' The distributor information is 'Distributed by: Trigen Laboratories, LLC, Alpharetta, GA 30005, GK-372-1 Rev. 01/2022'. A barcode is located on the right side of the label, with the number 'N 3 13811 68930 3' printed below it. A large pink area on the far right of the label is labeled 'NO VARNISH AREA'.

### Principal Display Panel - 200 mg Capsule Bottle Label

NDC 13811-690-30

Tramadol Hydrochloride CIV  
Extended-Release Capsules

200 mg per capsule

Once Daily

R<sub>x</sub> ONLY

TRIGEN  
LABORATORIES

30 Capsules

**NDC 13811-690-30**

**Tramadol Hydrochloride <sup>Ⓒ</sup>IV**  
**Extended-Release Capsules**

**200 mg per capsule**

Once Daily  
R<sub>x</sub> ONLY

**TRIGEN**  
LABORATORIES

**30 Capsules**

**Usual Adult Dose:**

- Refer to package insert.
- Swallow capsule whole.
- Do not crush or chew.

Dispense the accompanying Medication Guide to each patient.

**WARNING: KEEP OUT OF REACH OF CHILDREN.**

Each extended-release capsule contains: 200 mg of tramadol hydrochloride, USP

Store at 25°C; excursions permitted to 15°C - 30°C (59°F - 86°F).  
[See USP Controlled Room Temperature].

Distributed by: **TRIGEN LABORATORIES, LLC**  
Alpharetta, GA 30005  
[www.trigenlab.com](http://www.trigenlab.com)

Rev. 01/2022 GK-373-1

3 13811 69030 9

**NO VARNISH AREA**

### Principal Display Panel - 300 mg Capsule Bottle Label

NDC 13811-691-30

Tramadol Hydrochloride CIV  
Extended-Release Capsules

300 mg per capsule

Once Daily

R<sub>x</sub> ONLY

TRIGEN  
LABORATORIES

30 Capsules

NDC 13811-691-30

# Tramadol Hydrochloride <sup>IV</sup> Extended-Release Capsules

300 mg per capsule

Once Daily  
Rx ONLY



30 Capsules

**Usual Adult Dose:**

- Refer to package insert.
- Swallow capsule whole.
- Do not crush or chew.

Dispense the accompanying Medication Guide to each patient.

WARNING: KEEP OUT OF REACH OF CHILDREN.

Each extended-release capsule contains: 300 mg of tramadol hydrochloride, USP

Store at 25°C; excursions permitted to 15°C - 30°C (59°F - 86°F). [See USP Controlled Room Temperature].

Distributed by: TRIGEN LABORATORIES, LLC  
Alpharetta, GA 30005  
www.trigenlab.com

Rev. 01/2022 GK374-1



NO VARNISH  
AREA

## TRAMADOL HYDROCHLORIDE EXTENDED-RELEASE

tramadol hydrochloride capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:13811-689
Route of Administration	ORAL	DEA Schedule	CIV

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRAMADOL HYDROCHLORIDE (UNII: 9N7R477WCK) (TRAMADOL - UNII:39J1LGJ30J)	TRAMADOL HYDROCHLORIDE	100 mg

### Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SHELLAC (UNII: 46N107B71O)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SUCROSE STEARATE (UNII: 274KW0050M)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
TALC (UNII: 7SEV7J4R1U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
ETHYL ACRYLATE AND METHYL METHACRYLATE COPOLYMER (2:1; 750000 MW) (UNII: P2OM2Q86BI)	

## Product Characteristics

<b>Color</b>	WHITE (white)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	G;252;100
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13811-689-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/22/2015	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA022370	06/22/2015	

## TRAMADOL HYDROCHLORIDE EXTENDED-RELEASE

tramadol hydrochloride capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:13811-690
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CIV

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>TRAMADOL HYDROCHLORIDE</b> (UNII: 9N7R477WCK) (TRAMADOL - UNII:39J1LGJ30J)	TRAMADOL HYDROCHLORIDE	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>D&amp;C RED NO. 7</b> (UNII: ECW0LZ41X8)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SUCROSE STEARATE</b> (UNII: 274KW0050M)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>ETHYL ACRYLATE AND METHYL METHACRYLATE COPOLYMER (2:1; 750000 MW)</b> (UNII: P2OM2Q86BI)	

### Product Characteristics

<b>Color</b>	WHITE (white)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	22mm
<b>Flavor</b>		<b>Imprint Code</b>	G;253;200
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13811-690-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/22/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA022370	06/22/2015	

## TRAMADOL HYDROCHLORIDE EXTENDED-RELEASE

tramadol hydrochloride capsule

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:13811-691
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CIV

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>TRAMADOL HYDROCHLORIDE</b> (UNII: 9N7R477WCK) (TRAMADOL - UNII:39J1LGJ30J)	TRAMADOL HYDROCHLORIDE	300 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

<b>SHELLAC</b> (UNII: 46N107B71O)
<b>D&amp;C RED NO. 7</b> (UNII: ECW0LZ41X8)
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)
<b>SUCROSE STEARATE</b> (UNII: 274KW0O50M)
<b>POVIDONE K30</b> (UNII: U725QWY32X)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>ETHYL ACRYLATE AND METHYL METHACRYLATE COPOLYMER (2:1; 750000 MW)</b> (UNII: P2OM2Q86BI)

### Product Characteristics

<b>Color</b>	WHITE (white)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (CAPSULE)	<b>Size</b>	23mm
<b>Flavor</b>		<b>Imprint Code</b>	G;254;300
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13811-691-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/22/2015	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA AUTHORIZED GENERIC	NDA022370	06/22/2015	

**Labeler** - Trigen Laboratories, LLC (830479668)