

IBUPROFEN- ibuprofen tablet, film coated
Meijer Distribution Inc

Meijer 44-291

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - headache
 - toothache
 - backache
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- rash
- blisters
- shock
- skin reddening
- asthma (wheezing)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have problems or serious side effects from taking pain relievers or fever reducers
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have stomach pain that does not get better
 - have bloody or black stools
- you have symptoms of heart problems or stroke:
 - chest pain
 - leg swelling
 - trouble breathing
 - slurred speech
 - weakness in one part or side of body
- fever gets worse or lasts more than 3 days
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

meijer®

NDC 41250-991-03

COMPARE TO ADVIL® TABLETS

ACTIVE INGREDIENT†

TAMPER EVIDENT: Use Only if This Blister is Intact

Ibuprofen

Tablets USP, 200 mg

PAIN RELIEVER | FEVER REDUCER
(NSAID)

**10
TABLETS**

Actual Size

TAMPER EVIDENT: Use Only if This Blister is Intact

†This product is not manufactured or distributed by Haleon US Holdings LLC, owner of the registered trademark Advil® Tablets.

DIST. BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

**OUR QUALITY
GUARANTEE**

WWW.MEIJER.COM/SATISFACTION

The Meijer
Family

50844

REV1221E29103

meijer

NDC 41250-991-03

COMPARE TO ADVIL® TABLETS ACTIVE INGREDIENT†

Ibuprofen
Tablets USP, 200 mg

PAIN RELIEVER | FEVER REDUCER (NSAID)

10 TABLETS

Actual Size

TAMPER EVIDENT: Use Only if This Blister is Intact

**HEAT SEAL COATING ONLY
NO COPY AREA**

Drug Facts (continued)

When using this product • take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have stomach pain that does not get better
 - have bloody or black stools
- you have symptoms of heart problems or stroke:
 - chest pain
 - leg swelling
 - trouble breathing
 - slurred speech
 - weakness in one part or side of body
 - fever gets worse or lasts more than 3 days
 - pain gets worse or lasts more than 10 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • do not take more than directed

- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information • store between 20°-25°C (68°-77°F)

- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients • carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

†This product is not manufactured or distributed by Haleon US Holdings LLC, owner of the registered trademark Advil® Tablets.

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com 50844 REV1221E29103

Meijer 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:41250-991

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-991-03	1 in 1 PACKAGE	05/24/1988	
1		10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - Meijer Distribution Inc (006959555)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(41250-991)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41250-991)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41250-991)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(41250-991)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41250-991)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(41250-991)