

**NICOTINE POLACRILEX COATED FRUIT- nicotine polacrilex gum, chewing  
Publix Supermarkets, Inc.**

-----  
**Nicotine Polacrilex Gum 2 mg and 4 mg Coated Fruit - Publix**

***Drug Facts***

***Active ingredient (in each chewing piece) - 2 mg***

Nicotine Polacrilex 2 mg (nicotine)

***Active ingredient (in each chewing piece) - 4 mg***

Nicotine Polacrilex 4 mg (nicotine)

***Purpose***

Stop smoking aid

***Use***

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

***Warnings***

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may

need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.**

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center(1-800-222-1222) right away.

**Directions - 2 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**Directions - 4 mg**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

#### ***Other information - 2 mg***

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

#### ***Other information - 4 mg***

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20 - 25°C (68 - 77°F)
- protect from light

#### ***Inactive ingredients - 2 mg***

acacia, acesulfame potassium, carnauba wax, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide and xylitol.

#### ***Inactive ingredients - 4 mg***

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 lake, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide and xylitol.

## **Principal display panel 2 mg**

\*Compare to the active ingredient in Nicorette® Gum

SUGAR FREE

nicotine gum

NICOTINE POLACRILEX GUM USP, 2mg (nicotine)

STOP SMOKING AID

COATED FRUIT FLAVOR

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.

If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicotine Polacrilex Gum, 4 mg

2 mg

### **TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
- 2 **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.
  - Not for sale to those under 18 years of age
  - Proof of age required.
  - Not for sale in vending machines or from any source where proof of age cannot be verified.

Blister packages for your protect. Do not use if individual seals are open or torn.


\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

DISTRIBUTED BY PUBLIX SUPER MARKETS, INC.,

3300 PUBLIX CORPORATE PARKWAY,  
LAKELAND, FL 33811, 1-888-267-3037

**Package label 2 mg**





**PUBLIX Sugar Free Nicotine Coated Fruit Flavor**

**Principal display panel 4 mg**

\*Compare to the active ingredient in Nicorette® Gum

SUGAR FREE

nicotinegum

NICOTINE POLACRILEX GUM USP, 4 mg (NICOTINE)

STOP SMOKING AID

COATED FRUIT FLAVOR

FOR THOSE SMOKE THEIR FIRST CIGARETTE WITHIN 302 MINUTES OG WAKING UP.

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up use Nicotine Polacrilex Gum USP, 2 mg

4 mg

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivates to quit.
2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use with a Support Program** as directed in the enclosed User's Guide.
  - Not for sale to those under 18 years of age.
  - Proof of age required.
  - Not for sale in vending machines or from any source where proof of age cannot be verified

Blister packaged for your protection. Do not use if individual seals are open or torn.

\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., Distributor of Nicorette® Gum

DISTRIBUTED BY PUBLIX SUPERMARKETS, INC.,

3300 PUBLIX CORPORATE PARKWAY,  
LAKELAND, FL 33811, 1-888-267-3037

**Package label 4 mg**







Drug Facts (continued)	Warnings	Use	Active Ingredient (in each chewing piece)	Purpose						
<p>■ If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:</p> <table border="1"> <tr> <td>Weeks 1 to 6</td> <td>1 piece every 1 to 2 hours</td> </tr> <tr> <td>Weeks 7 to 9</td> <td>1 piece every 2 to 4 hours</td> </tr> <tr> <td>Weeks 10 to 12</td> <td>1 piece every 4 to 8 hours</td> </tr> </table> <p>■ Chew the gum slowly until it begins to melt. Then park it between your cheek and teeth. Do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece.</p> <p>■ Do not use more than 24 pieces a day.</p> <p>■ It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.</p>	Weeks 1 to 6	1 piece every 1 to 2 hours	Weeks 7 to 9	1 piece every 2 to 4 hours	Weeks 10 to 12	1 piece every 4 to 8 hours	<p>■ Ask a doctor before use if you have a heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.</p> <p>■ Ask a doctor or pharmacist before use if you are taking a non-contraceptive drug containing estrogen, your prescription drug may need to be adjusted.</p> <p>■ Stop use and ask a doctor if you need to be adjusted.</p>	<p>■ Nicotine gum reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.</p> <p>■ If you are pregnant or breast-feeding, only use the medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement products. The medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.</p>	<p><b>nicotinegum</b></p> <p>USP, 4 mg (nicotine)</p>	<p>■ To help you quit smoking, this medicine is used as a short-term treatment. It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.</p>
Weeks 1 to 6	1 piece every 1 to 2 hours									
Weeks 7 to 9	1 piece every 2 to 4 hours									
Weeks 10 to 12	1 piece every 4 to 8 hours									
<p><b>Other Information</b></p> <p>■ Each piece contains calcium 115 mg, sodium 8 mg</p> <p>■ Store at 20°-25°C (68°-77°F)</p> <p>■ Protect from light</p>	<p>■ Blood pressure.</p> <p>■ High blood pressure not controlled with medication. Nicotine can increase blood pressure.</p> <p>■ Stomach ulcer or diabetes</p> <p>■ History of seizures</p>	<p>■ Nicotine gum is not fully known.</p>		<p>■ Ask a doctor or pharmacist before use if you are taking a non-contraceptive drug containing estrogen, your prescription drug may need to be adjusted.</p> <p>■ Stop use and ask a doctor if you need to be adjusted.</p>						



### PUBLIX Nicotine Gum Fruit Flavor

## NICOTINE POLACRILEX COATED FRUIT

nicotine polacrilex gum, chewing

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:56062-503
<b>Route of Administration</b>	BUCCAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM OXIDE (UNII: 3A3U0G171G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XYLITOL (UNII: VCQ006KQ1E)	

### Product Characteristics

<b>Color</b>	brown (light brown)	<b>Score</b>	no score
--------------	---------------------	--------------	----------

<b>Shape</b>	SQUARE	<b>Size</b>	14mm
<b>Flavor</b>	FRUIT (coated fruit)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-503-44	10 in 1 CARTON	01/01/2014	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079044	01/01/2014	

## NICOTINE POLACRILEX COATED FRUIT

nicotine polacrilex gum, chewing

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:56062-504
<b>Route of Administration</b>	BUCCAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NICOTINE</b> (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>ACESULFAME POTASSIUM</b> (UNII: 230V73Q5G9)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>MAGNESIUM OXIDE</b> (UNII: 3A3U0GI71G)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>SODIUM CARBONATE</b> (UNII: 45P3261C7T)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>XYLITOL</b> (UNII: VCQ006KQ1E)	

### Product Characteristics

<b>Color</b>	brown (light tan)	<b>Score</b>	no score
<b>Shape</b>	SQUARE	<b>Size</b>	14mm
<b>Flavor</b>	FRUIT (coated fruit)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-504-44	10 in 1 CARTON	01/01/2014	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079038	01/01/2014	

**Labeler** - Publix Supermarkets, Inc. (006922009)

Revised: 8/2022

Publix Supermarkets, Inc.