## DEXBROMPHENIRAMINE MALEATE AND PHENYLEPHRINE HYDROCHLORIDEdexbrompheniramine maleate and phenylephrine hydrochloride tablet Westminster Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## Dexbrompheniramine Maleate and Phenylephrine HCl

#### **DRUG FACTS**

Active ingredients (in each tablet)	Purpose
Dexbrompheniramine Maleate 2 mg	Antihistamine
Phenylephrine HCL 10 mg	Nasal
Pilenylephinie HCL 10 liig	Decongestant

#### Uses

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

#### Warnings

#### Do not exceed recommended dosage.

#### Do not use this product

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to enlargment of the prostate gland
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

## When using this product

- excitability may occur, especially in children
- may cause drowsiness
- avoid alcoholic drinks

- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever

**If pregnant or breast-feeding,** ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

## Do not exceed recommended dosage.

	1 tablet every 4 to 6 hours, not to exceed 6 tablets in 24 hours or as directed by doctor.
Children 6 to under 12 years of age:	1/2 tablet every 4 to 6 hours, not to exceed 3 tablets in 24 hours or as directed by doctor.
Children under 6 years of age:	Consult a doctor.

#### Other Information

Store at 15°-30°C (59°-86°F). Supplied in a tight, light resistant container with a child-resistant cap. Dexbrompheniramine Maleate and Phenylephrine HCl tablets are purple, oval shaped, scored tablets, debossed "WP" bisect "197" on one side and plain on the other.

#### **Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, D&C Red #27, dibasic calcium phosphate dihydrate, FD&C Blue #1, magnesium stearate, and microcrystalline cellulose.

### **Questions? Comments?**

Call 1-844-221-7294

#### PRINCIPAL DISPLAY PANEL - 60 Tablet Bottle Label

NDC 69367-197-60

Dexbrompheniramine Maleate and Phenylephrine HCl


**Antihis tamine • Nas al Deconges tant** 

Each tablet contains: Dexbrompheniramine Maleate 2 mg

## Phenylephrine HCI 10 mg

60 Tablets

Westminster Pharmaceuticals



#### DRUG FACTS (continued) DRUG FACTS (continued) monoamine oxidase inhibitor (MAOI) (certain Other Information drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 Store at 15°-30°C (59°-86°F). Supplied in a tight, light resistant container with a child-resistant cap. weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an Dexbrompheniramine Maleate and Phenylephrine HCl tablets are purple, oval shaped, scored tablets, debossed "WP" bisect "197" on one side MAOI, ask a doctor or pharmacist before taking this product. and plain on the other. Ask a doctor before use if you have a breathing problem such as emphysema or Inactive ingredients chronic bronchitis ■ glaucoma ■ trouble urinating due to enlargment of the prostate gland ■ heart disease ■ high blood pressure colloidal silicon dioxide, croscarmellose sodium, D&C Red #27, dibasic calcium phosphate dihydrate, FD&C Blue #1, magnesium stearate, and microcrystalline cellulose. ■ thyroid disease ■ diabetes Ask a doctor or pharmacist before use it you are taking sedatives or tranquilizers. Questions? Comments? Call 1-844-221-7294 When using this product excitability may occur, especially in children may cause drowsiness avoid alcoholic drinks Manufactured for: Westminster Pharmaceuticals, LLC. ■ alcohol, sedatives, and tranquilizers may Olive Branch, MS 38654 Rev. 10/18 increase the drowsiness effect use caution when driving a motor vehicle or operating machinery DRUG FACTS Stop use and ask a doctor if Active ingredients Purpose ■ nervousness, dizziness, or sleeplessness (in each tablet) symptoms do not improve within 7 days or are accompanied by fever Dexbrompheniramine Maleate 2 mg. Antihistamine If pregnant or breast-feeding, ask a health professional before use. Phenylephrine HCL 10 mg... .. Nasal Decongestant Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or Directions: Do not exceed recommended dosage. Adults and children 12 1 tablet every 4 to 6 hours, not to exceed 6 tablets in 24 hours throat itchy, watery eyes nasal congestion reduces swelling of nasal passages years of age or as directed by doctor. and over 1/2 tablet every 4 to 6 hours, not to exceed 3 tablets in 24 hours Children 6 to under 12 Warnings Do not exceed recommended dosage years of age: or as directed by doctor. Do not use this product ■ if you are now taking a prescription Children under 6 years | Consult a doctor. of age: Lift Here

# DEXBROMPHENIRAMINE MALEATE AND PHENYLEPHRINE HYDROCHLORIDE

dexbrompheniramine maleate and phenylephrine hydrochloride tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69367-197
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
<b>DEXBROMPHENIRAMINE MALEATE</b> (UNII: BPA9 UT29 BS) (DEXBROMPHENIRAMINE - UNII:75T6 4B71RP)	DEXBROMPHENIRAMINE MALEATE	2 mg		
<b>PHENYLEPHRINE HYDRO CHLO RIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg		

Inactive Ingredients		
Ingredient Name	Strength	
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)		
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS 185U6K)		
DIBASIC CALCIUM PHO SPHATE DIHYDRATE (UNII: O7TSZ97GEP)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)		
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)		

Product Characteristics			
Color	PURPLE	Score	2 pieces
Shape	OVAL	Size	11mm
Flavor		Imprint Code	WP;197
Contains			

l	Packaging				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1 NDC:69367-197-	60 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/25/2018		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/25/2018	

## Labeler - Westminster Pharmaceuticals (079516651)

Revised: 1/2020 Westminster Pharmaceuticals