

RANITIDINE- ranitidine tablet
Shopko Stores Operating Co., LLC.

Drug Facts

Active ingredient (in each tablet)

Ranitidine 150 mg (as ranitidine hydrochloride 168 mg)

Purpose

Acid reducer

Uses

- **relieves** heartburn associated with acid indigestion and sour stomach
- **prevents** heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have:

- frequent **chest pain**
- unexplained weight loss
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent wheezing, particularly with heartburn
- nausea or vomiting

Stop use and ask a doctor if

- your heartburn continues or worsens
- stomach pain continues
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and older:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily -do not take more than 2 tablets in 24 hours
- children under 12 years of age: ask a doctor

Other information

- do not use if printed foil under cap is torn or open
- store at controlled room temperature 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- this product is sugar free
- *This product is not manufactured or distributed by Glaxosmithkline Consumer Healthcare, owner of the registered trademark Zantac® 150

Inactive Ingredients

colloidal silicon dioxide, croscarmellose sodium, diethyl phthalate, FD&C yellow # 6, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose and titanium dioxide.

Principal display panel

SHOPKO®

Maximum Strength HEARTBURN

Ranitidine Tablets, USP 150 mg

Acid Reducer

One tablet relieves: heartburn, acid indigestion & sour stomach

*Compare to Zantac 150® active ingredient

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Packaged by:

PL Developments

200 Hicks street

Westbury, NY 11590, USA

Questions or comments?

call toll free 1-877-753-3935

Product of INDIA

Product Label

NO INK
NO VARNISH

NO INK
NO VARNISH

Drug Facts
Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride 168 mg).....Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or have bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers. ■ if you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have: ■ frequent chest pain ■ unexplained weight loss ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with **lightheadedness, sweating, or dizziness** ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck, or shoulders, or lightheadedness. ■ frequent wheezing, particularly with heartburn ■ nausea or vomiting

Stop use and ask a doctor if ■ your heartburn continues or worsens ■ stomach pain continues ■ you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health care professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ adults and children 12 years of age and older:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 ■ can be used up to twice daily - do not take more than 2 tablets in 24 hours
 ■ children under 12 years of age: ask a doctor

Other information ■ do not use if printed foil under bottle cap is torn or open ■ store at controlled room temperature 20°-25°C (68°-77°F) ■ avoid excessive heat and humidity ■ this product is sugar free. ■ "This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, owner of the registered trademark Zantac® 150

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, diethyl phthalate, FD&C Yellow # 6, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, and titanium dioxide.

SHOPKO®

MAXIMUM STRENGTH HEARTBURN

ranitidine tablets, usp 150 mg

Acid Reducer

One Tablet Relieves:
Heartburn, Acid Indigestion
& Sour Stomach

50 Tablets (50 doses)

SHOPKO®

*COMPARE TO
ZANTAC®150
ACTIVE INGREDIENT

MAXIMUM STRENGTH HEARTBURN

ranitidine tablets, usp
150 mg

Acid Reducer

One Tablet Relieves:
Heartburn, Acid Indigestion
& Sour Stomach

50 Tablets (50 doses)

SHOPKO®

MAXIMUM STRENGTH HEARTBURN

ranitidine tablets, usp 150 mg

Acid Reducer
One Tablet Relieves:
Heartburn, Acid Indigestion
& Sour Stomach

50 Tablets (50 doses)

MAXIMUM STRENGTH HEARTBURN

ranitidine tablets, usp 150 mg

SHOPKO®

PLD-A
F127SHK50
Lot No.:
Exp. Date:

READ THE DIRECTIONS, CONSUMER INFORMATION LEAFLET AND WARNINGS BEFORE USE.
KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

Packaged by:
PL Developments
200 Hicks Street
Westbury, NY 11590, USA
Questions or comments?
Call toll free 1-877-753-3935



PRODUCT OF INDIA

SATISFACTION GUARANTEED
BY REFUND OR EXCHANGE



Ranitidine 150 mg

RANITIDINE

ranitidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:370 12-127
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIETHYL PHTHALATE (UNII: UF064M00AF)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	HEXAGON (6 SIDED)	Size	10mm
Flavor		Imprint Code	W;741
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:370 12-127-50	1 in 1 CARTON		
1		50 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA078653	07/11/2010	
------	------------	------------	--

Labeler - Shopko Stores Operating Co., LLC. (023252638)

Revised: 8/2012

Shopko Stores Operating Co., LLC.