

ACETAMINOPHEN- acetaminophen tablet
TIME CAP LABS INC

340R Timely 49483-340 Acetaminophen 325 mg Tablets

DRUG FACTS

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:

headache

muscular aches

backache

minor pain of arthritis

the common cold

toothache

premenstrual and menstrual cramps

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

adults and children 12 years and over

- take 2 tablets every 4 to 6 hours while symptoms last
- do not take more than 10 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children 6 years to under 12 years

- take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years

- ask a doctor

OTHER INFORMATION

Other information

- SODIUM FREE
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain this ingredient

NDC 49483-340-10
*Compare to the active ingredient in
Tylenol® Regular Strength

timely™

Regular Strength

Pain Reliever
Acetaminophen 325 mg

Pain Reliever / Fever Reducer

Contains no aspirin

THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN

1000 TABLETS

Actual Size



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient
(in each tablet)

Purpose

Acetaminophen 325 mg...Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - the common cold
 - toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

- Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if:
 - adult takes more than 4,000 mg of acetaminophen in 24 hours
 - child takes more than 5 doses in 24 hours, which is the maximum daily amount
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rashIf a skin reaction occurs, stop.

Drug Facts (continued)

use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

not notice any signs or symptoms.

Directions

See overdose warning

- Do not take more than directed.**
- adults and children 12 years and over
 - take 2 tablets every 4 to 6 hours while symptoms last
 - do not take more than 10 tablets in 24 hours, unless directed by a doctor
 - do not use for more than 10 days unless directed by a doctor

children 6 years to under 12 years

- take 1 tablet every 4 to 6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years

- ask a doctor

Other information


- SODIUM FREE**
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

Questions or comments? Call 1-877-290-4008
Manufactured by: Time-Cap Labs, Inc.,
7 Michael Avenue, Farmingdale, NY 11735

LOT #:

EXP. DATE:



5

3

Adhesive Area

Questions or comments?
Call 1-877-290-4008

STOP PEELING

Other information ■ SODIUM FREE ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

Directions ■ Do not take more than directed.

See overdose warning

adults and children 12 years and over

take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor

children 6 years to under 12 years

take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not use for more than 5 days unless directed by a doctor

children under 6 years

ask a doctor

Other information ■ SODIUM FREE ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

Directions ■ Do not take more than directed.

See overdose warning

adults and children 12 years and over

take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor

children 6 years to under 12 years

take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not use for more than 5 days unless directed by a doctor

children under 6 years

ask a doctor

Other information ■ SODIUM FREE ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

NDC 49483-340-01
*Compare to the active ingredient in
Tylenol® Regular Strength

timely™

Regular Strength

Pain Reliever
Acetaminophen 325 mg

Pain Reliever / Fever Reducer

Contains No Aspirin

100 TABLETS

Actual Size



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Acetaminophen 325 mg...Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - the common cold
 - toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether

Drug Facts (continued under label)

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc.,
McNeil Consumer Healthcare Division, owner of the registered trademark Tylenol®
Regular Strength.

Manufactured by: Time-Cap Labs, Inc.,
7 Michael Avenue, Farmingdale, NY 11735
JAPR 12/24



LOT:

EXP. DATE:

PEEL HERE

Varnish Omit Area

3 147 9 9 4 8 3 1 4 6 9 0 1 1

ACETAMINOPHEN

acetaminophen tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-340	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
Inactive Ingredients				
Ingredient Name			Strength	
POVIDONE (UNII: FZ989GH94E)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	TCL340	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-340-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/17/2018	
2	NDC:49483-340-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/03/2021	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		12/15/2011	

Labeler - TIME CAP LABS INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABS INC		037052099	manufacture(49483-340)