

**TUSSIN CF MULTI-SYMPTOM COLD- dextromethorphan hydrobromide,
guaifenesin, phenylephrine hydrochloride liquid
DeMoulas Market Basket**

Tussin CF - 504

DRUG FACTS

**Active ingredients
(in each 5 mL tsp)**

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Phenylephrine HCL, USP 5 mg

Purposes

Cough Suppressant

Expectorant

Nasal Decongestant

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses

- help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use Ask a doctor or pharmacist before use

- heart disease
- high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days, or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Directions

- Do not take more than 6 doses in any 24 hour period
- This adult strength product is not intended for use in children under 12 years of age

Age

adults and children 12 years and over

children under 12

Dose

2 teaspoons every 4 hours

do not use

Other information

- **Keep carton for full Direction for use.**
- store at 20-25 ° C (68-77 ° F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients

anhydrous citric acid, FD and C red no. 40, glycerin, menthol, e, natural and artificial flavor, propylene glycol purified water, sodium benzoate sorbitol solution, sucralose

Questions?

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Product Label

"MORE FOR YOUR DOLLAR"

MARKET BASKET®

***COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD MULTI-SYMPTOM COLD**

Tussin CF Multi-Symptom Cold

Dextromethorphan HBr/ Guaifenesin / Phenylephrine HCL

**COUGH SUPPRESSANT
EXPECTORANT/
NASAL DECONGESTANT**

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Relieves:

- Cough
- Mucus
- Nasal Congestion

NON-DROWSY

ADULT


for ages 12 and over

8 FL OZ (237 mL)

*This product is not manufactured or distributed by Pfizer, owner of the trademark Robitussin ®

DISTRIBUTED BY DEMOULAS SUPERMARKETS INC. 875 EAST STREET TEWKSBURG, MA 01876 BX-013





DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Active Ingredients (in each 5 mL tsp)

Purpose

Dextromethorphan HBr, USP 10 mg.....	Cough Suppressant
Guaifenesin, USP 100 mg.....	Expectorant
Phenylephrine HCl, USP 5 mg.....	Nasal Decongestant

Uses ■ helps loosen phlegm (mucus) and bronchial secretions to drain bronchial tubes
 ■ temporarily relieves these symptoms occurring with a cold: ■ nasal congestion
 ■ cough due to minor throat and bronchial irritation.

Warnings

Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Stop use and ask a doctor ■ you get nervous, dizzy or sleepless ■ symptoms do not get better within 7 days or are accompanied by fever ■ cough lasts more than 7 days, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 6 doses in any 24 hour period.

Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use

Other Information ■ See carton for for full labeling ■ store at 20-25°C (68-77°F)
 ■ do not refrigerate ■ dosage cup provided ■ sodium 3 mg per teaspoonful

Inactive Ingredients

inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**.

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®.

DISTRIBUTED BY

DEMOULAS SUPERMARKETS INC.

375 East Street

FEWKSBURY, MA 01876

LR-032

LOT:

EXP:

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)	Purposes
Dextromethorphan HBr, USP 10 mg	Cough Suppressant
Guaifenesin, USP 100 mg	Expectorant
Phenylephrine HCl, USP 5 mg	Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age

AGE	DOSE
Adults and children 12 years and over	2 teaspoons every 4 hours
Children under 12 years	do not use

Other information ■ **Keep carton for full Direction for use**

- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful


Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®
 DISTRIBUTED BY DEMOULAS SUPERMARKETS INC.
 875 EAST STREET TEWKSBURY, MA 01876

BX-013
Rev 01

res



DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Active Ingredients (in each 5 mL tsp)	Purpose
Dextromethorphan HBr, USP 10 mg.....	Cough Suppressant
Guaifenesin, USP 100 mg.....	Expectorant
Phenylephrine HCl, USP 5 mg.....	Nasal Decongestant

Uses ■ helps loosen phlegm (mucus) and bronchial secretions to drain bronchial tubes
 ■ temporarily relieves these symptoms occurring with a cold: ■ nasal congestion
 ■ cough due to minor throat and bronchial irritation.

Warnings

Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

Stop use and ask a doctor ■ you get nervous, dizzy or sleepless ■ symptoms do not get better within 7 days or are accompanied by fever ■ cough lasts more than 7 days, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 6 doses in any 24 hour period.

Age	Dose
adults and children 12 years and over	2 teaspoons every 4 hours
children under 12	do not use

Other Information ■ See carton for for full labeling ■ store at 20-25°C (68-77°F)

■ do not refrigerate ■ dosage cup provided ■ sodium 3 mg per teaspoonful

Inactive Ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**.

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®.

DISTRIBUTED BY

DEMOULAS SUPERMARKETS INC.

375 East Street

FEWKSBURY, MA 01876

LOT:

LR-032

EXP:

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)	Purposes
Dextromethorphan HBr, USP 10 mg	Cough Suppressant
Guaifenesin, USP 100 mg	Expectorant
Phenylephrine HCl, USP 5 mg	Nasal Decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age

AGE	DOSE
Adults and children 12 years and over	2 teaspoons every 4 hours
Children under 12 years	do not use

Other information ■ **Keep carton for full Direction for use**


- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients anhydrous citric acid, FD&C red no. 40, glycerin, menthol, natural & artificial flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin®
DISTRIBUTED BY DEMOULAS SUPERMARKETS INC.
 875 EAST STREET TEWKSBURY, MA 01876

BX-013
Rev 01



TUSSIN CF MULTI-SYMPOTOM COLD

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53942-504

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-504-26	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/20/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/20/2012	

Labeler - DeMoulas Market Basket (007869647)

Registrant - DeMoulas Market Basket (007869647)

Establishment

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(53942-504)