GELATO TOPICAL ANESTHETIC- benzocaine gel Mycone Dental Supply Co., Inc DBA Keystone Industries and Deepak Products Inc

Active Ingredients

Benzocaine 200mg (in each g)

Purpose:

Oral Anesthetic

Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings:

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away. **If pregnant or breast feeding,** ask a physician before use.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Keep out of reach of children.

Other Information:

Store at 59°-86°F (15°-30°C). Protect from freezing.

Inactive Ingredients:

Flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow#5 (tartrazine), yellow #6, as a color additive.

Questions or comments?

1-800-333-3131 or

keystoneindustries.com



mouth symptoms do not improve f swelling, rash or fever develops, directed by a physician. If sore redness persists or worsens; or in 7 days; irritation, pain, or

see your physician promptly. for teething

Do not exceed recommended in children under 2 years of age contact occurs, flush with water. When using this product Avoid contact with eyes. If

> may cause methemoglobinemia, a serious condition that must be

warning: Use of this product

Methemoglobinemia

Warnings

If pregnant or breast feeding, Poison Control Center right away. dosage. If more than used for pain is accidentally swallowed, get medical help or contact a ask a physician before use.

Keep out of reach of children Directions

attention if you or a child in your

occur even if you have used this

product before. Stop use and

seek immediate medical

reduces the amount of oxygen

treated promptly because it

carried in the blood. This can

 Apply only amount needed to children under 2 years of age: the oral mucosa to prevent or relieve pain.

pale, gray or blue colored skin

care develops:

do not use

Store at 59° - 86° F (15 - 30°C). Other information

dizziness or lightheadedness

 shortness of breath rapid heart rate

 headache cvanosis)

fatique or lack of energy

Protect from freezing.

(tartrazine), yellow #6, as a color blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, sodium saccharin. May contain flavoring, PEG 3350, PEG 400, nactive ingredients Allergy Alert: Do not use on

allergies to local anesthetics

patients with a history of

such as procaine, butacaine

benzocaine or other "caine'

anesthetics.

 for more than 7 days unless Do not use

Questions or comments?

additive.

keystoneindustries.com

-800-333-3131 or

GELATO TOPICAL ANESTHETIC

For oral mucosal use only, as

Oral Anesthetic

Purpose

directed by dentist. For the

temporary relief of pain due

to minor dental procedures.

benzocaine gel

Product Information

Benzocaine 200 mg (in each g)

Active ingredients

Drug Facts

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68400-354

Route of Administration DENTAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZ OCAINE** 200 mg in 1 g

Inactive Ingredients

Strength **Ingredient Name**

POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics			
Color	orange	Score	
Shape		Size	
Flavor	MANGO	Imprint Code	
Contains			

Ш	Packaging				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1 NDC:68400-354-	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	07/01/2013	

Labeler - Mycone Dental Supply Co., Inc DBA Keystone Industries and Deepak Products Inc (014769301)

Establishment				
Name	Address	ID/FEI	Business Operations	
Keystone Industries		014769301	manufacture(68400-354) , label(68400-354)	

Revised: 12/2023 Mycone Dental Supply Co., Inc DBA Keystone Industries and Deepak Products Inc