

**DAYTIME COLD AND FLU MULTI SYMPTOM RELIEF- acetaminophen,
dextromethorphan hydrobromide, phenylephrine hydrochloride liquid
Lidl US, LLC**

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - fever
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount.
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- a sodium-restricted diet
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough accompanied by excessive phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if

you are taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage.

Stop use and ask a doctor if

- redness or swelling is present
- new symptoms occur
- nervousness, dizziness or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs, with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause

liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL=milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **When using Day Time and Night Time products, carefully read each label to ensure correct dosing**

Other information

- **each 15 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate

Inactive ingredients

citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

*Compare to the active ingredients of Vicks® DayQuil® Cold & Flu

RELIEVES

Aches Fever & Sore Throat Cough Nasal Congestion

Non-Drowsy

Multi-Symptom

Daytime

Cold & Flu

RELIEF

Acetaminophen 325 mg

Pain reliever/Fever reducer

Dextromethorphan HBr 10 mg

Cough suppressant

Phenylephrine HCl 5 mL

Nasal decongestant

For ages 6 years and over

Alcohol-free

Antihistamine-free

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:

Lidl US, LLC

3500 S. Clark Street

Arlington, VA 22202

www.lidl.com

Package Label

Drug Facts (continued)

or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see Overdose warning)
 ■ do not take more than 4 doses in any 24-hour period
 ■ measure only with dosing cup provided. Do not use any other dosing device.
 ■ mL = milliliter ■ keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information ■ each 15 mL contains: sodium 12 mg
 ■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

Discard Seal, Empty & Replace Cup
 DISTRIBUTED BY LIDL US, LLC
 3500 S. Clark Street
 Arlington, VA 22202
 PLD-A380D LB004447 7710-005 ADO
 QUESTIONS? 1-877-753-3935
 www.lidl.com

PLASTIC BOTTLE PLASTIC CUP
 how2recycle.info
 2073 5913

*Compare to the active ingredients in Vicks® DayQuil® Cold & Flu
 NDC 71141-137-32

RELIEVES
 Aches
 Fever &
 Sore Throat
 Cough
 Nasal
 Congestion

**Non-Drowsy
 Multi-Symptom**

**Daytime
 cold & flu
 RELIEF**

Acetaminophen 325 mg
 Pain reliever/Fever reducer
 Dextromethorphan HBr 10 mg
 Cough suppressant
 Phenylephrine HCl 5 mg
 Nasal decongestant

For ages 6 years and over
 Alcohol-free
 Antihistamine-free
 12 FL OZ (355mL)

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredients (in each 15 mL)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms
 ■ minor aches and pains ■ headache ■ fever ■ sore throat
 ■ nasal congestion ■ cough due to minor throat and bronchial irritation

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease
 ■ high blood pressure ■ thyroid disease ■ diabetes ■ a sodium-restricted diet
 ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur
 ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children)

LIDL Daytime Cold and Flu Relief

DAYTIME COLD AND FLU MULTI SYMPTOM RELIEF

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71141-137
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71141-137-32	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/28/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/28/2017	

Labeler - Lidl US, LLC (079389709)

Revised: 4/2024

Lidl US, LLC