

RYMED- dexchlorpheniramine maleate and phenylephrine hydrochloride tablet, coated
EDWARDS PHARMACEUTICALS, INC.

RYMED TABLETS

Drug Facts

| <i>Active Ingredients (in each tablet)</i> | <i>Purpose</i> |
|---|-----------------------|
| Dexchlorpheniramine Maleate 2 mg | Antihistamine |
| Phenylephrine HCl 10 mg | Nasal Decongestant |

Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

Warnings

- **Do not exceed recommended dosage.**

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by a fever
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed recommended dosage.

| | |
|---|---|
| Adults and children 12 years of age and over: | 1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor |
| Children 6 to under 12 years of age: | 1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor |
| Children under 6 years of age: | Consult a doctor. |

Inactive ingredients

Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate

Questions or Comments?

Call 1-800-543-9560

PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label

E

NDC 0485-0080-01

RYMED TABLETS

Antihistamine • Nasal Decongestant

Each tablet contains:

Dexchlorpheniramine Maleate 2 mg
Phenylephrine HCl 10 mg
This bottle is not to be dispensed to consumer.
Dispense in a tight, light-resistant container
with a child-resistant cap.

Store at 59°-86° (15°-30°C) [see USP Controlled Room Temperature].
Tamper evident by foil seal under cap.
Do not use if foil seal is broken or missing.

Manufactured for:
EDWARDS
Pharmaceuticals, Inc.
Ripley, MS 38663

100 tablets

Drug Facts

Active Ingredients Purpose (in each tablet)

Dexchlorpheniramine
Maleate 2 mg Antihistamine
Phenylephrine HCl
10 mg Nasal Decongestant

Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes
- nasal congestion ■ reduces swelling of nasal passages

Peel Here



NDC 0485-0080-01

RYMED TABLETS

Antihistamine • Nasal Decongestant

Each tablet contains:

Dexchlorpheniramine Maleate 2 mg
Phenylephrine HCl 10 mg

This bottle is not to be dispensed to consumer.
Dispense in a tight, light-resistant container
with a child-resistant cap.

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature].
Tamper evident by foil seal under cap.
Do not use if foil seal is broken or missing.

Manufactured for:
EDWARDS
Pharmaceuticals, Inc.
Ripley, MS 38663

100 tablets



Lot: _____
Exp. Date: _____

Drug Facts (continued)

Warnings

■ **Do not exceed recommended dosage.**

Do not use this product

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ a breathing problem such as emphysema or chronic bronchitis
■ glaucoma ■ heart disease
■ high blood pressure ■ thyroid disease ■ diabetes mellitus
■ difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

■ excitability may occur, especially in children
■ may cause drowsiness
■ alcohol, sedatives and tranquilizers may increase the drowsiness effect
■ avoid alcoholic beverages
■ use caution when driving a

Drug Facts (continued)

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur
■ symptoms do not improve within 7 days or are accompanied by a fever ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed recommended dosage.

| | |
|---|---|
| Adults and children 12 years of age and over: | 1 tablet every 4 hours, not to exceed 6 tablets in 24 hours, or as directed by a doctor |
| Children 6 to under 12 years of age: | 1/2 tablet every 4 hours, not to exceed 3 tablets in 24 hours, or as directed by a doctor |
| Children under 6 years of age: | Consult a doctor. |

Inactive ingredients

Magnesium Stearate,
Microcrystalline Cellulose, Sodium Starch Glycolate

Questions or Comments?

RYMED

dexchlorpheniramine maleate and phenylephrine hydrochloride tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0485-0080 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------------|----------|
| DEXCHLORPHENIRAMINE MALEATE (UNII: B10YD955QW) (DEXCHLORPHENIRAMINE - UNII:3Q9Q0B929N) | DEXCHLORPHENIRAMINE MALEATE | 2 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | white | Score | 2 pieces |
| Shape | OVAL | Size | 12mm |
| Flavor | | Imprint Code | R;1 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0485-0080-01 | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 11/15/2011 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 11/15/2011 | |

Labeler - EDWARDS PHARMACEUTICALS, INC. (195118880)

Revised: 8/2024

EDWARDS PHARMACEUTICALS, INC.