

ORAJEL INSTANT PAIN RELIEF SEVERE- benzocaine 20%, menthol, benzalkonium chloride gel

Church Dwight Co., Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Orajel SEVERE Toothache & Gum Relief Plus

TRIPLE MEDICATED

Active ingredients

Benzocaine 20%

Menthol 0.5%

Benzalkonium Chloride 0.1%

Purpose

Oral pain reliever

Oral Pain reliever

Antiseptic

Use

- for the temporary relief of pain due to toothaches
- to help protect against infection in minor oral irritation

Warnings

⚠**Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

⚠**Do not use**

- more than directed
- for more than 7 days unless directed by a physician or healthcare provider

⚠**Stop use and ask a physician if**

- swelling, rash or fever develops
- irritation, pain or redness persists or worsens
- symptoms do not improve in 7 days

KEEP OUT OF REACH OF CHILDREN:

⚠In case of overdose or allergic reaction, get medical help or contact a Poison Control Center right away

⚠**Directions** - cut open tip of tube on score mark

- **Adults and children 2 years of age and over** - Apply a small amount of product to the cavity and around gum surrounding the teeth. Use up to 4 times daily or as directed by a physician or healthcare provider
- **Children under 12 years of age** - Should be supervised in the use of this product
- **Children under 2 years of age** - Ask a physician or healthcare provider

Other information

- do not use if tube tip is cut prior to opening
- this preparation is intended for use in cases of toothache, only as a temporary expedient until a physician can be consulted
- do not use continuously
- Orajel Severe Pain Formula will stay in place for extended duration of relief
- avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

Inactive ingredients

blue 1, cellulose gum, gelatin, methyl salicylate, mineral oil, pectin, petrolatum, polyethylene glycol, sodium saccharin

Questions or comments?

call us at ☎1-800-952-5080☎ M-F 9am-5pm ET or visit our website at www.oraljel.com

#1

ORAL PAIN

Reliever for Toothache

Now with 3X More

Active Ingredients vs. Store Brand

Orajel™

Instant Pain Relief

SEVERE

Toothache & Gum Relief Plus

TRIPLE MEDICATED

20% Benzocaine to Relieve Oral Pain

Antiseptic to Help Prevent Infection

Menthol to Soothe

FAST-ACTING

GEL

Oral Pain Reliever/Antiseptic NET WT. 0.25 OZ (7.0g)

OJFC-32503-06



RECEIVED FOR
TOOTHACHE

Orajel™

Instant Pain Relief

SEVERE

Toothache & Gum Relief Plus

TRIPLE MEDICATED

- ✓ 20% Benzocaine to Relieve Oral Pain
- ✓ Antiseptic to Help Prevent Infection
- ✓ Menthol to Soothe

SAFETY SEALED
TUBE TIP

Orajel
Instant Pain Relief
SEVERE
Toothache & Gum Relief Plus

TRIPLE MEDICATED

- ✓ 20% Benzocaine to Relieve Oral Pain
- ✓ Antiseptic to Help Prevent Infection
- ✓ Menthol to Soothe

ORAL PAIN RELIEVER/ANTISEPTIC

NET WT 0.25 OZ (7.0 g)



FAST-ACTING
GEL

Orajel
Instant Pain Relief
SEVERE
Toothache & Gum Relief Plus

Church & Dwight Co., Inc., Ewing, NJ 08620
©2013 Church & Dwight Co., Inc.
ORAJEL is a trademark of Church & Dwight Co., Inc.
The makers of Orajel® do not manufacture store brand
oral pain products.
Made in USA with US and Foreign Components
OJFC-32503-06 70506294



3 10310 32503 9

CHDW0003
090328Q1

Drug Facts

Active ingredients

Benzocaine 20%
Menthol 0.5%
Benzalkonium Chloride 0.1%
Oral pain reliever
Antiseptic

Purpose

Oral pain reliever
Antiseptic

Use ■ For the temporary relief of pain due to toothaches ■ To help protect against infection in minor oral irritation

Warnings ■ Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

Do not use ■ more than directed ■ for more than 7 days unless directed by a physician or healthcare provider

Stop use and ask a physician if ■ swelling, rash or fever develops ■ irritation, pain or redness persists or worsens ■ symptoms do not improve in 7 days

Keep out of reach of children. In case of overdose or allergic reaction, get medical help or contact a Poison Control Center right away.

Directions ■ Cut open tip of tube on score mark

Adults and children 2 years of age and over

Apply a small amount of product to the cavity and around gum surrounding the tooth. Use up to 4 times daily or as directed by a physician or healthcare provider

Children under 12 years of age

Should be supervised in the use of this product

Children under 2 years of age

Ask a physician or healthcare provider

Other information ■ Do not use if tube tip is cut prior to opening. ■ The preparation is intended for use in case of toothache, only as a temporary expedient until a physician can be consulted

■ Do not use continuously. ■ Orajel Severe Pain Formula will stay in place for extended duration of relief ■ Avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

Inactive ingredients Blue 1, cellulose gum, glycerin, methyl salicylate, mineral oil, pectin, petroleum, polyethylene glycol, sodium saccharin

Questions or comments? Call us at 1-800-952-5000 M-F 9am-5pm ET or visit our website at www.orajel.com

■ Yes, single medicated oral pain products

benzocaine 20%, menthol, benzalkonium chloride gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10237-752
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1 mg in 1 g
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	50 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
METHYL SALICYLATE (UNII: LAV5U5022Y)	
MINERAL OIL (UNII: T5L8T28FGP)	
PECTIN (UNII: 89NA02M4RX)	
PETROLATUM (UNII: 4T6H12BN9U)	
POLYETHYLENE GLYCOL 1000 (UNII: U076Q6Q621)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10237-752-25	1 in 1 CARTON	12/26/2016	02/27/2017
1		7 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	11/15/2013	

Labeler - Church Dwight Co., Inc. (001211952)

Establishment

Name	Address	ID/FEI	Business Operations
Church Dwight Co., Inc.		043690812	manufacture(10237-752)

