

**WINCO FOODS MAXIMUM STRENGTH ORAL ANALGESIC- benzocaine gel**  
**WinCo Foods, LLC**

-----  
**WinCo Max. Strength Oral Gel 0.42oz (24115)**

<b>Active ingredient</b>	<b>Purpose</b>
Benzocaine 20%.....	Oral pain reliever

**Uses**

For the temporary relief of pain due to:

- toothache
- canker sores
- gum pain
- denture pain
- cheek bites

**Warnings**

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy Alert:** do not use this product if your child has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

**Do not use**

- more than directed
- for more than 7 days unless directed by a doctor
- for teething
- in children under 2 years of age

**Stop use and ask a doctor if**

- sore mouth symptoms do not improve in 7 days
- irritation, pain or redness does not go away
- swelling, rash or fever develops

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- Adults and children 2 years of age and older: Remove cap and cut tip of tube. Apply a

small amount of product to the cavity and around gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor.

- Children under 12 years of age: Should be supervised in the use of this product.
- Children under 2 years of age: Do not use.

**Other information**

- store at room temperature 20-25°C (68-77°F)
- this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor can be consulted. Do not use continuously.

**Inactive ingredients**

glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

**DISTRIBUTED BY:**

WINCO FOODS, LLC

BOISE, ID 83704

MADE IN CHINA

WinCo  
FOODS

# MAXIMUM STRENGTH Oral Analgesic Gel

BENZOCAINE 20% / ORAL PAIN RELIEVER

WinCo  
FOODS

## MAXIMUM STRENGTH Oral Analgesic Gel

BENZOCAINE 20% / ORAL PAIN RELIEVER

Provides Immediate Relief  
From: Toothache, Denture  
& Gum Irritations,  
Cold & Canker Sores

NET WT 0.42 OZ (11.9 g)

GLUE AREA  
No Ink /  
No Varnish

GLUE AREA  
No Ink /  
No Varnish

WF2002



DISTRIBUTED BY:  
WINCO FOODS, LLC,  
BOISE, ID 83704  
MADE IN CHINA



The Supermarket Low Price Leader®

100% SATISFACTION  
GUARANTEE

Comments & Questions?  
800-824-1706 · www.wincofoods.com

Lot & Exp.

24115-ZDP-BX-R1  
OC-MD-ZDPI05-R3

WinCo  
FOODS  
MAXIMUM STRENGTH  
Oral Analgesic  
Gel

<b>Drug Facts</b>	
<b>Active ingredient</b>	Benzocaine 20%.....Oral pain reliever
<b>Uses</b>	■ for the temporary relief of pain due to: ■ toothache ■ canker sores ■ gum pain ■ denture pain ■ cheek bites
<b>Warnings</b>	Methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy <b>Allergy Alert:</b> do not use this product if your child has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.
<b>Do not use</b>	■ more than directed ■ for more than 7 days unless directed by a doctor ■ for teething ■ in children under 2 years of age
<b>Stop use and ask a doctor if</b>	■ sore mouth symptoms do not improve in 7 days ■ irritation, pain or redness does not go away ■ swelling, rash or fever develops
If pregnant or breast-feeding, ask a health professional for advice.	
<b>Drug Facts (continued)</b>	
<b>Purpose</b>	Oral pain reliever
<b>Keep out of reach of children.</b>	If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
<b>Directions</b>	Adults and children 2 years of age and older: Remove cap and cut tip of tube. Apply a small amount of product to the cavity and around gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor. Children under 12 years of age: Should be supervised in the use of this product. Children under 2 years of age: Do not use.
<b>Other information</b>	■ store at room temperature 20-25°C (68-77°F) ■ this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor can be consulted. Do not use continuously.
<b>Inactive ingredients</b>	polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol
<b>Questions?</b>	Monday - Friday 9am - 4pm MST Call 800-824-1706

TAMPER EVIDENT: DO NOT USE IF OUTER CARTON IS OPEN OR IF TUBE TIP IS CUT PRIOR TO OPENING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

# WINCO FOODS MAXIMUM STRENGTH ORAL ANALGESIC

benzocaine gel

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-198
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>POLYETHYLENE GLYCOL 4000</b> (UNII: 4R4HFI6D95)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>PEPPERMINT OIL</b> (UNII: AV092KU4JH)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:67091-198-11	1 in 1 CARTON	11/08/2016	
1		11.9 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M022	12/01/2014	

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 10/2025

WinCo Foods, LLC