WINCO FOODS MAXIMUM STRENGTH ORAL ANALGESIC- benzocaine gel WinCo Foods, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

WinCo Max. Strength Oral Gel 0.42oz (24115)

Active ingredient Purpose Benzocaine 20%...... Oral pain reliever

Uses

For the temporary relief of pain due to:

- toothache
- canker sores
- gum pain
- denture pain
- cheek bites

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allery Alert: do not use this product if your child has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- more than directed
- for more than 7 days unless directed by a doctor
- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain or redness does not go away
- swelling, rash or fever develops

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison

Control Center (1-800-222-1222) right away.

Directions

- Adults and children 2 years of age and older: Remove cap and cut tip of tube. Apply a small amount of product to the cavity and around gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor.
- Children under 12 years of age: Should be supervised in the use of this product.
- Children under 2 years of age: Do not use.

Other information

- store at room temperature 20-25°C (68-77°F)
- this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor can be consulted. Do not use continuously.

Inactive ingredients

glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

DISTRIBUTED BY:

WINCO FOODS, LLC BOISE, ID 83704 MADE IN CHINA



MAXIMUM STRENGTH Oral Analgesic Gel

BENZOCAINE 20% / ORAL PAIN RELIEVER

WinCo

MAXIMUM STRENGTH Oral Analgesic Gel

BENZOCAINE 20% / ORAL PAIN RELIEVER

Provides Immediate Relief From: Toothache, Denture & Gum Irritations,

Cold & Canker Sores

NET WT 0.42 OZ (11.9 g)



DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

MADE IN CHINA



The Supermarket Low Price Leader®

SATISFACTION GUARANTEE

Comments & Questions? 800-824-1706 · www.wincofoods.com

TSM mq4 - ma9 yam - 4pm MST Questions? Call 800-824-1706

polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol Inactive ingredients glycerin, peppermint oil,

can be consulted. Do not use continuously. (68-77-F) ■ this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor and the preparation of the preparati O°5S-0S arufer information ■ store at room temperature 20-25°C

/ears of age.	
Children under 2	Do not use.
/ears of age.	broduct.
Children under 12	Should be supervised in the use of this
	dentist or doctor.
	up to 4 times daily or as directed by a
older.	around gum surrounding the teeth. Use
/ears of age and	small amount of product to the cavity and
Adults and children 2	Remove cap and cut tip of tube. Apply a

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away. erore use. Drug Facts (continued)

Directions

It pregnant or breast-feeding, ask a health professional

away = swelling, rash or fever develops Stop use and ask a doctor if ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain or redness does not go

unless directed by a doctor

for teething in children under Do not use ■ more than directed ■ for more than 7 days

of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

in your care develops:

Stop use and seek immediate medical attention if you or a child blood. This can occur even if you have used this product before. methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in Methemoglobinemia warning: Use of this product may cause Warnings

■ csuker sores ■ gum pain ■ denture pain ■ cheek bites **NSGS** ■ for the temporary relief of pain due to: ■ toothache

Purpose Active ingredient Drug Facts

TAMPER EVIDENT: DO NOT USE IF OUTER CARTON IS OPEN OR IF TUBE TIP IS CUT PRIOR TO OPENING.

KEEP OUTER CARTON FOR COMPLETE VARNINGS AND PRODUCT INFORMATION.

24115-ZDP-BX-R1 OC-MD-ZDP105-R3

WINCO FOODS MAXIMUM STRENGTH ORAL ANALGESIC

benzocaine gel

Droduct	Inform	\ation
Product		ıatıvı

Product Type HUMAN OTC DRUG Item Code (Source) NDC:67091-198

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) BENZOCAINE 200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
PEPPERMINT OIL (UNII: AV092KU4JH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-198- 11	1 in 1 CARTON	11/08/2016	
1		11.9 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

ting Start Marketing E	
Date Date	End
4	
	1

Labeler - WinCo Foods, LLC (056098817)

Revised: 3/2022 WinCo Foods, LLC