

EASYLAX- polyethylene glycol powder
Marc Glassman, Inc.

Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350,
17 g (cap filled to line)

Purpose

Osmotic laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol.

Do not use

If you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

When using this product

you may have loose, watery, more frequent stools.

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
 - fill to top of line in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounce of beverage (cold, hot, or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredient

none

Principal Display Panel

*Compare to the active ingredient in MIRALAX®

EasyLAX

ORIGINAL STRENGTH

Polyethylene Glycol 3350

Powder for Oral Solution

Osmotic Laxative

- Relieves Occasional Constipation (irregularity)
- Softens stool
- Dissolves in any beverage
- Sugar Free
- Unflavored, grit-free powder

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Polyethylene glycol 3350 increases frequency of bowel movements and softens the stool

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MIRALAX®.

Distributed by: Marc Glassman, Inc.

West 130th Street, Cleveland, OH 44130

Package Label

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091077	08/31/2016	12/30/2021

Labeler - Marc Glassman, Inc. (094487477)

Revised: 11/2019

Marc Glassman, Inc.