

**EASOTIC OTIC- hydrocortisone aceponate, miconazole nitrate, and gentamicin sulfate suspension**  
**Virbac AH, Inc.**

-----  
**EASOTIC®**  
**Otic suspension**  
**(hydrocortisone aceponate, miconazole nitrate, gentamicin sulfate)**

Anti-inflammatory, antifungal, and antibacterial

**Rx**

*For Otic Use in Dogs Only*

**CAUTION**

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**DESCRIPTION**

EASOTIC® suspension contains 1.11 mg/mL hydrocortisone aceponate, 15.1 mg/mL miconazole nitrate and 1.5 mg/mL gentamicin sulfate. The inactive ingredient is a semi-liquid petroleum jelly.

**INDICATIONS**

EASOTIC® suspension is indicated for the treatment of otitis externa in dogs associated with susceptible strains of yeast (*Malassezia pachydermatis*) and bacteria (*Staphylococcus pseudintermedius*).

**DOSAGE AND ADMINISTRATION**

Verify that the tympanic membrane is intact. **Shake well before each use.**

**Priming the canister**

Prior to the first use of the dosing canister, prime the pump by depressing the pump 1 to 2 times to fill the clear canula (tip) with a full dose of product.

Carefully insert the canula into the affected external ear canal(s) and apply 1 mL (a single pump) of suspension once per day for 5 days. Wash hands after usage.

**CONTRAINDICATIONS**

Do not use in dogs with known tympanic membrane perforation.

EASOTIC® suspension is contraindicated in dogs with known or suspected hypersensitivity to corticosteroids, imidazole antifungals, or aminoglycoside antibiotics.

**WARNINGS**

**Human Warnings**

Not for use in humans. Keep this and all drugs out of reach of children. In case of accidental skin contact, wash area thoroughly with water. Avoid contact with eyes.

Humans with known or suspected hypersensitivity to hydrocortisone, aminoglycoside antibiotics, orazole antifungals should not handle this product.

In case of accidental ingestion by humans, contact a physician immediately. Physicians may contact a Poison Control Center for advice concerning cases of ingestion by humans.

### **Animal Warnings**

As a class, aminoglycoside antibiotics are associated with ototoxicity, vestibular dysfunction and renal toxicity. The use of EASOTIC<sup>®</sup> suspension in a dog with a damaged tympanic membrane can result in damage to the structures of the ear associated with hearing and balance or in transmission of the infection to the middle or inner ear. Immediately discontinue use of EASOTIC<sup>®</sup> suspension if hearing loss or signs of vestibular dysfunction are observed during treatment (see **ADVERSE REACTIONS**).

### **PRECAUTIONS**

Do not administer orally.

Concurrent administration of potentially ototoxic drugs should be avoided.

Use with caution in dogs with impaired hepatic or renal function (see **ANIMAL SAFETY**).

Long-term use of topical otic corticosteroids has been associated with adrenocortical suppression and iatrogenic hyperadrenocorticism in dogs (see **ANIMAL SAFETY**).

The safe use of EASOTIC<sup>®</sup> suspension in dogs used for breeding purposes, during pregnancy, or in lactating bitches, has not been evaluated.

### **ADVERSE REACTIONS**

In a field study conducted in the United States (see **EFFECTIVENESS**), there were no adverse reactions reported in 145 dogs administered EASOTIC<sup>®</sup> suspension.

In foreign market experience, reports of hearing loss and application site erythema have been received. In most reported cases, the hearing loss and erythema were transient and resolved with discontinuation of EASOTIC<sup>®</sup> suspension.

To report suspected adverse drug events, contact Virbac at 800-338-3659 or the FDA at 1-888-FDA-VETS.

For technical assistance or to obtain a Material Safety Data Sheet, call Virbac at 800-338-3659.

### **PHARMACOLOGY**

Hydrocortisone aceponate is a glucocorticoid with anti-inflammatory effects. Miconazole nitrate is an imidazole antifungal. Gentamicin sulfate is an aminoglycoside antibiotic.

In the target animal safety study, hydrocortisone aceponate, miconazole and gentamicin were shown to be systemically absorbed from the ears of healthy dogs (see **ANIMAL SAFETY**); increased systemic absorption may be observed in inflamed ears.

### **MICROBIOLOGY**

The compatibility and additive effect of each of the components in EASOTIC<sup>®</sup> suspension was demonstrated in a component effectiveness and non-interference study. An *in vitro* study of organisms collected from clinical cases of otitis externa in dogs and from dogs enrolled in the clinical effectiveness study for EASOTIC<sup>®</sup> suspension determined that miconazole nitrate and gentamicin sulfate inhibit the growth of bacteria and yeast commonly associated with otitis externa in dogs. No consistent synergistic or antagonistic effect of the two antimicrobials was demonstrated. The addition of hydrocortisone aceponate to the combination did not impair antimicrobial activity to any clinically-significant extent.

In a field study (see **EFFECTIVENESS**), the minimum of 10 isolates from successfully treated cases was met for *S. pseudintermedius* and *M. pachydermatis*.

## **EFFECTIVENESS**

The effectiveness of this drug was evaluated in 157 dogs with otitis externa. The study was a double-masked field study with a placebo control. One hundred and four dogs were treated with EASOTIC<sup>®</sup> suspension and 53 dogs were treated with the placebo control. Treatment was administered once daily for 5 consecutive days to the affected ear(s). The dogs were evaluated at 4 different intervals over the course of 1 month to determine response to therapy. The 6 clinical signs evaluated were: malodor, aural discharge, pruritus, erythema, swelling and pain. The individual clinical scores were assigned based on the severity of each sign. Success was based on clinical improvement at Day 28 ±2 days. The success rates of the 2 groups were significantly different (p=0.0179); 68.5% of dogs administered EASOTIC<sup>®</sup> suspension were successfully treated, compared to 21.8% of the dogs in the placebo control group.

## **ANIMAL SAFETY**

In the target animal safety study, EASOTIC<sup>®</sup> suspension was administered at 0×, 1×, 3× and 5× the recommended dose for 15 consecutive days (3 times the recommended treatment duration) in laboratory Beagles, with 8 dogs per group. Hypersensitivity reactions in the external ear canal and inner pinnae were seen in all EASOTIC<sup>®</sup> suspension groups and included mild to severe aural erythema (3× group), papules and ulceration (1× and 5× groups), otitis externa (3× and 5× groups), and otitis media (5× group). Renal tubular crystals were present in the cortex and medulla (0×, 1×, 3×, and 5× groups) and mild renal tubular basophilia and atrophy were present in one 5× group dog. Baseline cortisol values and the cortisol response to ACTH stimulation were lower in treated dogs compared to the control dogs. The ACTH stimulation test results are consistent with systemic absorption of topical corticosteroids causing suppression of the hypothalamic-pituitary-adrenal axis. Dogs in the 3× and 5× groups demonstrated elevations in AST and ALP, while dogs in the 1×, 3×, and 5× groups had elevated cholesterol, total protein, and albumin levels. Dogs in the 3× and 5× groups also had higher liver weights and greater food consumption.

## **STORAGE INFORMATION**

Store at temperatures between 20° C-25° C (68° F-77° F), with excursions permitted between 15° C-30° C (59° F-86° F).

## **HOW SUPPLIED**

EASOTIC<sup>®</sup> suspension is supplied in a polyethylene canister, with a soft applicator canula.

Each canister contains ten 1 mL doses.

Made in the U.S.A.

Distributed by:

Virbac AH, Inc.

Fort Worth, TX 76137 USA

NADA 141-330, Approved by FDA.

Revision Date 8/2011

302076-01

## **PRINCIPAL DISPLAY PANEL - 10 mL Carton**

*easOtic*<sup>®</sup>

**(hydrocortisone aceponate 1.11 mg /  
miconazole nitrate 15.1 mg /  
gentamicin sulfate 1.5 mg/mL)**  
Suspension for Dogs

**For otic use in dogs only.**

**CAUTION:**

**Federal law restricts this drug  
to use by or on the order of  
a licensed veterinarian.**

**Net Contents: 10 mL (10 doses)**

**NADA 141-330, Approved by FDA**

***Virbac***

**ANIMAL HEALTH**



## EASOTIC OTIC

hydrocortisone aceponate, miconazole nitrate, and gentamicin sulfate suspension

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:51311-360
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Hydrocortisone aceponate</b> (UNII: 2340UP1L2G) (Hydrocortisone - UNII:WI4X0X7BPJ)	Hydrocortisone aceponate	1.11 mg in 1 mL

<b>Miconazole Nitrate</b> (UNII: VW4H1CYW1K) (Miconazole - UNII:7NNO0D7S5M)	Miconazole Nitrate	15.1 mg in 1 mL
<b>Gentamicin Sulfate</b> (UNII: 8X7386QRLV) (Gentamicin - UNII:T6Z9V48IKG)	Gentamicin Sulfate	1.5 mg in 1 mL

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51311-360-10	10 mL in 1 BOTTLE, PUMP		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141330	02/07/2012	

**Labeler** - Virbac AH, Inc. (131568396)

### Establishment

Name	Address	ID/FEI	Business Operations
Virbac Corporation		829166276	manufacture

### Establishment

Name	Address	ID/FEI	Business Operations
Hovione FarmaCiencia SA		449818434	api manufacture

### Establishment

Name	Address	ID/FEI	Business Operations
FDC Limited Roha Plant		650441301	api manufacture

### Establishment

Name	Address	ID/FEI	Business Operations
Fujian Fukang Pharmaceutical Co. LTD		528180884	api manufacture

Revised: 7/2017

Virbac AH, Inc.