

## LOW DOSE ASPIRIN- aspirin 81 mg delayed release tablets tablet WALGREENS

-----  
**481R WALGREENS 0363 0690 Aspirin 81 mg Tablet**

### DRUG FACTS

#### **Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever

#### **Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric-coated 81 mg Aspirin

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problem
- take a blood thinning (anticoagulant) or steroid drug
- take other drug containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

#### **Ask a doctor before use if**

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for:

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

an allergic reaction occurs. Seek medical help right away.

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present

new symptoms occur

ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**


- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

Store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?** Call **1-877-290-4008**





Walgreens

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

120 TABLETS

NDC 0363-0690-12

## Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

## Purpose

Pain reliever

**Uses** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

## Warnings (See carton for complete warnings information)

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## Directions

■ Drink a full glass of water with each dose ■ Adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ Children under 12 years: consult a doctor

**Other information** ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Do not use if the imprinted foil seal over the mouth of the bottle is cut, torn, broken or missing.** ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

## Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Uses** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include ■ hives ■ facial swelling ■ shock ■ asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## Directions

■ Drink a full glass of water with each dose ■ Adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ Children under 12 years: consult a doctor

## Other information

■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Do not use if the imprinted foil seal over the mouth of the bottle is cut, torn, broken or missing.** ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

## Coating/Print Omit Area

Exp Date:

Lot No.:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE

Walgreens.com ©2022 Walgreen Co.

MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No. : Varnish Omit Area

Exp. Date:

DO NOT USE IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. ■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

ITEM 284619 481R 0222


DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

100% SATISFACTION GUARANTEE







# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

NDC 0363-0690-20

**RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)  
nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

**Warnings (See carton for complete warnings information)**

■ **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

■ **Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

■ **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

■ **Do not use**


- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

■ **Ask a doctor before use if**

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

**WARNING: DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

ITEM 284614 W00000-0000-D



3 11917 13641 7

Lot No.:  
Exp. Date:

**Drug Facts (continued)**

- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if:**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ red stool ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-525-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 6 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- Store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** anhydrous lactose, croscarmellose sodium, colloidal silicon dioxide, croscarmellose sodium, D4C-yellow P10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyvinylpyrrolidone K80, croscarmellose, sodium hydroxide, sodium lauryl sulfate, starch, lact, titanium dioxide, triethyl citrate

**Questions or comments?** Call 1-877-398-4000

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-1122

**SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**DO NOT USE THIS PRODUCT IF THE PREPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.**

**ITEM 284614 W00000-0000-D**

Lot No.:  
Exp. Date:

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

**200**  
TABLETS

W3ORG0422-F  
REV-





NDC 0363-0690-31

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

300

TABLETS

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*

**Purpose**  
Pain reliever

**Nonsteroidal anti-inflammatory drug**

**Uses**  
For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  

- ask your doctor about other uses for enteric-coated 81 mg Aspirin.
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Do not use
  - if you are allergic to aspirin or any other pain reliever/fever reducer
  - if you have ever had an allergic reaction to this product or any of its ingredients
- Ask a doctor before use if
  - you have asthma
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you are taking a diuretic
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- Ask a doctor or pharmacist before use if you are taking a prescription drug for:
  - gout
  - diabetes
  - arthritis
- Stop use and ask a doctor if
  - an allergic reaction occurs. Seek medical help right away
  - you experience any of the following signs of stomach bleeding:
    - feel faint
    - vomit blood
    - have bloody or black stools
    - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Directions**  

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**  

- Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
 anhydrous lactose, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
 Call 1-877-299-4898

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEED  
 MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS 481R 0222

NDC 0363-0690-31

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

300

TABLETS

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*

**Purpose**  
Pain reliever

**Nonsteroidal anti-inflammatory drug**

**Uses**  
For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  

- ask your doctor about other uses for enteric-coated 81 mg Aspirin.
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Do not use
  - if you are allergic to aspirin or any other pain reliever/fever reducer
  - if you have ever had an allergic reaction to this product or any of its ingredients
- Ask a doctor before use if
  - you have asthma
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you are taking a diuretic
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- Ask a doctor or pharmacist before use if you are taking a prescription drug for:
  - gout
  - diabetes
  - arthritis
- Stop use and ask a doctor if
  - an allergic reaction occurs. Seek medical help right away
  - you experience any of the following signs of stomach bleeding:
    - feel faint
    - vomit blood
    - have bloody or black stools
    - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Directions**  

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**  

- Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
 anhydrous lactose, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
 Call 1-877-299-4898

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEED  
 MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS ITEM 285642 W000100-0000-0

NDC 0363-0690-31

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

300

TABLETS

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*

**Purpose**  
Pain reliever

**Nonsteroidal anti-inflammatory drug**

**Uses**  
For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  

- ask your doctor about other uses for enteric-coated 81 mg Aspirin.
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Do not use
  - if you are allergic to aspirin or any other pain reliever/fever reducer
  - if you have ever had an allergic reaction to this product or any of its ingredients
- Ask a doctor before use if
  - you have asthma
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you are taking a diuretic
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- Ask a doctor or pharmacist before use if you are taking a prescription drug for:
  - gout
  - diabetes
  - arthritis
- Stop use and ask a doctor if
  - an allergic reaction occurs. Seek medical help right away
  - you experience any of the following signs of stomach bleeding:
    - feel faint
    - vomit blood
    - have bloody or black stools
    - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Directions**  

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**  

- Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
 anhydrous lactose, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
 Call 1-877-299-4898

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEED  
 MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS ITEM 285642 W000100-0000-0

NDC 0363-0690-31

**Walgreens**

# Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose
Enteric-Coated

300

TABLETS

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*

**Purpose**  
Pain reliever

**Nonsteroidal anti-inflammatory drug**

**Uses**  
For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

**Warnings**  

- ask your doctor about other uses for enteric-coated 81 mg Aspirin.
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- Do not use
  - if you are allergic to aspirin or any other pain reliever/fever reducer
  - if you have ever had an allergic reaction to this product or any of its ingredients
- Ask a doctor before use if
  - you have asthma
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you are taking a diuretic
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- Ask a doctor or pharmacist before use if you are taking a prescription drug for:
  - gout
  - diabetes
  - arthritis
- Stop use and ask a doctor if
  - an allergic reaction occurs. Seek medical help right away
  - you experience any of the following signs of stomach bleeding:
    - feel faint
    - vomit blood
    - have bloody or black stools
    - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Directions**  

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**  

- Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
 anhydrous lactose, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyacrylate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
 Call 1-877-299-4898

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEED  
 MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS ITEM 285642 W000100-0000-0

aspirin 81 mg delayed release tablets tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0690
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

## Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TALC (UNII: 7SEV7J4R1U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

## Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	Heart
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0690-12	1 in 1 CARTON	10/08/2021	
1		120 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0690-32	1 in 1 CARTON	06/15/2022	
2		32 in 1 BOTTLE; Type 0: Not a Combination		

4		Product		
3	NDC:0363-0690-29	2 in 1 CARTON	06/15/2022	
3		120 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0690-20	1 in 1 CARTON	06/15/2022	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0363-0690-31	1 in 1 CARTON	06/15/2022	
5		300 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/08/2021	

**Labeler** - WALGREENS (008965063)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

## Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(0363-0690)

Revised: 1/2025

WALGREENS