NICOTINE TRANSDERMAL PATCHES- aroamas nicotine transdermal patches stop smoking aid patch Ehy Holdings LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

81799-002

Nicotine Transdermal Patches

Active Ingredient(s)

Nicotine 6%.....Stop smoking aid menthol 0.3%.....Stop smoking aid

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your healthcare provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to safer than smoking. However, the risks to your child from this medicine are not full known.

Ask a doctor before use if you have

- *heart disease, recent heartattack, or irregular heartbeat. Nicotine can increase your heartrate.
- *high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- *an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- *stomach ulcer or diabetes.
- *history of seizures.

Ask a doctor or pharmacist before use if you are

- *using an on-nicotine stop smoking drug
- *taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted

Do not use

- *do not wear more than one patch at a time
- *do not cut patch in half or into smaller pieces
- *do not leave patch on for more than 24 hours because if may irritate your skin and loses strength after 24 hours

When using this product

*if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- *skin redness caused by the patch does not go away after four days, or if your skins wells, or you get arash
- *irregular heartbeat or palpitations occur
- *you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- *you have symptoms of an allergic reaction(such as difficulty breathing or rash)

Keep out of reach of children and pets.

Used patches have enough nicotine to poison children and pets.If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch di posal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- *if you are under 18 years of age, ask a doctor before use.*before using this product, read the enclosed self-help guide for complete directions and other information *begin using the patch on your guit day
- *if you smoke more than 10 cigarettes per day, use the following schedule below[]

STEP 1	STEP 2	STEP 3
Use one 21mg patch/day	Use one 14mg patch/day	Use one 7mg patch/day
Weeks 1-6	Weeks 7-8	Weeks 9-10

^{*}If you smoke 10 or less cigarettes per day, do not use Step 1(21mg) .Start with Step 2(14mg) for 6 weeks, then Step 3(7mg) for 2 weeks and then stop.

Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

- *the used patch should be removed and a new one applied to a different skin site at the sametime each day.
- *if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.
- *you may wear the patch for 18 to 24 hours
- *do not wear more than one patch at a time
- *do not cut patch in half or into smaller pieces
- *do not leave patch on for more than 24 hours because if may irritate your skin and

^{*}apply one new patch every 24 hours on skin that is dry, clean and hairless.

^{*}remove backing from patch and immed ately press on to skin. Hold for 10 seconds..wash hands after applying or removing patch.

loses strength after 24 hours

- *to avoid possible burns, remove patch before undergoing and MR(magnetic resonance imaging) procedures
- *it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your healthcare provider.

Other information

If you smoke more than 10 cigarettes per day, start with Step 1. If you smoke 10 or less cigarettes per day, Start with Step 2.

Inactive ingredients

codonopsis pilosula, chrysanthemum, clove, natural tobacco

Package Label - Principal Display Panel

81799-002-01 7mg 1Patch 21PATCH 1BOX





NICOTINE TRANSDERMAL PATCHES

aroamas nicotine transdermal patches - stop smoking aid patch

Product Information				
Product Type	HUMAN OTC DRUG	Item C	ode (Source)	NDC:81799-002
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name Basis of Strength			Strength	

MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.3 mg in 100 mg
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	6 mg in 100 mg

Inactive Ingredients			
Ingredient Name	Strength		
CHRYSANTHEMUM INDICUM FLOWER (UNII: 160ER6U04L)			
CLOVE (UNII: K48IKT5321)			
CODONOPSIS PILOSULA WHOLE (UNII: 76AI4Q0335)			
TOBACCO LEAF (UNII: 6YR2608RSU)			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:81799-002- 01	21 in 1 BOX	03/30/2020		
1		7 mg in 1 PATCH; Type 0: Not a Combination Product			

Marketing Information			
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
part348	03/30/2020		
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date	

Labeler - Ehy Holdings LLC (117322715)

Establishment				
Name	Address	ID/FEI	Business Operations	
Ehy Holdings LLC		117322715	manufacture(81799-002)	

Revised: 1/2023 Ehy Holdings LLC