BISACODYL - bisacodyl suppository Cardinal Health

Leader Bisacodyl Suppositories

Active ingredient (in each suppository)

Bisacodyl USP, 10mg

Purpose

Stimulant Laxative

Uses

- for temporary relief of occasional constipation
- this product generally produces bowel movement in 15 minutes to 1 hour

Warnings

For rectal use only.

Do not use unless directed by a doctor

- if abdominal pain, nausea or vomiting are present
- for longer than 1 week

Ask a Doctor before use if you have

noticed a sudden change in bowel habits that lasts over a perios of 2 weeks

When using this product

it may cause abdominal discomfort, faitness, rectal burning and mild cramps

Stop use and ask a doctor if

you have rectal bleeding or fail to have a bowel movement after using this product. These may indicate a serious condition.

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- detach one suppository from the strip
- remove wrapper before inserting into rectum
- carefully insert one suppository well into the rectum, pointed end first
- retain about 15 to 20 minutes

adults and children 12 years of age and over - 1 suppository once daily children 6 to under 12 years of age - 1/2 suppository once daily children under 6 - do not use

Other information

- Store at room remperature do not to exceed 30°C (86°F)
- Lot No. & Exp. Date: see wrapper or box

Inactive ingredients

hydrogenated vegetable oil

Questions or comments?

(800) 492-5988

Leader Bisacodyl Suppositories

LeaderBisSupp.jpg



(COMPORT SHAPED) DO NOT SWALLOW

Stimulant Laxative

Purpose

Consumer Health Care, owner of the registered trademark Dulcolax®. *This product is not manufactured or distributed by Boehringer Ingelheim

Questions or comments? 1-800-492-5988

Inactive ingredient hydrogenated vegetable oil

■ Lot No. & Exp. Date: see wrapper or box

Other information \blacksquare Store at room temperature. Do not exceed 30° C (86° F)

Drug Facts (continued)

	esu ton ob	children under 6 years of age
once daily	1/2 suppository	children 6 to under 12 years of age
nce daily	o ynotizoqqua f	adults and children 12 years of age and over
suppository well into the rectum, pointed end first Tretain about 15 to 20 minutes		

detach one suppository from the strip remove wrapper before inserting into rectum carefully insert one

Directions

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

If pregnant or breast-feeding, ask a health professional before use.

product. These may indicate a serious condition.

Stop use and ask a doctor if you have rectal bleeding or you fail to have a bowel movement after using this

When using this product it may cause abdominal discomfort, faintness, rectal burning and mild cramps

Ask a doctor before use if you have noticed a sudden change in bowel habits that lasts over a period of 2 weeks

■ if abdominal pain, nausea or vomiting are present ■ for longer than 1 week

Do not use unless directed by a doctor

For rectal use only.

Marnings

■ this product generally produces bowel movement in 15 minutes to 1 hour **USES** • for temporary relief of occasional constipation

Bisacodyl, USP 10 mg.....

Active ingredient (in each suppository)

Drug Facts

DO NOT USE IF WRAPPER IS TORN OR OPEN. INDIVIDUALLY SEALED FOR YOUR PROTECTION.

The product package shown above represents a sample of that currently in use. Additional packaging may also be available.

Bisacodyl Suppositories

Distributed by

Major Pharmaceuticals

Livonia, MI 48152

BISACODYL

bisacodyl suppository

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0573
Route of Administration	RECTAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	10 mg	

Inactive Ingredients	
Ingredient Name	Strength
HYDROGENATED PALM KERNEL OIL (UNII: FM8D1RE2VP)	

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:70000- 0573-2	4 in 1 CARTON	05/19/2021		
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:70000- 0573-1	1 in 1 CARTON	05/19/2021		
2		4 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	05/19/2021	

Labeler - Cardinal Health (063997360)

Registrant - DSC Laboratories, Inc. (097807374)

Establishment				
Name	Address	ID/FEI	Business Operations	
DSC Laboratories, Inc.		097807374	manufacture(70000-0573)	

Revised: 12/2024 Cardinal Health