GUMNUMB TOPICAL ANESTHETIC- benzocaine gel Crosstex International Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active Ingredient (in each gram)

Benzocaine 200mg

Purpose

Oral Anesthetic

Use

For oral mucosa use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy Alert: Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product Avoid contact with eyes. If contact occurs, flush with water.

Do not exceed recommended dosage. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breast feeding, ask a physician before use.

Keep out of reach of children.

Directions

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

Other Information

Store at 59° - 86° F (15° - 30° C).

Protect from freezing.

Inactive Ingredients

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

Questions or comments?

crosstex@crosstex.com or www.crosstex.com

PEEL BACK to review drug facts



REF UTS





UTS-BTL Rev. 7 - 7/18

Manufactured for: Crosstex International, Inc. 10 Ranick Road Hauppauge, NY 11788-4209



STRAVVBERRY 1 OZ. (30 G)

n 7 days; irritation, pain, or redness mouth symptoms do not improve persists or worsens; or if swelling, rash or fever develops, see your

in children under 2 years of age physician promptly. for teething

Avoid contact with eyes. If contact When using this product occurs, flush with water.

Do not exceed recommended dosage. If more than used for oain is accidentally swallowed get medical help or contact a

if pregnant or breast feeding, (eep out of reach of children. Poison Control Center right away ask a physician before use.

· Apply only amount needed to children under 2 years of age: the oral mucosa to prevent or Directions relieve pain.

do not use

Other information

Store at 59° - 86°F (15 - 30°C) Protect from freezing.

olue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, sodium saccharin. May contair **Inactive** ingredients lavoring, PEG 3350, PEG 400,

Ouestions or comments? crosstex@crosstex.com or additive.

tartrazine), yellow #6, as a color

99190006 Rev. 7 - 7/18

www.crosstex.com

Active ingredients in each gram) 3enzocaine 200 mg **Drug Facts**

Oral Anesthetic Purpose

For oral mucosal use only, as directed by dentist. For the

temporary relief of pain due to

minor dental procedures.

warning: Use of this product may Methemoglobinemia Warnings

treated promptly because it reduces the blood. This can occur even if you have used this product before. Stop use and seek immediate medical the amount of oxygen carried in attention if you or a child in your cause methemoglobinemia, a serious condition that must be

 pale, gray or blue colored skin care develops: (cyanosis)

 shortness of breath rapid heart rate headache

dizziness or lightheadedness

patients with a history of allergies Allergy Alert: Do not use on to local anesthetics such as fatigue or lack of energy

Do not use

procaine, butacaine, benzocaine or

other "caine" anesthetics

directed by a physician. If sore for more than 7 days unless

GUMNUMB TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:24794-225

DENTAL **Route of Administration**

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZ OCAINE** 200 mg in 1 g

Inactive Ingredients Ingredient Name Strength POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) SACCHARIN SODIUM (UNII: SB8ZUX40TY) FD&C YELLOW NO. 5 (UNII: I753WB2F1M)

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics				
Color	red	Score		
Shape		Size		
Flavor	STRAWBERRY	Imprint Code		
Contains				

F	Packaging				
#	tem Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:24794-225- 30	30 g in 1 JAR; Type 0: Not a Combination Product	07/01/2018		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph not final	part356	07/01/2013			
	part356	07/01/2013			

Labeler - Crosstex International Inc. (057728685)

Revised: 1/2022 Crosstex International Inc.