

PAIN RELIEF REGULAR STRENGTH- acetaminophen tablet
TopCo Associates LLC

TCR - 1001 - 2019-1016

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If

you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease.

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 tablets every 4 to 6 hours while symptoms last▪ do not take more than 10 tablets in 24 hours▪ do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">▪ take 1 tablet every 4 to 6 hours while symptoms last▪ do not take more than 5 tablets in 24 hours▪ do not use for more than 5 days unless directed by a doctor
children under 6 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients

povidone, pregelatinized starch, sodium starch glycolate, stearic acid

PRINCIPAL DISPLAY PANEL

NDC 36800-900-02

TopCare® Health™

Compare to Regular Strength Tylenol® Tablets Active Ingredient*

Regular Strength

Pain Relief

Acetaminophen 325 mg

Pain Reliever • Fever Reducer

Pharmacists Recommend

100 Tablets

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING

2 062721 00899C 0

83 RB2J2C21001011

DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING



This product is not manufactured or distributed by McKelvey Consumer Healthcare, distributor of Regular Strength Tylenol® Tablets.

Drug Facts

Active Ingredient (in each tablet) Acetaminophen 325 mg

Purpose Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ backache
■ minor pain of arthritis ■ toothache ■ muscular aches
■ menstrual and menstrual cramps
■ temporarily reduces fever

Warnings

User warnings: This product contains acetaminophen. Severe liver damage may occur if:
■ adult takes more than 4,000 mg of acetaminophen in 24 hours
■ child takes more than 5 doses in 24 hours, which is the maximum daily amount
■ taken with other drugs containing acetaminophen
■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if the user is allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if the user has liver disease.

Drug Facts (continued)

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days in adults
■ pain gets worse or lasts more than 5 days in children under 12 years
■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Directions

Do not take more than directed (see overdose warning)
adults and children 12 years and over
■ take 2 tablets every 4 to 6 hours while symptoms last
■ do not take more than 10 tablets in 24 hours
■ do not use for more than 10 days unless directed by a doctor
children 6-11 years
■ take 1 tablet every 4 to 6 hours while symptoms last
■ do not take more than 5 tablets in 24 hours
■ do not use for more than 5 days unless directed by a doctor
children under 6 years
■ ask a doctor

Drug Facts (continued)

Other information
■ store between 20-25°C (68-77°F)
■ retain carton for complete product information

Inactive ingredients povidone, polyethylene glycol, sodium starch glycolate, stearic acid



DISTRIBUTED BY
TOP CARE HEALTH LLC
EL PASO, TEXAS 75007
© TOPCO 44940521
QUESTIONS? 1-888-423-0139
topcarehealth.com
www.topcarehealth.com

Scan here for more information or call 1-888-423-0139



NDC 36800-900-02



COMPARE TO REGULAR STRENGTH TYLENOL® TABLETS ACTIVE INGREDIENT*

REGULAR STRENGTH

Pain Relief

ACETAMINOPHEN 325 mg

PAIN RELIEVER
FEVER REDUCER

100 TABLETS

actual size



REGULAR STRENGTH
Pain Relief

ACETAMINOPHEN 325 mg
PAIN RELIEVER • FEVER REDUCER

100 TABLETS

actual size

20-00-900-00899C:CN

CN

CN

CN

PAIN RELIEF REGULAR STRENGTH

acetaminophen tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:36800-900

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	M2A3;57344
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-900-02	1 in 1 CARTON	08/30/2011	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/30/2011	

Labeler - TopCo Associates LLC (006935977)

Revised: 12/2024

TopCo Associates LLC