

EQUALINE COMPLETE DUAL ACTION- famotidine, calcium carbonate and magnesium hydroxide tablet, chewable
Supervalu Inc

SuperValu Inc. Dual Action Complete Drug Facts

Active ingredients (in each chewable tablet)

Famotidine 10 mg

Calcium carbonate 800 mg

Magnesium hydroxide 165 mg

Purposes

Acid reducer

Antacid

Use

relieves heartburn associated with acid indigestion and sour stomach

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Antacids and acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- **do not swallow tablet whole: chew completely**
- to relieve symptoms, **chew** 1 tablet before swallowing
- do not use more than 2 chewable tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** calcium 330 mg; magnesium 80 mg
- Phenylketonurics: Contains Phenylalanine 2 mg per tablet
- read the directions and warnings before use
- read the bottle. It contains important information.
- store at 20-25°C (68-77°F)
- protect from moisture

Inactive ingredients

anhydrous lactose, artificial berry flavor, aspartame, D&C red no. 7 calcium lake, dextrates, FD&C blue no. 1 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyacrylate dispersion, polysorbate 80, povidone, pregelatinized starch, sodium starch glycolate, talc

Questions or comments?

1-855-423-2630

Principal Display Panel

compare to Pepcid Complete® active ingredients

dual action complete

famotidine 10mg, calcium carbonate 800mg, magnesium hydroxide 165mg tablets (chewable)

(acid reducer + antacid)

just one tablet relieves heartburn due to acid indigestion

SEE NEW WARNINGS

berry flavor

actual size
25 chewable tablets

EQUALINE®
dual action complete
famotidine 10mg, calcium carbonate 800mg, magnesium hydroxide 165mg tablets (chewable)
(acid reducer + antacid)
Just one tablet relieves heartburn due to acid indigestion
SEE NEW WARNINGS berry flavor
25 chewable tablets

Compare to Pepcid Complete* active ingredient*

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Warnings: Do not use if you are allergic to famotidine or other acid reducers. **Do not use** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ With other acid reducers

Use: relieves heartburn associated with acid indigestion and sour stomach

Warnings: Do not use if you are allergic to famotidine or other acid reducers. ■ If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ With other acid reducers

Ask a doctor before use if you have: ■ stomach or heart problems, ■ heartburn over 14 days, ■ history of stomach ulcers or bleeding, ■ kidney disease or diabetes, ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck, or shoulders, or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ frequent or ongoing nausea or vomiting ■ ulcers or other stomach problems

Ask doctor or pharmacist before use if you are taking: ■ any medicines that may interact with certain prescription drugs. ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

Stop use and ask a doctor if: ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

100% Quality GUARANTEED Use it or let us make it right. 855-472-2650

Divided into 500 mg (red) and 250 mg (white) tablets. Contains 250 mg of famotidine, 800 mg of calcium carbonate, and 165 mg of magnesium hydroxide. **US: In. No. 0801 JTS** **Div. Marketed by: JUNEI** **OL UTRM FREE** **Powdered, RI 02801 USA**

Directions or comments: 871-855-423-8530

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ Adults and children 12 years and over: ■ Take one swallow tablet twice daily with meals. ■ Do not use more than 2 swallow tablets in 24 hours

Other Information ■ Each tablet contains calcium 530 mg, magnesium 80 mg, and phenylephrine 2 mg. ■ Contains 2 mg of phenylephrine. ■ Read the directions and warnings before use. ■ Seal the package if broken. ■ Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30° (59°-86°) (77°F). ■ Protect from moisture

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ Adults and children 12 years and over: ■ Take one swallow tablet twice daily with meals. ■ Do not use more than 2 swallow tablets in 24 hours

Other Information ■ Each tablet contains calcium 530 mg, magnesium 80 mg, and phenylephrine 2 mg. ■ Contains 2 mg of phenylephrine. ■ Read the directions and warnings before use. ■ Seal the package if broken. ■ Store at 20°-25°C (68°-77°F); excursions permitted to 15°-30° (59°-86°) (77°F). ■ Protect from moisture

EQUALINE COMPLETE DUAL ACTION			
famotidine, calcium carbonate and magnesium hydroxide tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-321
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg	
CALCIUM CARBONATE (UNII: H0G9379FGK) (CARBONATE ION - UNII:7UJQ5OPE7D, CALCIUM CATION - UNII:2M83C4R6ZB)	CALCIUM CARBONATE	800 mg	
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	165 mg	
Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS LACTOSE (UNII: 3S Y5LH9 PMK)			
ASPARTAME (UNII: Z0H242BBR1)			
D&C RED NO. 7 (UNII: ECW0LZ41X8)			
DEXTRATES (UNII: G263MI44RU)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)			

MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	PINK (mottled)	Score	no score
Shape	ROUND (bi-layered)	Size	17mm
Flavor	BERRY	Imprint Code	L321
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-321-63	25 in 1 BOTTLE; Type 0: Not a Combination Product	09/02/2008	
2	NDC:41163-321-71	50 in 1 BOTTLE; Type 0: Not a Combination Product	09/02/2008	01/15/2017

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077355	09/02/2008	

Labeler - Supervalu Inc (006961411)