

**NOHIST LQ- chlorpheniramine maleate and phenylephrine hydrochloride liquid**  
**Larken Laboratories, Inc.**

-----  
**NoHist LQ**

**Drug Facts**

***Active Ingredients***

***(In each 5 mL teaspoonful)***

Chlorpheniramine Maleate, USP 4 mg

Phenylephrine HCl, USP 10 mg

***Purpose***

Chlorpheniramine Maleate Antihistamine

Phenylephrine HCl Nasal decongestant

***Uses***

Temporarily relieves these symptoms due to hay fever (allergic rhinitis):

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

***Warnings***

**Do not use**

- to sedate a child or make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this drug.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes

- glaucoma
- trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking any other nasal decongestant or stimulant
- taking sedatives or tranquilizers

**When using this product**

**Do not exceed recommended dosage.**

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur.
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of the reach of children**

In case of overdose, get medical help or contact a Poison Control Center immediately.

***Directions***

Do not exceed 6 doses in a 24-hour period

Age	Dose
Adults and children over 12 years of age	1 teaspoonsful (5 mL) every 4 hours
Children 6 to under 12 years of age	1/2 teaspoonsful (2.5 mL) every 4 hours
Children under 6 years of age	Ask your doctor

***Other Information***

- store at 20°-25°C (68°-77°F)
- very low sodium, contains 5 mg sodium per 5 mL teaspoonful

## **Inactive Ingredients**

bubblegum flavoring, citric acid, D&C Red #33, edetate disodium, glycerin, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, and sodium citrate dihydrate

## ***Questions or Comments***

Call 1-601-855-7678 weekdays from 9:00 am to 4:00 pm CST or go to <http://www.larkenlabs.com>.

## **Principal Display Panel**

**Figure 1:** 16 oz. Bottle Label

NDC 68047-185-16

# NOHIST·LQ

**ANTIHISTAMINE  
NASAL DECONGESTANT**

**SUGAR FREE / ALCOHOL FREE**

**Bubblegum Flavored Liquid**

**DO NOT USE IF FOIL SEAL UNDER  
THE CAP IS BROKEN OR MISSING.**

Distributed by:

**LARKEN  
LABORATORIES**

Canton, MS 39046

16 fl. oz. (473 mL)



Lot/Exp. date:

400734-06 Rev. 07/2016

## Drug Facts

**Active ingredients** *(in each 5 mL teaspoonful)* **Purpose**  
Chlorpheniramine Maleate, USP 4 mg ..... Antihistamine  
Phenylephrine HCl, USP 10 mg ..... Nasal decongestant

**Uses**  
temporarily relieves these symptoms due to hay fever (allergic rhinitis): ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ temporarily restores freer breathing through the nose

**Warnings**  
**Do not use**  
■ to sedate a child or make a child sleepy  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's Disease), or for two weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this drug.

**Ask a doctor before use if you have**  
■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**  
■ taking any other nasal decongestant or stimulant  
■ taking sedatives or tranquilizers

**When using this product**  
**Do not exceed recommended dosage.**  
■ marked drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children.

**Stop use and ask a doctor if**  
■ nervousness, dizziness, or sleeplessness occur.  
■ if symptom do not improve within 7 days or are accompanied by fever.

**PEEL**

## NOHIST LQ

chlorpheniramine maleate and phenylephrine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68047-185
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE -	PHENYLEPHRINE	10 mg

UNII:1WS297W6MV)

HYDROCHLORIDE

in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0KOOR)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68047-185-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/06/2011	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/06/2011	

**Labeler** - Larken Laboratories, Inc. (149484540)**Registrant** - Larken Laboratories, Inc. (149484540)

Revised: 10/2023

Larken Laboratories, Inc.