

MILK OF MAGNESIA- magnesium hydroxide suspension

Major Pharmaceuticals

Milk of Magnesia

Drug Facts

Active ingredient (in each 30 mL dose)

Magnesium hydroxide 2400 mg

Purpose

Antacid / Saline Laxative

Uses as Antacid

- relieves
 - heartburn
 - sour stomach
 - acid indigestion
 - and upset stomach associated with these symptoms

Uses as Saline Laxative

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 1/2 to 6 hours

Warnings for Antacid

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use If you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product

- do not take more than 60 mL (2 doses) in a 24-hour period
- do not use the maximum dosage for more than 2 weeks except under the advice and supervision of a doctor
- may cause a laxative effect

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.

Warnings for Saline Laxative

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use If you are

- presently taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions for Antacid

- shake well before use
- mL=milliliters
- adults and children 12 years of age or older: 30 mL (1 dose) up to 2 times a day or as directed by a doctor
- children under 12 years of age: ask a doctor

Directions for Saline Laxative

- do not exceed the maximum recommended daily dose of 60 mL in a 24 hour period
- shake well before use
- mL=milliliters
- drink a full glass (8 oz) of liquid with each dose
- adults and children 12 years of age or older: 30 mL to 60 mL (1 to 2 doses)
- children under 12 years of age: ask a doctor
- the dose may be taken as a single daily dose or in divided doses

Other information for Antacid

- each 30 mL contains: **magnesium 1 g**

- store at 20° - 25°C (68° - 77°F)
- protect from freezing
- **tamper-evident: do not use if seal on cup is missing or torn**

Other information for Saline Laxative

- each 30 mL contains: **magnesium 1 g**
- store at 20° - 25°C (68° - 77°F)
- protect from freezing
- **tamper-evident: do not use if seal on cup is missing or torn**

Inactive ingredients antifoam af emulsion, flavoring, hydroxypropyl methylcellulose, methylparaben, potassium citrate, propylene glycol, propylparaben, purified water, sodium saccharin, sorbitol

Questions or comments? call 1-800-845-8210

R03/25

Distributed by:
MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268

PRINCIPAL DISPLAY PANEL - 30 mL Cup Tray Label

MAJOR®
NDC 0904-7564-62

Milk of Magnesia USP
Magnesium Hydroxide 2400 mg
Antacid / Saline Laxative

2400 mg/ 30 mL

SHAKE WELL

Delivers 30 mL
See Drugs Facts Insert
For Institutional Use Only

MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268

F0431C300325

MAJOR®

NDC 0904-7564-62

Milk of Magnesia USP

Magnesium Hydroxide 2400 mg
Antacid / Saline Laxative

2400 mg/30 mL

SHAKE WELL



Delivers 30 mL

**See Drugs Facts Insert
For Institutional Use Only**

**MAJOR® PHARMACEUTICALS
Indianapolis, IN 46268**

F0431C300325

MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDc:0904-7564
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	2400 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CYCLOMETHICONE 4 (UNII: CZ227117JE)	
HYPROMELLOSE 2910 (4000 MPA.S) (UNII: RN3152OP35)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	white	Score
Shape		Size
Flavor	SPEARMINT	Imprint Code
Contains		

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7564-73	10 in 1 CASE	01/08/2026	
1		10 in 1 TRAY		
1	NDC:0904-7564-62	30 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	01/08/2026	

Labeler - Major Pharmaceuticals (191427277)

Revised: 1/2026

Major Pharmaceuticals