

BREACOLD CHILDREN- guaifenesin, dextromethorphan hbr liquid
Menper Distributors, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Active ingredients (in each 15mL tsp)

Guaifenesin 100 mg.....Expectorant
Dextromethorphan HBr 10 mg.....Cough suppressant

purpose

Expectorant
Cough Suppressant

Uses

Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold and help loosen phlegm, (mucus) and thin bronchial secretions to make coughs more productive.

Warnings Do not exceed recommended dosage.

- A persistent cough may be a sign of a serious condition. If cough persists for more than one week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

Do not use:

- This product for persistent or chronic cough, such as occurs with asthma, or if cough is accompanied by excessive phlegm (mucus), unless directed by a doctor.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Keep out of reach of children.

If you are pregnant or breast-feeding,

ask a health professional before use. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed 6 doses in 24 hours.

Age	Dose
Children 6 to under 12 years of age.....	1 teaspoonful (5 mL) every 4 hours
Children 2 to under 6 years of age.....	1/2 teaspoonful (2.5 mL) every 4 hours
Children under 2 years of age.....	consult your doctor

Other information

Store this product between 15°-30°C (59°-86°F)

Inactive ingredients

Citric acid, D&C red #33, glycerin, methylparaben, propylene glycol, propylparaben ,
raspberry flavor, sucrose, water

Questions?

Call 1-800-560-5223 M-F 9 AM to 4 PM Eastern.

BREA - COLD

Alivia:
• La Tos
• La Congestión de Pecho/Mucosa



Niños

Sabor a Frambuesa

DEXTROMETHORPHAN HBR
(Supresor de la Tos)

GUAIFENESINA
(Expectorante)

Contenido: 4 Fl. Oz (120 mL)

Factores Farmacológicos

Ingredientes activos (en cada 15 mL) Propósitos
Dextrometorphan Hbr 10 mg..... Alivio de la Tos
Guaifenesina 100 mg..... Expectorante

Usos

Alivio temporal de la tos debido a irritación de la garganta y bronquios, asociada al resfriado común o gripe y ayuda a aliviar la fiebre (mucosa) y secreciones bronquiales para hacer la tos más productiva.

Advertencias

No exceder la dosis recomendada.

Una tos persistente puede ser señal de una condición seria. Si la tos persiste por más de una semana, regresa o está acompañada por fiebre, dolor de cabeza o erupciones en la piel, consulte a su médico.

No usar

este producto para tos persistente o crónica, tal como ocurre si tiene asma, o si la tos es acompañada por flema excesiva (secreción), sólo por indicación médica.

Si usted está tomando ahora alguna medicina bajo prescripción que contenga inhibidores de la monoaminoxidasa (MAOI) (ciertas drogas para la depresión, esquizofrenia, o condiciones emocionales, o enfermedad de Parkinson) o pasadas 2 semanas después de haber ingerido medicamentos que contengan MAOI. Si usted no está seguro que el medicamento que usted está tomando contiene MAOI, consulte a su doctor o a un farmacéutico antes de usarlo.

Mantener fuera del alcance de los niños

Si está embarazada o lactando, pregunte a un profesional de la salud antes de usarlo. En caso de sobredosis accidental, busque ayuda médica o contacte a un Centro de Control de Envenenamiento inmediatamente.

Instrucciones

No exceda más de 6 dosis en 24 horas.

edad	dosis
Niños de 6 a menores de 12 años de edad	1 cucharada (15 mL) cada 4 horas
Niños de 2 a menores de 6 años de edad	1/2 cucharada (7.5 mL) cada 4 horas
Niños menores de 2 años de edad	consulte a su médico

Otra información

conservar este producto entre 15° a 30°C (59° a 86°F)

Ingredientes inactivos ácido cítrico, D&C rojo #33, glicerina, metilparabeno, propilenglicol, propilparabeno, sabor a frambuesa, sucrosa, agua

Preguntas? llame 1-800-560-5223 L-V 9 AM a 4 PM Este.

EVIDENCIA DE SEGURIDAD: NO USAR ESTE PRODUCTO SI EL SELLO IMPRESO SOBRE LA TAPA DE LA BOTELLA ESTA TORCIDO, ROTO O FALTA.

Distributed by/Distribuido por:
Menper Distributors, Inc.
11421 N.W. 107 Street, Suite 24, Miami, FL 33178
www.menperdistributors.com

BREA - COLD

Relieves:
• Cough
• Chest Congestion/Mucus



Children

Raspberry Flavor

DEXTROMETHORPHAN HBR
(Cough Suppressant)

GUAIFENESIN
(Expectorant)

Contents: 4 Fl. Oz (120 mL)

Drug Facts

Active ingredients (in each 15 mL) Purposes
Dextromethorphan Hbr 10 mg..... Cough suppressant
Guaifenesin 100 mg..... Expectorant

Uses

Temporarily relieves cough due to minor throat and bronchial irritation as may occur with the common cold and helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive.

Warnings

Do not exceed recommended dosage.

A persistent cough may be a sign of a serious condition. If cough persists for more than one week, tends to recur, or is accompanied by fever, rash or persistent headache, consult a doctor.

Do not use

this product for persistent or chronic cough, such as occurs with asthma, or if cough is accompanied by excessive phlegm (mucus), unless directed by a doctor.

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Keep out of reach of children

If pregnant or breast-feeding, ask a health professional before use. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed 6 doses in 24 hours.

age	dose
Children 6 to under 12 years of age	1 tablespoonful (15 mL) every 4 hours
Children 2 to under 6 years of age	1/2 tablespoonful (7.5 mL) every 4 hours
Children under 2 years of age	consult your doctor

Other information

store this product between 15° to 30°C (59° to 86°F)

Inactive ingredients citric acid, D&C red #33, glycerin, methylparaben, propylene glycol, propylparaben, raspberry flavor, sucrose, water

Questions? call 1-800-560-5223 M-F 9 AM to 4 PM Eastern.

TAMPER-EVIDENT: DO NOT USE THIS PRODUCT IF PRINTED CELLO-BAND OVER CAP IS TORN, BROKEN OR MISSING.



BREACOLD CHILDREN

guaifenesin, dextromethorphan hbr liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53145-007
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3R0TS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

CITRIC ACID ACETATE (UNII: DSO12WL7AU)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
RASPBERRY (UNII: 4N14V5R27W)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53145-007-04	1 in 1 CARTON	02/21/2012	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/21/2012	

Labeler - Menper Distributors, Inc. (101947166)

Revised: 2/2022

Menper Distributors, Inc.