

**IBUPROFEN MINIS- ibuprofen capsule, liquid filled  
EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### **Other information**

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use
- store at 20°-25°C (68°-77°F)
- avoid excessive heat above 40°C (104°F). Protect from light
- swallow whole; do not crush, chew, or dissolve

### **Inactive ingredients**

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol, solution, white ink

### **Questions or comments?**

Call **1-888-287-1915**

### **Principal Display Panel**

Compare to Advil® Liqui-Gels® Minis active ingredient†

**Ibuprofen** Capsules, 200 mg

**Pain Reliever/Fever Reducer (NSAID)**

MINIS

- Smaller capsule
- Same strength

LIQUID-FILLED CAPSULES

**TAMPER-EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

†This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Advil® Liqui-Gels® Minis.

DISTRIBUTED BY: Walmart Inc.,  
Bentonville, AR 72716

**Product Label**

**equate™**  
**Ibuprofen**  
Capsules, 200mg

**Pain Reliever/  
Fever Reducer  
(NSAID)**

**MINIS**  
**200 mg**  
**300 LIQUID-FILLED CAPSULES**

Actual Size

Compare to Advil® Liqui-Gels® Minis active ingredient

NDC 49035-265-30

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**Drug Facts**

**Active ingredient (in each capsule)**  
Sulbuzited ibuprofen equal to 200 mg ibuprofen (NSAID)\*.....Pain reliever/fever reducer (present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves minor aches and pains due to:  
muscular aches ■ minor pain of arthritis ■ toothache  
backache ■ the common cold ■ menstrual cramps  
temporarily reduces fever

**Warnings**  
**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
hives ■ facial swelling ■ asthma (wheezing)  
shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Drug Facts (continued under label)**

Satisfaction guaranteed – Or we'll replace it or give you your money back. For questions or comments please call 1-888-287-1915.

483524

8113135431 8

Lot No.: Exp. Date:

**DISTRIBUTED BY: Walmart Inc.,**  
Bentonville, AR 72716  
PRODUCT OF CHINA  
\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare Holdings (US) LLC, distributor of Advil® Liqui-Gels® Minis.  
PDU-C560C 1B008193

**PEEL HERE**

**Drug Facts (continued)**

**Ask a doctor before use if**  
the stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product,** take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if**  
you experience any of the following signs of stomach bleeding:  
feel faint ■ vomit blood ■ have bloody or black stools  
have stomach pain that does not get better ■ you have symptoms of heart problems or stroke:  
chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days  
redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

**Other information**  
each capsule contains: potassium 20 mg  
read all warnings and directions before use ■ store at 20°-25°C (68°-77°F) ■ avoid excessive heat above 40°C (104°F). Protect from light. ■ swallow whole; do not crush, chew, or dissolve

**Inactive ingredients**  
FD&C green #9, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution, white ink

**Questions or comments?**  
1-888-287-1915

**EQUATE Ibuprofen 200 mg Softgels Minis**

**IBUPROFEN MINIS**

ibuprofen capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-265	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg	
<b>Inactive Ingredients</b>				
	<b>Ingredient Name</b>		<b>Strength</b>	
	FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)			
	GELATIN (UNII: 2G86QN327L)			
	POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
	POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)			
	WATER (UNII: 059QF0KO0R)			
	SORBITOL (UNII: 506T60A25R)			
	SORBITAN (UNII: 6O92ICV9RU)			
	MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)			
<b>Product Characteristics</b>				
<b>Color</b>	green	<b>Score</b>	no score	
<b>Shape</b>	CAPSULE	<b>Size</b>	12mm	
<b>Flavor</b>		<b>Imprint Code</b>	P33	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:49035-265-16	160 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/31/2019	
2	NDC:49035-265-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/31/2019	
3	NDC:49035-265-80	80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/31/2019	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
ANDA	ANDA206568	12/31/2019		

**Labeler** - EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC) (051957769)