

**MOTION SICKNESS RELIEF- dimenhydrinate tablet**  
**Amerisource Bergen**

-----  
**Good Neighbor Pharmacy 44-198**

***Active ingredient (in each tablet)***

Dimenhydrinate 50 mg

***Purpose***

Antiemetic

***Uses***

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

***Warnings***

**Do not use**

for children under 2 years of age unless directed by a doctor.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- marked drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

**Other information**

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from moisture
- see end flap for expiration date and lot number

**Inactive ingredients**

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

**Questions or comments?**

**1-800-426-9391**

**Principal display panel**

**GOOD  
NEIGHBOR  
PHARMACY®**

**NDC 46122-536-53**

Compare to Dramamine® Original  
Formula active ingredient\*

**Motion Sickness Relief**

dimenhydrinate 50 mg / antiemetic

*Original Formula*

Prevents Nausea, Vomiting & Dizziness  
Children & Adults

**12** Tablets | 50 mg Each

Actual Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula.

50844 ORG051819802

Distributed By  
AmerisourceBergen  
1 West First Avenue  
Conshohocken, PA 19428  
Questions or Concerns?  
[www.mygnp.com](http://www.mygnp.com)




Compare to Dramamine® Original Formula active ingredient\*

# Motion Sickness Relief

dimenhydrinate 50 mg / antiemetic

**Original Formula**

Prevents Nausea, Vomiting & Dizziness For Children & Adults



12 Tablets | 50 mg Each

Actual Size



Compare to Dramamine® Original Formula active ingredient\*


NDC 46122-536-53

# Motion Sickness Relief

dimenhydrinate 50 mg / antiemetic

**Original Formula**

Prevents Nausea, Vomiting & Dizziness For Children & Adults



12 Tablets | 50 mg Each

Actual Size



7-Fold

#7732RZ

B-0045-198-02-R  
OR0051819802

Distributed By  
AmenSourceBergen  
1 West First Avenue  
Conshohocken, PA 19428  
www.mygnp.com  
Questions or Concerns?



\*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine® Original Formula.  
50844 ORG051819802

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

no print / no varnish area  
lot no. & exp. date

8:30 AM-4:00 PM ET, Monday-Friday  
**Questions or comments?** 1-800-426-9391

**Inactive ingredients**  
 croscarmellose sodium, dibasic calcium phosphate hydrate, magnesium stearate, microcystalline cellulose, silicon dioxide, stearic acid

**Other information**  
 Each tablet contains calcium 35 mg  
 TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGING IS OPENED OR BLISTER IS TORN OR BROKEN  
 marked drowsiness may occur  
 alcohol, sedatives, and tranquilizers may increase drowsiness  
 use caution when driving a motor vehicle or operating machinery  
 avoid alcoholic beverages  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Uses** for prevention and treatment of these symptoms associated with motion sickness:  
 nausea ■ vomiting ■ dizziness

**Warnings**  
 Do not use for children under 2 years of age unless directed by a doctor.  
 Ask a doctor before use if you have  
 a breathing problem such as emphysema or chronic bronchitis ■ glaucoma  
 difficulty in urination due to enlargement of the prostate gland  
 Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.  
 When using this product  
 marked drowsiness may occur  
 alcohol, sedatives, and tranquilizers may increase drowsiness  
 use caution when driving a motor vehicle or operating machinery  
 avoid alcoholic beverages  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.


**adults and children**  
 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours; or as directed by a doctor

**children 6 to under 12 years**  
 ½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours; or as directed by a doctor


**children 2 to under 6 years**  
 ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours; or as directed by a doctor

see end flap for expiration date and lot number  
 protect from moisture  
 store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 protect from moisture

ABC#: 10294647 REV: 08/24



3 46122 53653 0



44-198

## MOTION SICKNESS RELIEF

dimenhydrinate tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46122-536
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIMENHYDRINATE</b> (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	44;198
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-536-53	2 in 1 CARTON	12/01/1992	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	12/01/1992	

**Labeler** - Amerisource Bergen (007914906)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(46122-536)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(46122-536) , pack(46122-536)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(46122-536)

Revised: 9/2025

Amerisource Bergen