

# **KARBINAL ER- carbinoxamine maleate suspension, extended release**

## **Aytu Therapeutics, LLC**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use KARBINAL<sup>®</sup> ER safely and effectively. See full prescribing information for KARBINAL<sup>®</sup> ER.**

### **Karbinal<sup>®</sup> ER (carbinoxamine maleate) extended-release oral suspension**

**Initial U.S. Approval: 1953**

### **-----RECENT MAJOR CHANGES-----**

Contraindications, Nursing Mothers (4)

Removed 3/2021

### **-----INDICATIONS AND USAGE-----**

Karbinal ER is indicated for adults and pediatric patients 2 years of age and older for the symptomatic treatment of:

- Seasonal and perennial allergic rhinitis ( 1)
- Vasomotor rhinitis ( 1)
- Allergic conjunctivitis due to inhalant allergens and foods ( 1)
- Mild, uncomplicated allergic skin manifestations of urticaria and angioedema ( 1)
- Dermatographism ( 1)
- As therapy for anaphylactic reactions *adjunctive* to epinephrine and other standard measures after the acute manifestations have been controlled ( 1)
- Amelioration of the severity of allergic reactions to blood or plasma ( 1)

### **-----DOSAGE AND ADMINISTRATION-----**

Adults and Adolescents 12 years of age and older ( 2.3):

7.5 mL to 20 mL (6 to 16 mg) every 12 hours

Pediatric patients 2-11 years of age (approximately 0.2 to 0.4 mg/kg/day) ( 2.4):

2 to 3 years – 3.75 mL to 5 mL (3 to 4 mg) every 12 hours

4 to 5 years – 3.75 mL to 10 mL (3 to 8 mg) every 12 hours

6 to 11 years – 7.5 mL to 15 mL (6 to 12 mg) every 12 hours

### **-----DOSAGE FORMS AND STRENGTHS-----**

Extended-release oral suspension: 4 mg carbinoxamine maleate per 5 mL ( 3)

### **-----CONTRAINDICATIONS-----**

- Children younger than 2 years of age ( 4)
- Patients with known hypersensitivity to the drug or any of the inactive ingredients ( 4)
- Monoamine oxidase inhibitors (MAOI) ( 4)

### **-----WARNINGS AND PRECAUTIONS-----**

- Activities requiring mental alertness: Avoid engaging in hazardous tasks requiring complete mental alertness such as driving or operating machinery. ( 5.2)
- Anticholinergic actions: Use with caution in patients with increased intraocular pressure, narrow angle glaucoma, hyperthyroidism, cardiovascular disease, hypertension, stenosing peptic ulcer, symptomatic prostatic hypertrophy, bladder neck obstruction, pyloroduodenal obstruction. ( 5.3)
- Contains sodium metabisulfite, a sulfite that may cause anaphylaxis including life-threatening or less severe asthmatic episodes in susceptible individuals. ( 5.4)

### **-----ADVERSE REACTIONS-----**

Most common adverse reactions are: sedation, sleepiness, dizziness, disturbed coordination, epigastric distress, and thickening of bronchial secretions. ( 6)

**To report SUSPECTED ADVERSE REACTIONS, contact Cerecor, Inc., at 1-866-416-9637 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

---

## DRUG INTERACTIONS

---

- Monoamine oxidase inhibitors (MAOIs): Prolong and intensify the anticholinergic (drying) effects. ( 4 and 7)
- Alcohol and CNS depressants (hypnotics, sedatives, tranquilizers, etc.): Avoid concomitant use due to additive adverse effects. ( 7)

## USE IN SPECIFIC POPULATIONS

---

- Lactation: Advise not to breastfeed. ( 8.2)
- Contraindicated in children younger than 2 years of age. ( 4 and 8.4)
- May cause sedation or excitation in young children. ( 8.4)
- May cause dizziness, sedation, and hypotension in elderly patients. Start elderly patients on lower doses and observe closely for confusion and over-sedation. ( 8.5)

Revised: 8/2021

---

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Overview

#### 2.2 Administration

#### 2.3 Recommended Dosage for Adults and Adolescents 12 years of age and older:

#### 2.4 Recommended Dosage for Pediatric Patients 2 to 11 years of age (approximately 0.2 to 0.4 mg/kg/day):

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Pediatric Mortality

#### 5.2 Somnolence and Impaired Mental Alertness

#### 5.3 Concomitant Medical Conditions

#### 5.4 Allergic Reactions due to Sulfites, including Anaphylaxis

### 6 ADVERSE REACTIONS

### 7 DRUG INTERACTIONS

### 8 USE IN SPECIFIC POPULATIONS

#### 8.1 Pregnancy

#### 8.2 Lactation

#### 8.4 Pediatric Use

#### 8.5 Geriatric Use

### 10. OVERDOSAGE

### 11. DESCRIPTION

### 12. CLINICAL PHARMACOLOGY

#### 12.1 Mechanism of Action

#### 12.3 Pharmacokinetics

### 13. NONCLINICAL TOXICOLOGY

#### 13.1. Carcinogenesis, Mutagenesis, Impairment of Fertility

### 14. CLINICAL STUDIES

### 16. HOW SUPPLIED/STORAGE AND HANDLING

#### Administration

#### Activities Requiring Mental Alertness

MAOIs  
Lactation

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Karbinal ER is indicated for adults and pediatric patients 2 years of age and older for the symptomatic treatment of:

- Seasonal and perennial allergic rhinitis
- Vasomotor rhinitis
- Allergic conjunctivitis due to inhalant allergens and foods
- Mild, uncomplicated allergic skin manifestations of urticaria and angioedema
- Dermatographism
- As therapy for anaphylactic reactions *adjunctive* to epinephrine and other standard measures after the acute manifestations have been controlled
- Amelioration of the severity of allergic reactions to blood or plasma

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Overview**

The dosage of Karbinal ER should be individualized based on the severity of the condition and the response of the patient. Start with lower doses and increase as needed and tolerated.

#### **2.2 Administration**

Administer Karbinal ER by the oral route only. Measure Karbinal ER with an accurate milliliter measuring device. A household teaspoon is not an accurate measuring device and could lead to overdosage. A pharmacist can provide an appropriate measuring device and can provide instructions for measuring correct dose.

#### **2.3 Recommended Dosage for Adults and Adolescents 12 years of age and older:**

7.5 mL to 20 mL (6 to 16 mg) every 12 hours administered orally

#### **2.4 Recommended Dosage for Pediatric Patients 2 to 11 years of age (approximately 0.2 to 0.4 mg/kg/day):**

2 to 3 years: 3.75 mL to 5 mL (3 to 4 mg) every 12 hours administered orally

4 to 5 years: 3.75 mL to 10 mL (3 to 8 mg) every 12 hours administered orally

6 to 11 years: 7.5 mL to 15 mL (6 to 12 mg) every 12 hours administered orally

### **3 DOSAGE FORMS AND STRENGTHS**

Extended-release oral suspension: 4 mg carbinoxamine maleate per 5 mL

### **4 CONTRAINDICATIONS**

Karbinal ER is contraindicated in:

- children younger than 2 years of age because deaths have been reported in this age group (*see Warnings and Precautions (5.1)*)
  - patients who are hypersensitive to carbinoxamine maleate or any of the inactive ingredients in Karbinal ER [*see Warnings and Precautions (5.1)*]
- patients who are taking monoamine oxidase inhibitors (MAOI) [*see Drug Interactions(7)*].

### **5 WARNINGS AND PRECAUTIONS**

#### **5.1 Pediatric Mortality**

Deaths have been reported in children less than 2 years of age who were taking carbinoxamine-containing drug products; therefore, Karbinal ER is contraindicated in children younger than 2 years of age.

#### **5.2 Somnolence and Impaired Mental Alertness**

Karbinal ER may produce marked drowsiness and impair the mental or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery. Advise patients to avoid engaging in hazardous tasks requiring mental alertness and motor coordination after ingestion of Karbinal ER. Avoid concurrent use of Karbinal ER with alcohol or other central nervous system depressants because additional impairment of central nervous system performance may occur.

#### **5.3 Concomitant Medical Conditions**

Karbinal ER has anticholinergic (atropine-like) properties and, therefore, should be used with caution in patients with: increased intraocular pressure, narrow angle glaucoma, hyperthyroidism, cardiovascular disease, hypertension, stenosing peptic ulcer, symptomatic prostatic hypertrophy, bladder neck obstruction, or pyloroduodenal obstruction.

#### **5.4 Allergic Reactions due to Sulfites, including Anaphylaxis**

Karbinal ER contains sodium metabisulfite, a sulfite that may cause allergic-type reactions, including anaphylaxis and life-threatening or less severe asthmatic episodes, in susceptible individuals. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic individuals.

### **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are described elsewhere in the

labeling:

- Somnolence and Impaired Mental Alertness [see *Warnings and Precautions (5.2)*].
- Allergic Reactions due to Sulfites, including Anaphylaxis [see *Warnings and Precautions (5.2)*].

The most frequent adverse reactions include: sedation, sleepiness, dizziness, disturbed coordination, epigastric distress, and thickening of bronchial secretions. In clinical use, younger children and older adults may be particularly sensitive to adverse reactions [see *Pediatric Use (8.4)* and *Geriatric Use (8.5)*].

The following adverse reactions, listed by body system, have been identified in case reports and during the use of carbinoxamine in observational studies. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: Urticaria, drug rash, anaphylactic shock, photosensitivity, excessive perspiration, chills, dryness of mouth, nose and throat.

Cardiovascular: Hypotension, headache, palpitations, tachycardia, extrasystoles.

Central Nervous System: Fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, paresthesia, blurred vision, diplopia, vertigo, tinnitus, acute labyrinthitis, hysteria, neuritis, convulsions.

Gastrointestinal: Anorexia, nausea, vomiting, diarrhea, constipation.

Hematologic: Hemolytic anemia, thrombocytopenia, agranulocytosis.

Laboratory: Increase in uric acid levels.

Respiratory: Tightness of chest and wheezing, nasal stuffiness.

Urogenital: Urinary frequency, difficult urination, urinary retention, early menses.

## **7 DRUG INTERACTIONS**

- Do not use of Karbinal ER in patients who are taking monoamine oxidase inhibitors (MAOIs), which prolong and intensify the anticholinergic (drying) effects of antihistamines.
- Avoid use of Karbinal ER with alcohol and other CNS depressants (hypnotics, sedatives, tranquilizers, etc.) due to additive effects.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

Published data over decades of use of antihistamines, including carbinoxamine, have not identified a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. However, published data specifically evaluating the risk of carbinoxamine were not found. Animal reproductive studies have not been conducted with carbinoxamine maleate.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risks of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## **8.2 Lactation**

### Risk Summary

Based on the physical properties of carbinoxamine, it is likely that carbinoxamine is present in breastmilk. There are published reports of drowsiness and irritability in infants exposed to antihistamines via breastmilk. There are post-marketing reports of deaths in children under 2 years of age exposed to carbinoxamine by oral administration. There are no available data on the effects on milk production. It is not recommended to breastfeed during treatment with Karbinal ER [see *Warnings and Precautions (5.1)*] and *use in Specific Populations (8.4)*].

## **8.4 Pediatric Use**

Karbinal ER is contraindicated in pediatric patients younger than 2 years of age because deaths have been reported in this patient population who were taking carbinoxamine-containing drug products.

The safety and effectiveness of Karbinal ER in pediatric patients aged 2 years and older have been established and based on demonstration of bioequivalence to the immediate-release reference product [see *Clinical Pharmacology (12.3)*]. Carbinoxamine may diminish mental alertness or produce sedation in children. Paradoxical reactions with excitation are more likely in younger children.

## **8.5 Geriatric Use**

Karbinal ER may cause dizziness, hypotension, confusion, or over-sedation in the elderly. Start elderly patients on lower doses and observe closely.

## **10. OVERDOSAGE**

Overdosage with carbinoxamine may cause central nervous system depression or stimulation, hallucinations, convulsions, and death. Atropine-like signs and symptoms – dry mouth; fixed, dilated pupils; flushing; and gastrointestinal symptoms may also occur.


The treatment of overdosage consists of discontinuation of Karbinal ER and institution of symptomatic and supportive therapy. Vital signs (including respiration, pulse, blood pressure, and temperature) and EKG should be monitored. Induction of vomiting is not recommended. Activated charcoal should be given and gastric lavage should be considered after ingestion of a potentially life-threatening amount of drug. In the presence of severe anticholinergic effects, physostigmine may be useful. Vasopressors may be used to treat hypotension.

## **11. DESCRIPTION**


Each 5 mL of Karbinal ER Extended-release Oral Suspension contains carbinoxamine

complexed with polistirex equivalent to 4 mg carbinoxamine maleate and the following inactive ingredients: citric acid anhydrous, strawberry-banana flavor, glycerin, high fructose corn syrup, methylparaben, modified food starch, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, triacetin, and xanthan gum.

Carbinoxamine maleate is freely soluble in water. The chemical name is 2-[(4-chlorophenyl)-2-pyridinylmethoxy]- *N,N*-dimethylethanamine (Z)-2-butenedioate (1:1), which has the following structure:



The drug-polistirex complex is formed with the active ingredient (carbinoxamine maleate, USP) and sodium polystyrene sulfonate, USP, which has the following structure:



## 12. CLINICAL PHARMACOLOGY

## 12.1 Mechanism of Action

Carbinoxamine is an H<sub>1</sub> receptor antagonist (antihistamine) that exhibits anticholinergic (drying) and sedative properties.

Antihistamines compete with histamine for receptor sites on effector cells.

## 12.3 Pharmacokinetics

Karbinal ER after single-dose administration of 16 mg was bioequivalent to the reference carbinoxamine immediate-release oral solution after the administration of two doses of 8 mg six hours apart under fasting conditions. The carbinoxamine mean (SD) peak plasma concentration (C<sub>max</sub>) was 28.7 (5.3) ng/mL at 6.7 hours after Karbinal ER administration. The plasma half-life of carbinoxamine was 17.0 hours. There was no effect of food on the pharmacokinetic parameters.

Karbinal ER after multiple-dose administration of 16 mg every 12 hours for 8 days was bioequivalent to the reference carbinoxamine immediate-release oral solution after multiple-dose administration of 8 mg every 6 hours. The mean (SD) steady-state C<sub>max</sub> was 72.9 (24.4) ng/mL at 5.6 hours after Karbinal ER administration. Carbinoxamine mean (SD) minimum plasma concentration at steady-state was 51.8 (20.3) ng/mL.

## 13. NONCLINICAL TOXICOLOGY

### 13.1. Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term studies in animals have been performed to determine the possible effects of carbinoxamine on carcinogenesis, mutagenesis, and fertility.

## 14. CLINICAL STUDIES

The effectiveness and safety of Karbinal ER is based on demonstration of bioequivalence to the immediate-release reference product [see *Pharmacokinetics (12.3)*].

## 16. HOW SUPPLIED/STORAGE AND HANDLING

Karbinal ER Oral Suspension contains 4 mg carbinoxamine maleate per 5 mL. It is a light beige to tan viscous suspension with strawberry-banana flavor and is supplied as follows:

NDC 23594-101-05 Bottles of 16 fl oz (480 mL)

NDC 23594-101-01 Bottles of 1 fl oz (30 mL) Physician Samples

Store at 25°C (77°F); excursions permitted from 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.]

Dispense in tight, light-resistant container with child-resistant closure.

## 17. Patient Counseling Information

## **Administration**

Advise patients to measure Karbinal ER with an accurate milliliter measuring device. A household teaspoon is not an accurate measuring device and could lead to overdosage. *[see Dosage and Administration (2.2)].*

## **Activities Requiring Mental Alertness**

Advise patients to use caution when driving a motor vehicle or operating machinery. Karbinal ER may produce marked drowsiness and impair the mental or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery *[see Warnings and Precautions (5.2)].*

## **MAOIs**

Advise patients to not use MAOIs while taking Karbinal ER. MAOIs may prolong and intensify the anticholinergic (drying) effects *[see Contraindications (4.4) and Drug Interactions (7)].*

## **Lactation**

Advise women that breastfeeding is not recommended during treatment with Karbinal ER *[see Warnings and Precautions (5.1) and Use in Specific Populations (8.2 and 8.4)].*

## **PRINCIPAL DISPLAY PANEL - 480 mL Bottle Label**

**NDC 23594-101-05**

***Karbinal*<sup>™</sup> ER**

**(carbinoxamine maleate)**

**Extended-release**

**Oral Suspension**

**4 mg/5 ml**

Shake Well Before Use

Dose every 12 hours

Dispense with an accurate

milliliter measuring device

Strawberry Banana Flavored

**Rx only 16 fl oz. (480 mL)**

NDC 23594-101-05

**Karbinal<sup>®</sup> ER**  
(carbinoxamine maleate)

**Extended-release  
Oral Suspension**

**4 mg/5 mL**

Shake Well Before Use

Dose every 12 hours

Dispense with an accurate milliliter  
measuring device

Strawberry Banana Flavored

**Rx only**

**16 fl oz. (480 mL)**

**Usual Dosage:** See prescribing  
information.

**Pharmacist:** Dispense in a  
tight, light-resistant container  
with child-resistant closure.

**Warning:** Keep this and all  
medications out of the reach  
of children.

Store at 25°C (77°F); excursions  
permitted from 15° to 30°C (59°  
to 86°F) [see USP Controlled  
Room Temperature].

Manufactured By:  
**Tris Pharma, Inc.**  
Monmouth Junction, NJ 08852

for:



**Aytu Therapeutics, LLC**  
373 Inverness Parkway, Suite 206  
Englewood, CO 80112  
www.aytubio.com

LB8594 Rev. 01 06/2021



LB8594 Rev. 01 06/2021  
GTIN 00323594101053

Manufactured by:

Tris Pharma, Inc.

Monmouth Junction, NJ 08852

Distributed by:



Aytu Therapeutics

Englewood, CO 80112

www.aytubio.com

LB8594

Rev 01

06/2021

**KARBINAL ER**

carbinoxamine maleate suspension, extended release

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:23594-101
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CARBINOXAMINE MALEATE</b> (UNII: 02O55696WH) (CARBINOXAMINE - UNII:982A7M02H5)	CARBINOXAMINE MALEATE	4 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SODIUM METABISULFITE</b> (UNII: 4VON5FNS3C)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM POLYSTYRENE SULFONATE</b> (UNII: 1699G8679Z)	
<b>TRACETIN</b> (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	STRAWBERRY (Strawberry Banana)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23594-101-05	480 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/03/2014	
2	NDC:23594-101-01	2 in 1 CARTON	01/03/2014	
2		30 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA022556	01/03/2014	

**Labeler** - Aytu Therapeutics, LLC (117244733)

Revised: 1/2025

Aytu Therapeutics, LLC