

**NIGHT-TIME CHERRY MULTI-SYMPTOM COLD/FLU RELIEF- acetaminophen,  
dextromethorphan hydrobromide, doxylamine succinate liquid**

**Chain Drug Consortium, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Active ingredients**

**Drug Facts**

**Active ingredients (in each 30 mL dose cup)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine Succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Keep out of reach of children.**

**Uses**

temporarily relieves cold/flu symptoms:

- sore throat
- headache
- minor aches and pain
- fever
- runny nose and sneezing
- cough due to minor sore throat and bronchial irritation

**Warnings**

**Failure to follow these warnings could result in serious consequences**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take \* more than 4 doses in 24 hours, which is the maximum daily amount \*with other drugs containing acetaminophen \*3 or more alcoholic drinks every day while using this product.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- to make a child sleepy

#### **Ask a doctor before use if you have**

- a sodium restricted diet
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

#### **Ask a doctor or pharmacist before use**

- if you are taking sedatives or tranquilizers
- if you are taking the blood thinning drug warfarin

#### **When using this product**

- **do not use more than directed**
- avoid alcoholic drinks
- excitability may occur, especially in children
- marked drowsiness may occur
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

#### **Stop use and ask a doctor if**

- redness or swelling is present
- symptoms do not get better within 7 days or are accompanied by a fever
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough lasts more than 7 days, comes back, or occurs with fever, rash, headache that lasts
- These could be signs of a serious condition.

#### **If pregnant or breast-feeding**

ask a health professional before use.

#### **Overdose warning**

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- take only as recommended-see Overdose warnings
- Use dose cup or tablespoon (TBSP)

- do not exceed 4 doses per 24 hours
- If taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours

adults & children 12 years and over	30 mL (2 TBS <sup>P</sup> ) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

**Other information**

- **each 30 mL dose cup contains:**
- sodium 45 mg
- store at room temperature

**Inactive ingredients**

citric acid, FD and C Blue No. 1, FD and C Rde No. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose

**Questions?**

Call weekdays

**1-877-798-5944**

**Product Label**

**NDC 68016-146-01**

**COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® NYQUIL®**

**Premier Value®**

**Cherry**

**Night-Time**

**MULTI-SYMP TOM COLD/FLU RELIEF**

Acetaminophen.....Pain Reliever/Fever Reducer  
 Dextromethorphan HBr.....Cough Suppressant  
 Doxylamine Succinate .....Antihistamine

12 FL OZ (354 mL)

INDEPENDENTLY TESTED SATISFACTION GUARANTEED PV

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

\*This product is not manufactured or distributed by Procter and Gamble, owner of the registered trademark Vicks® NyQuil®

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

DISTRIBUTED BY  
 CHAN DRUG CONSORTIUM  
 3301 NW BOCO RATON BLVD  
 SUITE 101, BOCA RATON FL 33431  
 LR-026

**Drug Facts (continued)**  
**Stop use and ask a doctor if** • redness or swelling is present  
 • symptoms do not get better within 7 days or are accompanied by a fever  
 • fever gets worse or lasts more than 3 days • new symptoms occur  
 • cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.  
**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**  
**Overdose warning:** Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions** • take only as recommended—see Overdose warning  
 • use dose cup or tablespoon (TBS) • do not exceed 4 doses per 24 hours  
 • if taking Night Time at night and Day Time during the day, limit total to 4 doses per 24 hours.  

adults & children 12 years and over	30 mL (2 TBS) every 6 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

  
**Other information** • each 30 mL dose cup contains: sodium 45 mg  
 • store at room temperature  
**Inactive ingredients:** citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium citrate, sucrose  
**Questions?** Call weekdays 1-877-798-5944  
 \*This product is not manufactured or distributed by Procter & Gamble owner of the registered trademark Vicks® NyQuil®  
 DISTRIBUTED BY: CHAN DRUG CONSORTIUM  
 3301 NW BOCO RATON BLVD  
 SUITE 101, BOCA RATON, FL 33431  
 LR-026  
 If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

**DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN**  
 Failure to follow these warnings could result in serious consequences

**Drug Facts**  
**Active ingredients (in each 30 mL dose cup)** Purpose  
 Acetaminophen 650 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 30 mg.....Cough suppressant  
 Doxylamine Succinate 12.5 mg.....Antihistamine  
**Uses** temporarily relieves cold/flu symptoms: • sore throat • headache  
 • minor aches and pain • fever • runny nose and sneezing  
 • cough due to minor throat and bronchial irritation  
**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: • more than 4 doses in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.  
**Sore throat warning:** If sore throat is severe, persists for more than two days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleepy  
**Ask a doctor before use if you have** • a sodium restricted diet  
 • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema  
 • trouble urinating due to enlarged prostate gland.  
**Ask a doctor or pharmacist before use** • if you are taking sedatives or tranquilizers • if you are taking the blood thinning drug warfarin  
**When using this product** • do not use more than directed  
 • avoid alcoholic drinks • excitability may occur, especially in children  
 • marked drowsiness may occur  
 • be careful when driving a motor vehicle or operating machinery  
 • alcohol, sedatives, and tranquilizers may increase drowsiness

## NIGHT-TIME CHERRY MULTI-SYMP TOM COLD/FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-146
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	

<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCROSE</b> (UNII: C151H8M554)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY (Cherry)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-146-00	236 mL in 1 BOTTLE		
2	NDC:68016-146-01	354 mL in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/01/2012	

**Labeler** - Chain Drug Consortium, LLC (101668460)

**Registrant** - Aptapharma Inc. (790523323)

### Establishment

Name	Address	ID/FEI	Business Operations
Aptapharma Inc.		790523323	manufacture(68016-146)