

**ALL DAY PAIN RELIEF- naproxen sodium tablet, film coated  
WALMART INC.**

-----  
**Equate 44-604**

***Active ingredient (in each caplet)***

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - the common cold
  - toothache
  - menstrual cramps
  - backache
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- blisters
- shock
- rash
- hives
- asthma (wheezing)
- skin reddening
- facial swelling

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin,

ibuprofen, naproxen, or others]

- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - leg swelling
  - chest pain
  - slurred speech
  - trouble breathing
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

## **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- adults and children 12 years and older
  - take 1 caplet every 8 to 12 hours while symptoms last
  - for the first dose you may take 2 caplets within the first hour
  - do not exceed 2 caplets in any 8- to 12-hour period
  - do not exceed 3 caplets in a 24-hour period
- children under 12 years: ask a doctor

## ***Other information***

- **each caplet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
- see end flap for expiration date and lot number

## ***Inactive ingredients***

croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

## ***Questions or comments?***

**1-888-287-1915**

## ***Principal display panel***

**equate™**

NDC 79903-005-10

Compare to Aleve® Caplets active ingredient\*\*

**All Day**

**Pain Relief**

Naproxen Sodium

Tablets, 220 mg

Pain reliever/Fever reducer (**NSAID**)

**12 HOURS**

Actual Size

**220**

mg

EACH

**40**

CAPLETS†

†Capsule-Shaped Tablets

Satisfaction guaranteed – Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call **1-888-287-1915**.

**DISTRIBUTED BY: Walmart Inc.,  
Bentonville, AR 72716**

PRODUCT OF CHINA AND INDIA

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve® Caplets.

50844 REV1221A60410



### Equate 44-604

## ALL DAY PAIN RELIEF

naproxen sodium tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-005
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NAPROXEN SODIUM</b> (UNII: 9TN8753A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FD&amp;C BLUE NO. 2 ALUMINUM LAKE</b> (UNII: 4AQJ3LG584)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	44;604
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-005-20	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	
2	NDC:79903-005-10	1 in 1 CARTON	10/22/2020	
2		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:79903-005-75	1 in 1 CARTON	10/22/2020	
3		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:79903-005-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	
5	NDC:79903-005-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	
6	NDC:79903-005-25	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	
7	NDC:79903-005-90	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	
8	NDC:79903-005-27	270 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	
9	NDC:79903-005-99	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/22/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204872	10/22/2020	

**Labeler** - WALMART INC. (051957769)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(79903-005)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(79903-005) , pack(79903-005)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(79903-005)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(79903-005)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(79903-005)

Revised: 9/2024

WALMART INC.